


Gold- and Indium-Catalyzed Synthesis of 3- and 6-Sulfonylindoles from *ortho*-Alkynyl-*N*-sulfonylanilines

Itaru Nakamura,* Uichiro Yamagishi, Dschun Song, Sayaka Konta, and Yoshinori Yamamoto



Sulfonylindoles are found in a wide variety of biologically active compounds, such as L-737126^[1] and RO4368554.^[2] However, it is difficult to synthesize sulfonylindoles directly from the corresponding unsubstituted indoles by electrophilic substitution because the electrophilicity of sulfonyl groups is much lower than that of acyl groups and halogens.^[1] Although Yadav et al. recently reported that the indium-catalyzed reaction of indoles with sulfonyl chlorides leads to 3-sulfonylindoles,^[3,4] the development of an efficient and robust method to synthesize sulfonylindoles, which could lead to the discovery of new bioactive compounds,^[5] is still a challenge in organic synthesis.

Several groups, including ourselves, have recently reported that the reactions of *ortho*-alkynylanilines,^[6] *ortho*-alkynylphenyl ethers,^[6a,b,7] and *ortho*-alkynylphenyl sulfides^[8] with a migrating group (E) on the heteroatom (Y) give the corresponding 2,3-disubstituted indoles, benzofurans, and benzothiophenes, respectively, in the presence of transition-metal catalysts [Eq. (1)].




Acyl, allyl, α -alkoxyalkyl, and methoxyphenylmethyl (MPM) groups can be employed as migrating groups. In this article, we report that the reactions of the *ortho*-alkynyl-*N*-sulfonylanilines **1b-o** give the corresponding 3-sulfonylindoles **2b-o** in good to excellent yields in the presence of a catalytic amount of AuBr₃ [Eq. (2)].

In addition, with InBr₃ as catalyst the cyclization of 2-alkynyl-6-methoxy-*N*-sulfonylanilines **1p-x** proceeds by an unprecedented 1,7-migration of the sulfonyl group to produce the 6-sulfonylindoles **3** as the major product in good to high yields [Eq. (3)].



First, the catalytic activity of transition-metal compounds was tested with *N*-methyl-2-(1-pentynyl)-*N*-tosylaniline (**1a**) as substrate [Eq. (4)]. AuBr₃ showed the highest selectivity among the transition-metal complexes examined. Thus, the reaction of **1a** in the presence of 10 mol % AuBr₃ in toluene at 80 °C for one hour yielded *N*-methyl-2-propyl-3-tosylindole (**2a**) in 60% yield along with *N*-methyl-2-propyl-6-tosylindole (**3a**)^[9] and *N*-methyl-2-propyl-4-tosylindole (**4a**), which are derived from an unprecedented sulfonyl migration to the benzene ring of the indole skeleton, in 17% and 12% yields, respectively. In our previous metal-catalyzed migration of *ortho*-alkynylacetanilides,^[6d] *ortho*-alkynylphenyl acetals,^[7f] and *ortho*-alkynylphenyl sulfides,^[8] we did not observe substitution of the benzene ring by the migrating groups. The reaction of **1a** in the presence of PtCl₂ or PdCl₂ produced a 1:1 mixture of **2a** and **3a**, while the use of indium complexes led to a mixture of **2a**, **3a**, and **4a** (see the Supporting Information).

Next, we changed the tosyl group for a mesyl group. The results of the reaction of *ortho*-alkynyl-*N*-mesylanilines **1b-l**




[*] Dr. I. Nakamura, U. Yamagishi, Dr. D. Song, S. Konta, Prof. Dr. Y. Yamamoto
Department of Chemistry
Graduate School of Science
Tohoku University
Sendai 980-8578 (Japan)
Fax: (+81) 22-795-6784
E-mail: itaru-n@mail.tains.tohoku.ac.jp

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

with AuBr₃ as catalyst are summarized in Table 1. The reaction of *N*-mesyl-*N*-methyl-2-(1-pentynyl)aniline (**1b**) in the presence of 10 mol % of AuBr₃ in toluene at 80 °C for one

Table 1: AuBr₃-catalyzed cyclization of **1b-o**.^[a]




Entry	1	R ¹	R ²	R ³	Yield [%] 2 ^[b]	3 and 4 ^[c]
1	1b	<i>n</i> Pr	Me	Me	95	–
2	1c	cyclohexyl	Me	Me	62	25
3	1d	<i>t</i> Bu	Me	Me	38 ^[d]	10
4	1e	Ph	Me	Me	92	2
5	1f	<i>p</i> -tolyl	Me	Me	87	–
6	1g	<i>p</i> -anisyl	Me	Me	81	5
7	1h	<i>p</i> -F ₃ CC ₆ H ₄	Me	Me	51	20
8	1i	H	Me	Me	71	–
9	1j	CO ₂ Et	Me	Me	decomp ^[e]	–
10	1k	<i>n</i> Pr	Me	Bn	44	–
11	1l	<i>n</i> Pr	Me	<i>i</i> Pr	60	–
12	1m	<i>n</i> Pr	Ph	Me	52	22
13	1n	<i>n</i> Pr	<i>p</i> -anisyl	Me	85	7
14	1o	<i>n</i> Pr	<i>p</i> -AcC ₆ H ₄	Me	80	5

[a] The reaction of **1b-o** (0.25 mmol) was carried out in the presence of 10 mol % of AuBr₃ in toluene at 80 °C for 1 h. [b] Yield of isolated product. [c] Yield of an inseparable mixture of **3** and **4** determined by GC. [d] Substrate **1d** was recovered in 29% yield. [e] A mixture of unidentified products was obtained.

hour gave 3-mesyl-1-methyl-2-propylindole (**2b**) in 95% yield (Table 1, entry 1). No other regioisomers derived from aromatic substitution were obtained. The reaction of **1b** in the presence of AuCl₃ or PtCl₂ gave **2b** in 85% and 93% yield, respectively. Substrates **1c** and **1d**, where R¹ is a cyclohexyl or *tert*-butyl group, were converted into **2c** and **2d** in 62% and 38% yields, respectively (Table 1, entries 2 and 3). The reaction of the 2-(arylethynyl)-*N*-tosylanilines **1f** and **1g**, where R¹ is an electron-rich aromatic ring, gave **2f** and **2g** in good yields, while the reaction with **1h**, which is substituted with an electron-deficient aromatic ring, produced **2h** in lower yield along with significant amounts of the other regioisomers (Table 1, entries 5–7). The terminal alkyne **1i** was converted into the 3-mesyl-substituted indole **2i** in 71% yield, while the reaction of the ynoate **1j** led to a mixture of unidentified products (Table 1, entries 8 and 9, respectively). The reaction of *N*-benzyl-*N*-sulfonylaniline (**1k**) and *N*-isopropyl-*N*-sulfonylaniline (**1l**) afforded the corresponding indoles **2k** and **2l** in moderate yields (Table 1, entries 10 and 11, respectively). The reaction of the arylsulfonamides **1m-o** gave **2m-o** along with considerable amounts of the other regioisomers (Table 1, entries 12–14).

Many attempts to synthesize 4- or 6-sulfonylindoles as the major products by varying the reactions conditions (solvent, ligands, additives, and temperature) only led to a mixture of **2**, **3**, and/or **4**. We then modified the substrate by adding functional groups. Among the substrates prepared, 2-alkynyl-6-methoxysulfonamides **1p-x**, which have a methoxy group at the 6-position of the aniline moiety, were mainly converted into the corresponding 6-sulfonylindoles **3p-x** in the presence of catalytic amounts of InBr₃ [Eq. (3)]. The results are summarized in Table 2. The reaction of **1p** in the presence of 5 mol % of InBr₃ in toluene at 80 °C for two hours gave an 87:13 mixture of **3p** and **2p** in 95% combined yield (Table 2,

Table 2: InBr₃-catalyzed cyclization of **1p-x**.^[a]




Entry	1	R ¹	R ²	Yield [%] ^[b]	3/2 ^[c]
1	1p	<i>n</i> Pr	Me	95	87:13
2	1q	<i>n</i> Pr	MeO	99	84:16
3	1r	<i>n</i> Pr	H	98	78:22
4	1s	<i>n</i> Pr	NO ₂	99	66:34
5	1t	cyclohexyl	Me	97	87:13
6	1u	Ph	Me	99	90:10
7	1v	<i>p</i> -MeOC ₆ H ₄	Me	98	85:15
8	1w	<i>p</i> -CF ₃ C ₆ H ₄	Me	88	83:17
9	1x	H	Me	73	54:46


[a] The reactions of **1p-x** were carried out in the presence of 5 mol % of InBr₃ in toluene at 80 °C. [b] Yield of the isolated mixture of **2** and **3**. [c] The ratio was determined by ¹H NMR spectroscopy.

entry 1). The reaction of **1p** in the presence of AuBr₃ or PdBr₂ gave **3p** and **2p** with lower regioselectivities (Supporting Information). The reaction of **1q**, which has an electron-donating methoxy group on the arylsulfonamide moiety, produced the 6-sulfonylindole **3q** with higher regioselectivity than that of **1s**, which contains an electron-withdrawing nitro group (Table 2, entries 2 and 4, respectively). Substrate **1t**, which has a cyclohexyl group at R¹, was converted into **3t** with high regioselectivity (Table 2, entry 5). The ratio of **3** to **2** was not affected by the electronic properties of the R¹ substituent (Table 2, entries 6–8), although it was lower when terminal alkyne **1x** was employed as substrate (Table 2, entry 9). The reaction of the *N*-mesylaniline **1y** afforded a mixture of unidentified products, and **5a-c**, which have a methoxy group at the 3-, 4-, and 5-position, respectively, reacted sluggishly to give inseparable mixtures of unidentified products.

To find out if the migration of the sulfonyl group occurs in an intramolecular or intermolecular fashion, we performed crossover experiments [Eqs. (5) and (6)]. The reaction of a 1:1 mixture of **1e** and **1n** in the presence of a catalytic amount of




AuBr₃ gave the corresponding products **2e** and **2n** in 66 % and 65 % yields, respectively [Eq. (5)]; the crossover products were not detected by GC-MS or NMR spectroscopy.




Furthermore, the InBr₃-catalyzed reaction of a 1:1 mixture of **1q** and **1w** afforded the products **3q** and **2q**, derived from **1q**, and **3w** and **2w**, derived from **1w** [Eq. (6)]; again, no crossover products were obtained. These results clearly indicate that the present reaction proceeds in an intramolecular manner.


The isolated products **2a**, **3a**, and **4a** remain unchanged in the presence of InBr₃ or AuBr₃ in toluene at 80 °C for two




hours, thereby suggesting that interconversion between the products does not take place under the reaction conditions [Eq. (7)].



Mixing indoles **6a** and **6b** with tosyl chloride in the presence of AuBr₃ or InBr₃ did not give the corresponding sulfonylindoles **2**, **3**, and **4** [Eq. (8)].^[3] Accordingly, it is unlikely that electrophilic substitution of the indole with tosyl halides occurs under these reaction conditions.



The above experimental results led us to propose the mechanism for the cyclization of **1** shown in Scheme 1. The Lewis-acidic transition metal coordinates to the triple bond of **1** to form the π -complex **7**. Nucleophilic attack of the nitrogen atom to the alkynyl moiety then leads to the cyclized intermediate **8**. For the gold-catalyzed reaction of **1a–o**, the sulfonyl group intramolecularly migrates to the 3-position of



Scheme 1. Proposed mechanism for the catalytic formation of **2** and **3** from **1**.

the indole skeleton (cycle **A**),^[10] and elimination of AuBr₃ from **9** then gives the 3-sulfonylindole **2**. In the indium-catalyzed reaction of substrates **1p–x**, unprecedented consecutive 1,7-sulfonyl and 1,5-proton shifts take place instead (cycle **B**).^[11] Elimination of InBr₃ from the resulting intermediate **11** then gives the 6-sulfonylindoles **3**. An interaction between the benzene ring on the sulfonyl group and the indium catalyst might play a crucial role in selectively producing 6-sulfonylindoles **3**, since the InBr₃-catalyzed reaction of *N*-mesylaniline **1y** gives a complex mixture of unidentified products.

The present reaction proceeds by formal addition of a nitrogen–sulfur bond to a triple bond, a so-called amino-sulfonylation.^[12] It is therefore likely that this method could be applicable in an efficient and environmentally benign synthesis of a wide variety of 3- and 6-sulfonylindoles.^[13]

Experimental Section

AuBr₃-catalyzed cyclization of 1b: Toluene (0.5 mL) was added to a mixture of AuBr₃ (0.025 mmol) and **1b** (0.25 mmol) in a pressure vial under argon. After stirring at 80 °C for 1 h, the reaction mixture was filtered through a short SiO₂ pad. The crude product was purified by silica gel column chromatography with hexane/ethyl acetate as eluent to afford **2b** (95 %).

InBr₃-catalyzed cyclization of 1p: Toluene (1 mL) was added to a mixture of InBr₃ (0.0125 mmol) and **1p** (0.25 mmol). After stirring at 80 °C for 2 h, the reaction mixture was purified by Florisil column chromatography with hexane/ethyl acetate as eluent to afford **3p** (83 %) and **2p** (12 %). Further purification was performed by gel permeation chromatography.

Received: September 30, 2006

Revised: December 16, 2006

Keywords: cyclization · fused-ring systems · gold · indium · synthetic methods

- [1] a) T. M. Williams, T. M. Ciccarone, S. C. MacTough, C. S. Rooney, S. K. Balani, J. H. Condra, E. A. Emini, M. E. Goldman, W. J. Greenlee, L. R. Kauffman, J. A. O'Brien, V. V. Sardana, W. A. Schleif, A. D. Theoharides, P. S. Anderson, *J. Med. Chem.* **1993**, *36*, 1291–1294; b) R. Silvestri, G. De Martino, G. La Regina, M. Artico, S. Massa, L. Vargiu, M. Mura, A. G. Loi, T. Marceddu, P. La Colla, *J. Med. Chem.* **2003**, *46*, 2482–2493.

- [2] C. K. Lieben, A. Blokland, A. Sik, E. Sung, P. van Nieuwenhuizen, R. Schreiber, *Neuropsychopharmacology* **2005**, *30*, 2169–2179.
- [3] J. S. Yadav, B. V. S. Reddy, A. D. Krishna, T. Swamy, *Tetrahedron Lett.* **2003**, *44*, 6055–6058.
- [4] We have repeated this reaction several times but so far we have not been able to obtain 3-sulfonylindoles in good yields—the reaction of indole and tosyl chloride under the conditions mentioned in reference [3] gave 3-tosylindole in only 16 % yield in our hands.
- [5] a) R. Ragno, M. Artico, G. De Martino, G. La Regina, A. Coluccia, A. Di Pasquali, R. Silvestri, *J. Med. Chem.* **2005**, *48*, 213–223; b) R. Bernotas, S. Lenicek, S. Antane, G. M. Zhang, D. Smith, J. Coupet, B. Harrison, L. E. Schechter, *Bioorg. Med. Chem. Lett.* **2004**, *14*, 5499–5502.
- [6] Migration of allyl groups: a) S. Cacchi, G. Fabrizi, P. Pace, *J. Org. Chem.* **1998**, *63*, 1001–1011; b) A. Fürstner, P. W. Davies, *J. Am. Chem. Soc.* **2005**, *127*, 15024–15025. Propargyl groups: c) S. Cacchi, G. Fabrizi, L. Moro, *Tetrahedron Lett.* **1998**, *39*, 5101–5104. Acyl groups: d) T. Shimada, I. Nakamura, Y. Yamamoto, *J. Am. Chem. Soc.* **2004**, *126*, 10546–10547.
- [7] Allyl groups: a) A. Arcadi, S. Cacchi, M. D. Rosario, G. Fabrizi, F. Marinelli, *J. Org. Chem.* **1996**, *61*, 9280–9288; b) S. Cacchi, G. Fabrizi, L. Moro, *Synlett* **1998**, 741–745; c) N. Monterio, G. Balme, *Synlett* **1998**, 746–747; d) A. Fürstner, H. Szillat, F. Stelzer, *J. Am. Chem. Soc.* **2000**, *122*, 6785–6786; e) A. Fürstner, F. Stelzer, H. Szillat, *J. Am. Chem. Soc.* **2001**, *123*, 11863–11869. α -Alkoxyalkyl groups: f) I. Nakamura, Y. Mizushima, Y. Yamamoto, *J. Am. Chem. Soc.* **2005**, *127*, 15022–15023 and reference [6b].
- [8] I. Nakamura, T. Sato, Y. Yamamoto, *Angew. Chem.* **2006**, *118*, 4585–4587; *Angew. Chem. Int. Ed.* **2006**, *45*, 4473–4475.
- [9] CCDC 622538 (**3a**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [10] For migration of sulfonyl groups, see: a) S. Roy, I. Das, K. Bhanuprakash, B. D. Gupta, *Tetrahedron* **1994**, *50*, 1847–1858; b) J. J. González, P. M. Nieto, P. Prados, A. M. Echavarren, J. de Mendoza, *J. Org. Chem.* **1995**, *60*, 7419–7423; c) K. Oda, R. Hiratsuka, M. Machida, *Heterocycles* **1996**, *43*, 463–470; d) Y. Horino, M. Kimura, Y. Wakamiya, T. Okajima, Y. Tamaru, *Angew. Chem.* **1999**, *111*, 123–126; *Angew. Chem. Int. Ed.* **1999**, *38*, 121–124; e) Y. Horino, M. Kimura, M. Naito, S. Tanaka, Y. Tamaru, *Tetrahedron Lett.* **2000**, *41*, 3427–3431.
- [11] K. K. Park, J. J. Lee, J. Ryu, *Tetrahedron* **2003**, *59*, 7651–7659.
- [12] Z. Liu, R. C. Larock, *J. Am. Chem. Soc.* **2005**, *127*, 13112–13113.
- [13] For a synthesis of 6-sulfonylindoles, see: J. Caixach, R. Capell, C. Galvez, A. Gonzalez, N. Roca, *J. Heterocycl. Chem.* **1979**, *16*, 1631–1635.