


# A Highly Enantioselective Intramolecular Michael Reaction Catalyzed by N-Heterocyclic Carbenes\*\*


Eric M. Phillips, Manabu Wadamoto, Audrey Chan, and Karl A. Scheidt\*

The conjugate addition of enolate or enol nucleophiles to activated double bonds (the Michael reaction) is an efficient method for the synthesis of 1,5-dicarbonyl compounds.<sup>[1]</sup> The potential for the direct formation of a new carbon–carbon bond with control of up to three new stereogenic centers has driven continued and significant development of this reaction. Two central strategies are the use of metalloenolates and the addition of latent nucleophiles such as enol silanes in combination with a Lewis acid.<sup>[2]</sup> Studies by the research groups of Yamaguchi,<sup>[3]</sup> MacMillan,<sup>[4]</sup> Jørgensen,<sup>[5]</sup> and List<sup>[6]</sup> have demonstrated that secondary amines catalyze Michael reactions by the generation of activated unsaturated electrophiles through iminium ions. The corresponding strategy to the iminium approach is the catalytic generation of an enolate or enol nucleophile.<sup>[7,8]</sup> Herein we report that N-heterocyclic carbenes (NHCs) are highly selective catalysts for the intramolecular Michael reaction of substrates **1** to afford dicarbonyl compounds **2** after the addition of an exogenous nucleophile [Eq. (1)].



The Michael reaction traditionally relies on the stoichiometric generation of an enolate or enol. In our recent studies that involved NHCs and carbonyl compounds,<sup>[9]</sup> the opportunity to catalyze the formation of enol intermediates became apparent.<sup>[10]</sup> We anticipated that intramolecular Michael reactions catalyzed by NHCs could be achieved by using a conjugate acceptor with the general structure of **1**. The

proposed pathway for this process involves the addition of the NHC to an  $\alpha,\beta$ -unsaturated aldehyde to afford the extended diene intermediate **I** (Scheme 1).<sup>[11]</sup> The key enol nucleophile is revealed by  $\beta$ -protonation of **I** followed by a Michael addition to generate the enol **III**. An intramolecular acylation event releases the NHC catalyst to afford the bicyclic acylated enol **IV** that readily opens to give products such as **2** on exposure to mild nucleophiles.<sup>[12]</sup>



Scheme 1. Proposed catalytic pathway.

The search for optimal conditions for this process started with the parent enal **1a**, imidazolium salt **A**, and typical reaction conditions for NHC catalysis (Table 1, entry 1, Mes = 2,4,6-trimethylphenyl). Unfortunately, only a small amount of the desired product **3** was observed. When the base was changed from DBU to *i*Pr<sub>2</sub>EtN, we were pleased to observe an increased yield of **3** with greater than 20:1 selectivity favoring the *cis* diastereomer (entry 3).<sup>[13]</sup> The use of the achiral triazolium salt **B** in toluene with THF as a cosolvent to induce homogeneity of the reaction provided good yields (entry 4). Solvents that are not Lewis basic, such as toluene or CH<sub>2</sub>Cl<sub>2</sub>, were crucial to promote the  $\beta$ -protonation process with all the azolium precatalysts. This protonation step in turn generates the desired enol which participates in the reaction. When the concentration of **1a** in the reaction was decreased, the intramolecular manifold was favored with a resulting increase in yield (entry 9).


Importantly, NHCs derived from the triazolium salts **C** and **D** were found to be the most efficient catalysts and provided a platform to control the stereochemical outcome

[\*] E. M. Phillips, Dr. M. Wadamoto, A. Chan, Prof. K. A. Scheidt  
Department of Chemistry  
Northwestern University  
2145 Sheridan Rd, Evanston, IL 60208 (USA)  
Fax: (+1) 847-467-2184  
E-mail: scheidt@northwestern.edu

[\*\*] Research support was generously provided by the NIH/NIGMS (RO1 GM73072), Abbott Laboratories, Amgen, 3M, and Boehringer-Ingelheim. A.C. is the recipient of a Dow Chemical Company Fellowship. We thank T. Reynolds and C. Stern (NU) for assistance with X-ray crystallography. Funding for the NU Analytical Services Laboratory has been furnished in part by the NSF (CHE-9871268).


Supporting information for this article (detailed experimental procedures and full characterization of new compounds) is available on the WWW under <http://www.angewandte.org> or from the author.

**Table 1:** Optimization of the Michael reaction.



Entry	Catalyst	Conditions <sup>[a]</sup>	Yield <sup>[b]</sup>	ee [%] <sup>[c]</sup>
1	<b>A</b>	DBU, THF	6	–
2	<b>A</b>	KN(SiMe <sub>3</sub> ) <sub>2</sub> , THF <sup>[d]</sup>	11	–
3	<b>A</b>	<i>i</i> Pr <sub>2</sub> EtN, THF, 45 °C	50	–
4	<b>B</b>	<i>i</i> Pr <sub>2</sub> EtN, toluene/THF <sup>[e]</sup>	61	–
5	<b>C</b>	<i>i</i> Pr <sub>2</sub> EtN, toluene/THF <sup>[e,f]</sup>	61	93
6	<b>C</b>	Et <sub>3</sub> N (0.1 M), toluene/THF <sup>[e]</sup>	62	93
7	<b>C</b>	<i>i</i> Pr <sub>2</sub> EtN (0.1 M), CH <sub>2</sub> Cl <sub>2</sub> , –20 °C	49	93
8	<b>D</b>	<i>i</i> Pr <sub>2</sub> EtN (0.1 M), toluene/THF <sup>[e]</sup>	53	99
9	<b>D</b>	<i>i</i> Pr <sub>2</sub> EtN (0.05 M), toluene/THF <sup>[e]</sup>	66	99
10	<b>D</b>	<i>i</i> Pr <sub>2</sub> EtN (0.05 M), CH <sub>2</sub> Cl <sub>2</sub>	68	99
11	<b>D</b> <sup>[g]</sup>	<i>i</i> Pr <sub>2</sub> EtN (0.05 M), CH <sub>2</sub> Cl <sub>2</sub>	68	99

[a] Base (20 mol%), **1a** (0.2 M) at 23 °C unless otherwise noted. DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene. [b] Yield of isolated product. [c] Determined by HPLC (Chiracel AD-H). Absolute and relative configuration of **3** assigned by X-ray crystallography.<sup>[16]</sup> See the Supporting Information for details. [d] Carbene generated prior to addition of substrate. [e] 10:1 toluene/THF. [f] Base (1.2 equiv). [g] **D** (10 mol%).




for this process. Accordingly, the use of the phenylalanine-derived salt **C** afforded good yields of **3** with 93% *ee* (entries 5–7). When the structure of the catalyst was tuned to that from the salt **D**, which was derived from amino indanol and was first disclosed by Bode and co-workers,<sup>[14]</sup> we observed excellent levels of enantioselectivity with catalyst loadings of 10 mol % (99% *ee*, entry 11). It is important to note that the inclusion of *N*-mesityl substitution on the azolium salts was required for any reasonable conversion into the desired product.

Once the triazolium salt **D** had been identified as the most selective precatalyst, we surveyed potential substrates for this intramolecular process (Table 2). The use of methanol to quench the reaction avoids the propensity for several of the bicyclic products to undergo hydrolysis when purified on silica gel. The optimized reaction conditions allowed both electron-withdrawing and -donating groups on the enone (entries 1–3), and additionally, electron-withdrawing and -donating substituents could be placed on the aromatic tether (entries 6 and 7). This intramolecular reaction was not restricted to the use of aromatic substituents. The  $\alpha,\beta$ -unsaturated methyl ketone **1d** provided a moderate yield of the cyclopentane product with excellent enantioselectivity (entry 4). The bisaldehyde **1e** underwent an interesting desymmetrization reaction in which one aldehyde became the nucleophile when exposed to an NHC while the other unsaturated moiety became the conjugate acceptor (entry 5). The cyclization of the aliphatic substrate **13** (entry 8) proceeded in good yield after ten hours with a catalyst loading of 20 mol %. When the tether length was

increased to access six-membered rings, cyclohexene products were afforded but with reduced enantioselectivity and yield (62% *ee*, 52%; entry 9).<sup>[15]</sup> Interestingly, product **16** did not open after the addition of methanol unlike the cyclopentane compound.

The efficient formation of methyl esters by the simple addition of methanol to the reaction in Table 2 indicated that bicyclic intermediates such as **3** are good acylating agents. Accordingly, substituted cyclopentyl amides, such as primary amide **17** and secondary amide **18**, could be accessed in good yield directly by the addition of the corresponding primary and secondary amines to the reaction mixture after the starting material **1a** had been consumed (Scheme 2).



**Scheme 2.** Amide formation from the acylated enol products.

In summary, a highly diastereo- and enantioselective intramolecular Michael reaction catalyzed by *N*-heterocyclic carbenes has been developed. The addition of the carbene catalyst to an  $\alpha,\beta$ -unsaturated aldehyde, followed by subsequent  $\beta$ -protonation generated a reactive enol intermediate that underwent addition to a pendant conjugate acceptor. Aryl and alkyl substituents are suitable for the reaction, and high enantioselectivity is achieved when chiral enantiopure triazolium salts are used. The turnover of the catalyst is facilitated by the generation of a cyclic *O*-acylated enol which is intercepted by alcohols and amines to provide different esters and amides. Further studies to generate nucleophiles by using *N*-heterocyclic carbenes are ongoing and will be reported in due course.

## Experimental Section

The azolium salt **D** (4.2 mg, 0.01 mmol) and the corresponding enal (0.1 mmol) were added to a flame-dried round-bottom flask (10 mL) containing a magnetic stirring bar. The flask was sealed under a positive pressure of N<sub>2</sub>. The heterogeneous mixture was then diluted with CH<sub>2</sub>Cl<sub>2</sub> (2 mL, 0.05 M). Once the material had dissolved, diisopropylethylamine (2  $\mu$ L, 0.01 mmol) was added through a syringe. The reaction mixture was stirred at 23 °C under N<sub>2</sub> until the enal had been completely consumed (as observed by TLC). Methanol (5 mL) was then added and the reaction mixture stirred at 23 °C under N<sub>2</sub> for 5 h. The reaction mixture was partially concentrated under reduced pressure and the remaining residue was purified by chromatography (silica gel, 5% EtOAc/Hexanes) to afford the pure methyl ester. Analytical data for **4**: IR (film):  $\tilde{\nu}$  = 3024, 2947, 1730, 1685, 1442, 1365 cm<sup>–1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  = 7.95 (d, *J* = 7.3 Hz, 2H), 7.57 (t, *J* = 7.3 Hz, 1H), 7.46 (t, *J* = 7.6 Hz, 2H), 7.27 (d, *J* = 6.4 Hz, 1H), 7.19 (m, 3H), 4.28 (q, *J* = 7.3 Hz, 1H),

Table 2: Substrate Scope.

Entry	Substrate	Product	Yield [%] <sup>[a]</sup>	ee [%] <sup>[b]</sup>
1	<b>1a</b> R = Ph	<b>4</b>	69	99
2	<b>1b</b> R = 4-BrC <sub>6</sub> H <sub>4</sub>	<b>5</b>	62	99
3	<b>1c</b> R = 4-MeC <sub>6</sub> H <sub>4</sub>	<b>6</b>	80	99
4	<b>1d</b>	<b>7</b>	59	99
5	<b>1e</b>	<b>8</b>	68	99
6	<b>9</b>	<b>10</b>	68	99
7	<b>11</b>	<b>12</b>	73	99
8 <sup>[c]</sup>	<b>13</b>	<b>14</b>	66	99
9 <sup>[d]</sup>	<b>15</b>	<b>16</b>	52	62

[a] Yield of isolated product after purification. [b] Determined by HPLC on Chiralcel AD-H or OD-H columns. [c] **D** (20 mol %) with the enantioselectivity determined by GC (BetaDex column). [d] **D** (20 mol %).

3.59 (m, 4H), 3.42 (m, 2H), 3.15 ppm (m, 2H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ = 198.72, 174.72, 144.87, 137.26, 133.32, 128.84, 128.26, 127.53, 127.08, 124.81, 124.35, 51.90, 47.69, 42.75, 40.81, 34.73 ppm; LRMS (ES): calcd for C<sub>18</sub>H<sub>16</sub>O<sub>3</sub> [M]<sup>+</sup>, 294.32; found [M+H]<sup>+</sup>, 295.5; [α]<sub>D</sub> = −16.6 (CH<sub>2</sub>Cl<sub>2</sub>, c = 1.0, 99.5:0.5 *er*). The enantiomeric ratio was determined by HPLC (Chiralcel AD-H, 15% 2-propanol/hexanes, 1 mL min<sup>−1</sup>, t<sub>r1</sub> = 8.84, t<sub>r2</sub> = 13.88).

Received: December 27, 2006

Revised: February 2, 2007

Published online: March 15, 2007

**Keywords:** acylation · asymmetric catalysis · Michael addition · N-heterocyclic carbenes · synthetic methods

- [1] a) A. Michael, *J. Prakt. Chem.* **1887**, 36, 349–356; b) E. D. Bergmann, D. Ginsburg, R. Pappo, *Org. React.* **1959**, 10, 179–555; c) D. A. Oare, C. A. Heathcock, In *Topics in Stereochemistry*, Vol. 20 (Eds.: E. L. Eliel, S. H. Wilen), Wiley, New York, **1991**, pp. 124–170; d) R. D. Little, M. R. Masjedizadeh, *Org. React.* **1995**, 47, 315–552; e) N. Krause, A. Hoffmann-Röder, *Synthesis* **2001**, 171–196.
- [2] a) K. Narasaka, K. Soai, T. Mukaiyama, *Chem. Lett.* **1974**, 1223–1224; b) T. Mukaiyama, S. Kobayashi, *Org. React.* **1994**,

- 46, 1–103; c) D. A. Evans, K. A. Scheidt, J. N. Johnston, M. C. Willis, *J. Am. Chem. Soc.* **2001**, 123, 4480–4491, and references therein; d) *Comprehensive Asymmetric Catalysis* (Eds.: E. N. Jacobsen, A. Pfaltz, H. Yamamoto), Springer, New York, **1999**, chap. 31.
- [3] a) M. Yamaguchi, T. Shiraishi, M. Hiram, *Angew. Chem.* **1993**, 105, 1243–1245; *Angew. Chem. Int. Ed. Engl.* **1993**, 32, 1176–1178; b) M. Yamaguchi, T. Shiraishi, M. Hiram, *J. Org. Chem.* **1996**, 61, 3520–3530.
- [4] S. P. Brown, N. C. Goodwin, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2003**, 125, 1192–1194.
- [5] a) P. Melchiorre, K. A. Jørgensen, *J. Org. Chem.* **2003**, 68, 4151–4157; b) N. Halland, P. S. Aburel, K. A. Jørgensen, *Angew. Chem.* **2003**, 115, 685–689; *Angew. Chem. Int. Ed.* **2003**, 42, 661–665; c) N. Halland, P. S. Aburel, K. A. Jørgensen, *Angew. Chem.* **2004**, 116, 1292–1297; *Angew. Chem. Int. Ed.* **2004**, 43, 1272–1277.
- [6] a) B. List, *Tetrahedron* **2002**, 58, 5573–5590; b) M. T. Hechavarria Fonseca, B. List, *Angew. Chem.* **2004**, 116, 4048–4050; *Angew. Chem. Int. Ed.* **2004**, 43, 3958–3960; c) For an excellent review of enamine catalysis, see: B. List, *Acc. Chem. Res.* **2004**, 37, 548–557.
- [7] For examples of metal-catalyzed enolate generation, see: a) N. Yoshikawa, Y. M. A. Yamada, J. Das, H. Sasai, M. Shibasaki, *J. Am. Chem. Soc.* **1999**, 121, 4168–4178; b) B. M. Trost, H. Ito, *J. Am. Chem. Soc.* **2000**, 122, 12003–12004; c) D. A. Evans, J. S. Tedrow, J. T. Shaw, C. W. Downey, *J. Am. Chem. Soc.* **2002**, 124, 392–393; d) D. A. Evans, R. J. Thomson, *J. Am. Chem. Soc.* **2005**, 127, 10506–10507.
- [8] For examples of amine-catalyzed enamine generation, see: a) B. List, R. A. Lerner, C. F. Barbas III, *J. Am. Chem. Soc.* **2000**, 122, 2395–2396; b) J. M. Betancort, K. Sakthivel, R. Thayumanavan, C. F. Barbas III, *Tetrahedron Lett.* **2001**, 42, 4441–4444; c) B. List, *Synlett* **2001**, 1675–1686; d) P. I. Dalko, L. Moisan, *Angew. Chem.* **2001**, 113, 3840–3864; *Angew. Chem. Int. Ed.* **2001**, 40, 3726–3748; e) B. List, *Tetrahedron* **2002**, 58, 5573–5590; f) Y. Hayashi, H. Gotoh, T. Tamura, H. Yamaguchi, R. Masui, M. Shoji, *J. Am. Chem. Soc.* **2005**, 127, 16028–16029.
- [9] a) A. E. Mattson, A. R. Bharadwaj, K. A. Scheidt, *J. Am. Chem. Soc.* **2004**, 126, 2314; b) A. R. Bharadwaj, K. A. Scheidt, *Org. Lett.* **2004**, 6, 2465; c) A. E. Mattson, K. A. Scheidt, *Org. Lett.* **2004**, 6, 4363; d) M. C. Myers, A. R. Bharadwaj, B. C. Milgram, K. A. Scheidt, *J. Am. Chem. Soc.* **2005**, 127, 14675; e) A. Chan, K. A. Scheidt, *Org. Lett.* **2005**, 7, 905; f) A. Chan, K. A. Scheidt, *J. Am. Chem. Soc.* **2006**, 128, 4558.
- [10] Enols have also been implemented as intermediates in reactions that combine NHCs and aldehydes; see: a) N. T. Reynolds, T. Rovis, *J. Am. Chem. Soc.* **2005**, 127, 16406–16407; b) M. He, J. R. Struble, J. W. Bode, *J. Am. Chem. Soc.* **2006**, 128, 8418–8420.

- [11] The potential hetero-Diels–Alder pathway is unlikely given the constrained nature of the intramolecular substrates and the high levels of *cis* diastereoselectivity. For a related intermolecular process using triazolium salt **D** as the precatalyst which invokes a hetero-Diels–Alder pathway, see: M. He, G. J. Uc, J. W. Bode, *J. Am. Chem. Soc.* **2006**, *128*, 15088–15089.
- [12] The high diastereoselectivity in this reaction may arise because of a reversible Michael reaction followed by an irreversible lactonization occurring through the *cis* isomer. Investigations of these mechanistic pathways are ongoing.
- [13] For an approach to the *trans* diastereomers of compounds such as **4** using the Hantzsch ester and a chiral secondary amine, see J. W. Yang, M. T. H. Fonseca, B. List, *J. Am. Chem. Soc.* **2005**, *127*, 15036–15037.
- [14] See Ref [10b]. For relevant examples of chiral triazolium salts used in asymmetric catalysis, see: a) D. Enders, K. Breuer, J. Runsink, J. H. Teles, *Helv. Chim. Acta* **1996**, *79*, 1899–1902; b) M. S. Kerr, J. R. de Alaniz, T. Rovis, *J. Am. Chem. Soc.* **2002**, *124*, 10298–10299; c) D. Enders, U. Kallfass, *Angew. Chem.* **2002**, *114*, 1822–1824; *Angew. Chem. Int. Ed.* **2002**, *41*, 1743; d) N. T. Reynolds, J. R. de Alaniz, T. Rovis, *J. Am. Chem. Soc.* **2004**, *126*, 9518; e) D. Enders, T. Balensiefer, *Acc. Chem. Res.* **2004**, *37*, 534; f) J. R. de Alaniz, T. Rovis, *J. Am. Chem. Soc.* **2005**, *127*, 6284; g) D. Enders, O. Niemeier, T. Balensiefer, *Angew. Chem.* **2006**, *118*, 1491; *Angew. Chem. Int. Ed.* **2006**, *45*, 1463.
- [15] The remaining mass balance is primarily the five-membered ring which resulted from the conjugate addition of the homoenolate formed in situ.
- [16] CCDC-631931 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).