



Rh-Catalyzed Transannulation of Pyridotriazoles with Alkynes and Nitriles**


Stepan Chuprakov, Frank W. Hwang, and Vladimir Gevorgyan*

Transition-metal-catalyzed annulations are widely used in the synthesis of heterocyclic compounds.^[1] One of the most efficient methods for the construction of five-membered oxygen-containing heterocycles involves the annulation of diazocarbonyl compounds with alkynes and nitriles. Thus, Davies et al.^[2] and Padwa et al.^[3] have employed this method^[4] for the synthesis of furans (X = CH), and Helquist et al.^[5] for the preparation of oxazoles (X = N) [Eq. (1)]. In contrast, analogous transformations of α -imino diazo compounds, which may lead to the formation of pyrrole and imidazole rings, are unknown. Herein we report an efficient, direct, Rh-catalyzed transannulation of pyridotriazoles with alkynes and nitriles that leads to indolizines (X = CH) and imidazopyridines (X = N), respectively [Eq. (2)].




It has been shown that 2-pyridyl diazo compounds **1**^[6] transform into their cyclic triazole form **2**^[7] upon storage [Eq. (3)], and it is also known that some of these cyclic triazoles can still undergo transformations that are characteristic of diazo compounds.^[8] This phenomenon has been attributed to the closed/open form equilibrium of N-fused triazoles in solution,^[9] which can produce trace to significant

amounts of **1**. The position of this equilibrium depends on the temperature and the substitution pattern of the triazole.^[9b] Thus, introduction of a halogen substituent at C7 (R¹ = Cl) shifts the equilibrium to the left, which has been explained in terms of nonbonding repulsion between the lone pair of the halogen and that of the nitrogen in the *peri* position.^[10]



To evaluate the feasibility of using triazoles as precursors of Rh carbenoids we investigated the reaction of triazoles **3a** and **3b** with triethylsilane in the presence of a catalytic amount of rhodium(II) acetate, which is a method developed by Doyle and coworkers^[11] for the efficient trapping of Rh carbenoids [Eq. (4)]. Not surprisingly, pyridotriazoles **3a** and **3b** behave differently under these reaction conditions. Thus, while the 7-H derivative **3a** remains unaffected, the 7-chloro-substituted compound **3b** is smoothly converted into **4**, which is the product of carbenoid insertion into the Si–H bond. These experiments clearly indicate that 7-halo-substituted pyridotriazoles can indeed serve as convenient precursors of Rh carbenoids.




Next, to test our hypothesis regarding the annulation of α -imino diazo compounds with alkynes to form a pyrrole ring, we treated triazole **3b** with phenylacetylene in the presence of rhodium(II) acetate. This reaction proceeded smoothly to produce a mixture of cyclopropene **5** and indolizine **6a** with yields of 68% and 28% of isolated product, respectively [Eq. (5)]. Surprisingly, cyclopropene **5** does not undergo further isomerization into indolizine **6a** under these reaction conditions.^[12] The ratio of these products remained constant throughout the course of the reaction, thereby suggesting an independent path for the formation of **6a**.

We found, however, that the selectivity of the transannulation (**6** over **5**) could be dramatically improved by using

[*] S. Chuprakov, F. W. Hwang, Prof. V. Gevorgyan
Department of Chemistry
University of Illinois at Chicago
845 West Taylor Street, Chicago, IL 60607 (USA)
Fax: (+1) 312-355-0836
E-mail: vlad@uic.edu
Homepage: <http://www.chem.uic.edu/vggroup>

[**] Financial support from the National Institutes of Health (GM-64444) is gratefully acknowledged.

Supporting Information for this article is available on the WWW under <http://www.angewandte.org> or from the author.



rhodium(II) heptafluorobutyrate as catalyst.^[13] Thus, transannulation of **3b** with a series of aryl and alkenyl alkynes^[14] proceeded highly chemoselectively (90:10 to 95:5 vs. cyclopropene) to produce indolizines **6**^[15] in good yields (Table 1). Electron-rich, electron-deficient, and sterically hindered aryl alkynes were nearly equally effective in this reaction.

Table 1: Rhodium(II)-catalyzed transannulation of triazole **3b** with alkynes.

Entry	Alkyne	Product	Yield [%] ^[a]
1		6a	78
2		6b	80
3		6c	73
4		6d	85
5		6e	70
6		6f	65
7		6g	57

[a] Yield of isolated product. Indolizines **6** were accompanied by 5–10% of the corresponding cyclopropenes **5**; these compounds were readily separable by column chromatography.

Inspired by the successful formation of an N-fused pyrrole ring from the transannulation of triazoles with alkynes, we examined the formation of an N-fused imidazole ring in the reaction of **3** with nitriles and found that pyridotriazoles **3** react smoothly with a variety of aryl, alkyl, and alkenyl nitriles in the presence of $\text{Rh}_2(\text{OAc})_4$ (1 mol %) in toluene at 60 °C (Table 2) to afford N-fused imidazopyridines **7** in reasonable to high yields.


Both 3-carbomethoxy- (Table 2, entries 1–9) and 3-aryl- (Table 2, entry 10) pyridotriazoles are equally efficient in this reaction. Moreover, 7-bromo- (Table 2, entry 11) and even 7-methoxy-substituted (Table 2, entry 12) triazoles proved to be good substrates for this transannulation reaction.

We propose the following mechanism for this novel Rh-catalyzed transformation (Scheme 1). First, pyridotriazole **3** undergoes closed/open form equilibrium^[9] to produce small amounts of diazo compound **1** which, upon reaction with rhodium(II) carboxylate, generates the Rh-carbenoid species **I**. A direct nucleophilic attack^[18] of alkyne or nitrile **8** on species **I** produces ylide species **II**, according to path A, which

Table 2: Rhodium(II)-catalyzed transannulation of triazoles with nitriles.

Entry	R ¹	R ²	Triazole	R ³	Product	Yield [%] ^[a]
1	Cl	CO ₂ Me	3b	<i>p</i> -Tol	7a	89
2	Cl	CO ₂ Me	3b	Ph	7b	83
3	Cl	CO ₂ Me	3b	<i>p</i> -Me(O)CC ₆ H ₄	7c	54
4	Cl	CO ₂ Me	3b	Bn	7d	63
5	Cl	CO ₂ Me	3b	<i>n</i> Pr	7e	75
6	Cl	CO ₂ Me	3b	<i>c</i> Pr	7f	74
7	Cl	CO ₂ Me	3b	<i>t</i> Bu	7g	69
8	Cl	CO ₂ Me	3b		7h	66
9	Cl	CO ₂ Me	3b	CH ₂ SiMe ₃	7i	70
10	Cl	<i>p</i> -CF ₃ C ₆ H ₄	3c	Ph	7j	82
11	Br	<i>p</i> -CF ₃ C ₆ H ₄	3d	<i>p</i> -Tol	7k	73
12	OMe	<i>p</i> -CF ₃ C ₆ H ₄	3e	<i>n</i> Pr	7l	51

[a] Yield of isolated product.



Scheme 1. Plausible mechanisms for the Rh-catalyzed transannulation of pyridotriazoles with alkynes and nitriles. Y = N, CR^{''}.

then cyclizes to form **6** or **7** via cyclic zwitterion **III**. Alternatively (path B), [2+2] cycloaddition of **I** and **8** leads to metallacyclobutene **IV**, which can also be formed by cyclization of **II**.^[19] Rhodacycle **IV** then undergoes metathesis

to produce Rh carbenoid **V** which, upon 6π -electrocyclization and subsequent reductive elimination, furnishes product **6** or **7**. [2+1] Cycloaddition of **I** with **8** (path C) accounts for the formation of cyclopropene **5** in the presence of rhodium(II) acetate [see Eq. (5)]. As discussed above, **5** does not transform into heterocycle **6** under these reaction conditions.^[12]

In summary, we have developed an efficient Rh-catalyzed transannulation of pyridotriazoles for the formation of pyrrolo- and imidazopyridines, which are important fused heterocyclic scaffolds.^[20] We have also demonstrated that some of these pyridotriazoles can serve as stable^[13] and convenient^[21] precursors of Rh carbenoids.

Received: February 22, 2007

Published online: May 11, 2007

Keywords: annulation · nitrogen heterocycles · rhodium · synthetic methods · triazoles

- [1] For recent reviews, see: a) I. Nakamura, Y. Yamamoto, *Chem. Rev.* **2004**, *104*, 2127; b) G. Zeni, R. C. Larock, *Chem. Rev.* **2004**, *104*, 2285; c) M. Rubin, A. W. Sromek, V. Gevorgyan, *Synlett* **2003**, 2265.
- [2] a) H. M. L. Davies, K. R. Romines, *Tetrahedron* **1988**, *44*, 3343; b) H. M. L. Davies, W. R. Cantrell, Jr., K. R. Romines, J. S. Baum, *Org. Synth.* **1992**, *70*, 93.
- [3] a) F. R. Kinder, A. Padwa, *Tetrahedron Lett.* **1990**, *31*, 6835; b) A. Padwa, J. M. Kassir, S. L. Xu, *J. Org. Chem.* **1991**, *56*, 6971; c) A. Padwa, F. R. Kinder, *J. Org. Chem.* **1993**, *58*, 21.
- [4] For recent applications, see: a) V. Gettwert, F. Krebs, G. Maas, *Eur. J. Org. Chem.* **1999**, 1213; b) C. Batsila, G. Kostakis, L. P. Hadjiarapoglou, *Tetrahedron Lett.* **2002**, *43*, 5997.
- [5] R. Connell, F. Scavo, P. Helquist, *Tetrahedron Lett.* **1986**, *27*, 5559.
- [6] For cyclopropanation with 2-pyridyl diazo compounds, see: H. M. L. Davies, R. J. Townsend, *J. Org. Chem.* **2001**, *66*, 6595.
- [7] M. Regitz, *Angew. Chem.* **1967**, *79*, 786; *Angew. Chem. Int. Ed. Engl.* **1967**, *6*, 733.
- [8] See, for example: B. Abarca-González, *J. Enzym. Inhib. Med. Chem.* **2002**, *17*, 359, and references therein.
- [9] a) M. Regitz, B. Arnold, D. Danion, H. Schubert, G. Fusser, *Bull. Soc. Chim. Belg.* **1981**, *90*, 615; b) G. L'abbé, *Bull. Soc. Chim. Belg.* **1990**, *99*, 281; c) G. L'abbé, F. Godts, S. Toppet, *J. Chem. Soc. Chem. Commun.* **1985**, 589; d) G. L'abbé, I. Luyten, S. Toppet, *J. Heterocycl. Chem.* **1992**, *29*, 713; e) G. L'abbé, F. Godts, S. Toppet, *Bull. Soc. Chim. Belg.* **1986**, *95*, 679.
- [10] B. Abarca-González, R. Ballesteros, F. Mojarred, G. Jones, D. J. Mouat, *J. Chem. Soc. Perkin Trans. 1* **1987**, 1865.
- [11] V. Bagheri, M. P. Doyle, J. Taunton, E. E. Claxton, *J. Org. Chem.* **1988**, *53*, 6158.
- [12] For a related transformation of 3-carbonyl cyclopropenes into furans, see refs. [2a] and [3a].
- [13] See the Supporting Information for details.
- [14] The use of aliphatic alkynes resulted in sluggish and incomplete reactions.
- [15] Halo-substituted N-fused heterocycles can be further functionalized by cross-coupling reactions.^[16] We have also found that the chloride in **6a** can be quantitatively removed by treatment with $\text{PdCl}_2/\text{HSiEt}_3$.^[17]
- [16] For a review of coupling reactions of aryl chlorides, see: A. F. Littke, G. C. Fu, *Angew. Chem.* **2002**, *114*, 4350; *Angew. Chem. Int. Ed.* **2002**, *41*, 4176.
- [17] R. Boukherroub, C. Chatgililoglu, G. Manuel, *Organometallics* **1996**, *15*, 1508.
- [18] A. Padwa, D. J. Austin, A. T. Price, M. A. Semones, M. P. Doyle, M. N. Protopopova, W. R. Winchester, A. Tran, *J. Am. Chem. Soc.* **1993**, *115*, 8669.
- [19] T. R. Hoyer, C. J. Dinsmore, D. S. Johnson, P. F. Korkowski, *J. Org. Chem.* **1990**, *55*, 4518.
- [20] For selected reports of the biological activity of indolizines, see: a) S. Hagishita, M. Yamada, K. Shirahase, T. Okada, Y. Murakami, Y. Ito, T. Matsura, M. Wada, T. Kato, M. Ueno, Y. Chikazawa, K. Yamada, T. Ono, I. Teshirogi, M. Ohtani, *J. Med. Chem.* **1996**, *39*, 3636; b) L.-L. Gundersen, K. E. Malterud, A. H. Negussie, F. Rise, S. Teklu, O. B. Østby, *Bioorg. Med. Chem.* **2003**, *11*, 5409; for reports on the biological activity of imidazopyridines, see: c) D. Kim, L. Wang, J. J. Hale, C. L. Lynch, R. J. Budhu, M. MacCoss, S. G. Mills, L. Malkowitz, S. L. Gould, J. A. DeMartino, M. S. Springer, D. Hazuda, M. Miller, J. Kessler, R. C. Hrin, G. Carver, A. Carella, K. Henry, J. Lineberger, W. A. Schleif, E. A. Emini, *Bioorg. Med. Chem. Lett.* **2005**, *15*, 2129; d) S. Nakahara, A. Kubo, Y. Mikami, J. Ito, *Heterocycles* **2006**, *68*, 515.
- [21] The use of **3** does not require special slow-addition techniques as the concentration of **1** in the reaction mixture is always low.^[9]