

Alkylation of Aryl *N*-(2-Pyridylsulfonyl)aldimines with Organozinc Halides: Conciliation of Reactivity and Chemoselectivity**

Jorge Esquivias, Ramón Gómez Arrayás,* and Juan Carlos Carretero*

The direct addition of organometallic reagents to imines is a very convergent route to α -branched primary and secondary amines.^[1] Consequently, a number of synthetically useful addition reactions for the formation of carbon–carbon bonds have been developed with organolithium,^[2] Grignard,^[3] dialkyl zinc,^[4] alkenyl zirconium,^[5] aryl tin,^[6] aryl titanium,^[7] and aryl boron^[8] reagents as nucleophiles, including some very efficient catalytic enantioselective procedures.^[9] Despite this great progress, limitations in the scope of this reaction remain, especially with regard to functional-group compatibility. Highly reactive organometallic species, such as organolithium or Grignard reagents, have typically been used for the alkylation of imines, as the reactions occur under mild conditions with such reagents. However, drawbacks associated with these organometallic reagents are the competitive formation of reduction products and difficulties in the reconciliation of reactivity and chemoselectivity when functional groups are present in either the nucleophile or the imine substrate. The use of softer boron, tin, or silicon reagents has been limited mainly to arylation reactions^[6–8] and resonance-stabilized allyl metal species,^[10] which are more reactive than ordinary alkyl organometallic reagents. Organozinc reagents show an optimal compromise in terms of reactivity and wide functional-group tolerance, but only a limited number of simple dialkyl zinc reagents (typically dimethylzinc or diethylzinc) have been used for the alkylation of imines.^[4] Owing to their straightforward preparation^[11] and commercial availability, alkyl zinc halides offer a unique opportunity to satisfy the need for a general method for the alkylation of imines that combines high reactivity and wide functional-group tolerance. Although an increasing number of methodologies benefit from the unique reactivity/selectivity profile of $RZnX$ reagents,^[12] alkyl zinc halides have been scarcely used as nucleophiles in 1,2-addition reactions to

imines,^[13] probably as a result of the poor electrophilicity of the imine group.

In recent years, we^[14] and others^[3f,15] have demonstrated that *N*-(heteroarylsulfonyl) imines show better reactivity and/or selectivity than other *N*-sulfonyl imines, such as *N*-tosylimines, in some addition and cycloaddition reactions. Furthermore, the *N*-(heteroarylsulfonyl) group facilitates the often problematic deprotection of the amine functionality. Herein, we report a novel procedure for the alkylation of aromatic *N*-sulfonyl imines with alkyl zinc bromides. Key features of this method, which relies on the presence of a heteroarylsulfonyl coordinating group on the imine nitrogen atom, are high reactivity and broad functional-group compatibility, as well as straightforward deprotection to generate the free amine.

To find a suitable sulfonyl substituent, we treated a representative set of imines, **1a–g**, with *N*-hexylzinc bromide (2 equiv) in CH_2Cl_2 at room temperature (Table 1). Sub-

Table 1: Influence of the *N*-sulfonyl group on the reactivity of the imine.

Entry	Ar	1	<i>t</i>	Product	Yield [%] ^[a]
1	<i>p</i> -tolyl	1a	12 h	— ^[b]	—
2	<i>p</i> -NO ₂ C ₆ H ₄	1b	12 h	— ^[b]	—
3	<i>o</i> -NO ₂ C ₆ H ₄	1c	12 h	— ^[b]	—
4	2-thienyl	1d	12 h	— ^[b]	—
5	8-quinolyl	1e	15 min	2e	80
6	2-pyridyl	1f	5 min	2f	97
7	3-pyridyl	1g	12 h	— ^[b]	—

[a] After chromatographic purification. [b] Only the products of imine hydrolysis were recovered after aqueous workup. Tf = trifluoromethanesulfonyl.

strates **1a** with a tosyl group, **1b** with a *p*-nosyl group, **1c** with an *o*-nosyl group, and **1d** with a 2-thienylsulfonyl group did not undergo the desired reaction in the presence of $Cu(OTf)_2$ (10 mol %; Table 1, entries 1–4).^[16] Only products of imine hydrolysis were recovered following an aqueous workup. In contrast, the (8-quinolylsulfonyl)imine **1e**^[14d] and (2-pyridylsulfonyl)imine **1f**^[14a,b] underwent smooth addition reactions under the catalysis of $Cu(OTf)_2$ (10 mol %) to afford the corresponding adducts **2e** and **2f** in 80 and 97 % yield, respectively, in 15 min or less (Table 1, entries 5 and 6). Imine **1f** underwent addition with *N*-hexylzinc bromide even in the absence of a copper salt, although the reaction became much slower (8 h at room temperature), and the yield of **2f**

[*] J. Esquivias, Dr. R. Gómez Arrayás, Prof. Dr. J. C. Carretero
Departamento de Química Orgánica
Facultad de Ciencias
Universidad Autónoma de Madrid (UAM)
Cantoblanco 28049 Madrid (Spain)
Fax: (+34) 91-497-3966
E-mail: ramon.gomez@uam.es
juancarlos.carretero@uam.es

[**] This research was supported by the Ministerio de Educación y Ciencia (MEC, project CTQ2006-01121) and UAM–Consejería de Educación de la Comunidad Autónoma de Madrid (CAM, project CCG06-UAM/PPQ-0557). J.E. thanks the MEC for a predoctoral fellowship. We thank Prof. J. L. Mascareñas and Dr. F. López for helpful discussions.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

decreased to 64 % as a result of partial hydrolysis of the imine. Interestingly, the (3-pyridylsulfonyl)imine **1g**, an isomer of **1f**, proved to be unreactive under the optimized reaction conditions (Table 1, entry 7), which suggests that a chelating interaction of the heteroaryl moiety with the metal center may be responsible for the dramatic increase in reactivity observed with imines **1e** and **1f**.

Next, the influence of catalyst loading was investigated for the addition of the secondary alkyl zinc reagent 2-pentylzinc bromide to **1f** (Table 2). The addition was equally effective with 5 mol % (Table 2, entry 1) and 1 mol % (entry 2) of

Table 2: Effect of the catalyst loading and scaling up on the reaction.

Entry	Quantity of 1f [mmol]	x	t [min]	Yield [%] ^[a]
1	0.2	5	5–10	92
2	2	1	15	90
3	4	0.1	75	77
4	0.2	0	270	66

[a] After chromatographic purification.

$\text{Cu}(\text{OTf})_2$: In both cases the product **3f** was formed in over 90 % yield in less than 15 min.^[18] As a further demonstration of the potential of this method, the addition of 2-pentylzinc bromide to imine **1f** on a gram scale (4 mmol) in the presence of 0.1 mol % of $\text{Cu}(\text{OTf})_2$ (molar ratio S/C = 1000) led to **3f** in 77 % yield after 75 min at room temperature (Table 2, entry 3). In the absence of a copper catalyst, the reaction afforded **3f** in 66 % yield after 4.5 h (Table 2, entry 4).

The scope of this Cu-catalyzed reaction was examined with a variety of aryl and heteroaryl *N*-(2-pyridylsulfonyl)aldimines (Table 3). Aliphatic imines were not investigated, as to date we have not succeeded in developing a general method for the isolation of this more labile type of imine.^[19] For practical reasons, this study was undertaken with 5 mol % of $\text{Cu}(\text{OTf})_2$, although we confirmed later that the catalyst is equally effective in a loading of 1 mol % (Table 3, entry 27). A wide variety of alkyl zinc bromide reagents of different substitution patterns and degrees of functionalization participated in the addition reaction to provide the corresponding protected amines in less than 10 min in yields typically in the range of 80–90 %. Primary and secondary alkyl zinc bromide reagents with a broad variety of functional groups, including alkene, ether, acetal, chloride, ester, and nitrile groups, were used successfully in this procedure. Even an unprotected indole substituent with an acidic NH group proved to be compatible (Table 3, entry 13). Only the very bulky reagent *tert*-butylzinc bromide was unreactive under these conditions (Table 3, entry 5).

With respect to the imine substrate, aryl moieties with donor or acceptor substituents at the *ortho*, *meta*, or *para* position are tolerated well (Table 3, entries 14–21). Heteroaromatic 2-furyl, 2-thienyl, and 2-pyrrolyl imine derivatives also

Table 3: Scope of the direct addition of alkyl zinc bromides to imines.


Entry	R ¹	Imine	R ²	Product	Yield [%] ^[a]
1	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	2f	93
2	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	3f	92
3	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	13f	88
4	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	14f	81
5	Ph	1f	<i>t</i> Bu	—	— ^[b]
6	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	15f	90
7	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	16f	84
8	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	17f	92
9	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	18f	95
10	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	19f	88
11	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	20f	90
12	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	21f	90
13	Ph	1f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	22f	82
14	4-MeOC ₆ H ₄	4f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	23f	80
15	4-MeOC ₆ H ₄	4f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	24f	83
16	4-FC ₆ H ₄	5f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	25f	88
17	4-FC ₆ H ₄	5f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	26f	85
18	2-BrC ₆ H ₄	6f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	27f	87
19	3-MeC ₆ H ₄	7f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	28f	84
20	2-naphthyl	8f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	29f	91
21	2-naphthyl	8f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	30f	89
22	2-furyl	9f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	31f	92
23	2-furyl	9f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	32f	90
24	2-thienyl	10f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	33f	80
25	Indol-2-yl	11f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	34f	68 ^[c]
26	4-MeC ₆ H ₄	12f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	35f	77
27	4-MeC ₆ H ₄	12f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	36f	83 ^[d]
28	4-MeC ₆ H ₄	12f	CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ CH ₃	37f	85

[a] After chromatographic purification. [b] The product was not observed after 16 h at room temperature. [c] Reaction time: 3 h. [d] $\text{Cu}(\text{OTf})_2$: 1 mol %.

proved to be suitable (Table 3, entries 22–25). The more challenging substrate **12f**, with a potentially competitive ketone group, underwent imine alkylation with complete chemoselectivity (Table 3, entries 26–28).

In a competition experiment that highlights the exceptional reactivity of the *N*-(2-pyridylsulfonyl)imines, an equimolar mixture of benzaldehyde and imine **1f** was treated with

4-chlorobutylzinc bromide (2.0 equiv) under the optimized reaction conditions. After a reaction time of 5 min, the sulfonamide adduct **18f** was obtained as the only product. No trace of the product of addition to the aldehyde was detected by NMR spectroscopy, and benzaldehyde was recovered (Scheme 1).



Scheme 1. Chemoselectivity competition experiment between (2-pyridylsulfonyl)imine **1f** and benzaldehyde. Py = pyridyl.

In connection with our studies on the conjugate addition of Et_2Zn to α,β -unsaturated *N*-(2-pyridylsulfonyl)ketimines,^[14a] we examined the addition of alkyl zinc bromides to **38**, the (2-pyridylsulfonyl)imine of chalcone (Scheme 2).



Scheme 2. Conjugate addition of alkyl zinc bromides to the *N*-(2-pyridylsulfonyl)imine of chalcone.

Only conjugate addition was observed,^[20] and the corresponding enamides **39–41** were obtained in high yields (88–97%) and with very good *Z* stereoselectivity.^[21] In contrast, chalcone itself and its *N*-tosylimine were recovered unaltered after treatment under the same conditions, even after prolonged reaction times.

The exceptional reactivity of 2-pyridylsulfonylimines prompted us to study the mode of their coordination to copper. Initial attempts to isolate a copper complex of aldimine **1f** failed as a result of its high propensity to undergo hydrolysis. However, the treatment of the more stable ketimine **38** with 1 equivalent of $[\text{Cu}(\text{CH}_3\text{CN})_4]\text{PF}_6$ ^[22] in CH_2Cl_2 at room temperature produced an instantaneous color change from colorless to deep red, accompanied by nearly quantitative formation of complex **42**, which was isolated as an air-stable red solid. X-ray diffraction analysis of suitable crystals^[23] revealed a distorted trigonal-pyramidal $\text{N}_2\text{N}_2\text{N}_2$ -tetracoordinated copper complex^[24] in which the Cu^{I} atom coordinates to two molecules of **38** through the imino and pyridyl nitrogen atoms (Figure 1). The observation that the bonds between Cu^{I} and the imine nitrogen atoms (2.34 Å) are much longer than those between Cu^{I} and the pyridine nitrogen atoms (1.93 Å) reflects the higher affinity of the metal center to the more basic pyridine nitrogen atoms. The bidentate N_2N chelation of *N*-(2-pyridylsulfonyl)imines through a favorable five-membered chelate ring^[25] with the electrophilic copper salt provides a reasonable explanation for the much higher reactivity of 2-pyridylsulfonylimines over that of nonchelating *N*-sulfonyl imines.^[26]





Figure 1. X-ray crystal structure of complex **42**. (The PF_6^- anion and the hydrogen atoms have been omitted).

The 2-pyridylsulfonyl group was removed by the simple treatment of the sulfonamide adducts with an excess of Mg in MeOH at room temperature for 2 h.^[27] The corresponding primary amines were obtained in 90–94% yield (Scheme 3).



Scheme 3. Deprotection of the sulfonamide adducts and cyclization reactions.

This smooth deprotection reaction is compatible with sensitive functional groups, such as ketals, halides, and esters. The free chloroamine **44** was converted into the piperidine derivative **45** in 78% yield by subsequent treatment with K_2CO_3 . The deprotection of the sulfonamide esters **19f** and **36f** led to the δ -lactams **46** and **47**, respectively, in excellent yields (> 90%) through the spontaneous cyclization of the corresponding free amines under the reaction conditions.

In conclusion, readily available alkyl zinc bromide reagents are excellent nucleophilic species for the alkylation of aromatic *N*-(2-pyridylsulfonyl)aldimines with unprecedented functional-group compatibility. Functionalized free amines can be accessed in very good yields upon facile deprotection under mild conditions. We attribute the exceptional reactivity of these imines to their N_2N -bidentate character with respect to metal coordination, evidence for which was obtained from an X-ray crystallographic study of a Cu^{I} complex.

Received: July 19, 2007

Revised: September 5, 2007

Published online: October 30, 2007

Keywords: alkylation · copper · N_2N ligands · sulfonyl imines · zinc

- [1] For general reviews, see: a) D. Enders, U. Reinhold, *Tetrahedron: Asymmetry* **1997**, 8, 1895; b) R. Bloch, *Chem. Rev.* **1998**, 98, 1407; for a review on *N*-phosphinoyl imines, see: c) S. M. Weinreb, R. K. Orr, *Synthesis* **2005**, 1205–1227; for a review on *N*-acyl imines, see: M. Petrini, E. Torregiani, *Synthesis* **2007**, 159.
- [2] See, for example: a) S. E. Denmark, C. M. Stiff, *J. Org. Chem.* **2000**, 65, 5875; b) S. Arrasate, E. Lete, N. Sotomayor, *Tetrahedron: Asymmetry* **2001**, 12, 2077; c) C. Cimarrelli, G. Palmieri, E. Volpini, *J. Org. Chem.* **2003**, 68, 1200; d) A. A. Boezio, G. Solberghe, C. Lauzon, A. B. Charette, *J. Org. Chem.* **2003**, 68, 3241.
- [3] See, for example: a) S. V. Kolotuchin, A. I. Meyers, *J. Org. Chem.* **2000**, 65, 3018; b) T. Takahashi, Y. Liu, C. Xi, S. Huo, *Chem. Commun.* **2001**, 31; c) V. Gandon, P. Bertus, J. Szymoniak, *Eur. J. Org. Chem.* **2001**, 3677; d) S. Saito, K. Hatanaka, H. Yamamoto, *Synlett* **2001**, 1859; e) B. Z. Lu, C. Senanayake, N. Li, Z. Han, R. P. Bakale, S. A. Wald, *Org. Lett.* **2005**, 7, 2599; f) H. Sugimoto, S. Nakamura, M. Hattori, S. Ozeki, N. Shibata, T. Toru, *Tetrahedron Lett.* **2005**, 46, 8941; g) A. Krasovskiy, F. Kopp, P. Knochel, *Angew. Chem.* **2006**, 118, 511; *Angew. Chem. Int. Ed.* **2006**, 45, 497; h) M. Hatano, S. Suzuki, K. Ishihara, *J. Am. Chem. Soc.* **2006**, 128, 9998.
- [4] For recent examples, see: a) C. J. Wang, M. Shi, *J. Org. Chem.* **2003**, 68, 6229; b) T. Soeta, K. Nagai, H. Fujihara, M. Kuriyama, K. Tomioka, *J. Org. Chem.* **2003**, 68, 9723; c) L. C. Akullian, J. R. Porter, J. F. Traverse, M. L. Snapper, A. H. Hoveyda, *Adv. Synth. Catal.* **2005**, 347, 417; d) C. Lauzon, A. B. Charette, *Org. Lett.* **2006**, 8, 2743; e) T. Nishimura, Y. Yasuhara, T. Hayashi, *Org. Lett.* **2006**, 8, 979; f) S. Basra, M. W. Fennie, M. C. Kozlowski, *Org. Lett.* **2006**, 8, 2659.
- [5] a) P. Wipf, C. Kendall, C. R. J. Stephenson, *J. Am. Chem. Soc.* **2001**, 123, 5122; b) A. Kakuuchi, T. Taguchi, Y. Hanzawa, *Tetrahedron Lett.* **2003**, 44, 923.
- [6] a) T. Hayashi, M. Ishigedani, *J. Am. Chem. Soc.* **2000**, 122, 976; b) T. Hayashi, M. Ishigedani, *Tetrahedron* **2001**, 57, 2585.
- [7] T. Hayashi, M. Kawai, N. Tokunaga, *Angew. Chem.* **2004**, 116, 6251; *Angew. Chem. Int. Ed.* **2004**, 43, 6125.
- [8] a) M. Kuriyama, T. Soeta, X. Hao, Q. Chen, K. Tomioka, *J. Am. Chem. Soc.* **2004**, 126, 8128; b) N. Tokunaga, Y. Otomaru, K. Okamoto, K. Ueyama, R. Shintani, T. Hayashi, *J. Am. Chem. Soc.* **2004**, 126, 13584; c) Y. Otomaru, N. Tokunaga, R. Shintani, T. Hayashi, *Org. Lett.* **2005**, 7, 307; d) R. B. C. Jagt, P. Y. Touleec, D. Geerdink, J. G. de Vries, B. L. Feringa, A. J. Minnaard, *Angew. Chem.* **2006**, 118, 2855; *Angew. Chem. Int. Ed.* **2006**, 45, 2789.
- [9] For recent reviews on catalytic enantioselective addition to imines, see: a) T. Vilaivan, W. Bhanthumnavin, Y. Sritananant, *Curr. Org. Synth.* **2005**, 2, 1315; b) G. K. Friestad, A. K. Mathies, *Tetrahedron* **2007**, 63, 2541.
- [10] For reviews on the addition of allylic organometallic reagents to imines, see: a) H. Ding, G. K. Friestad, *Synthesis* **2005**, 2815; b) P. Merino, T. Tejero, J. I. Delso, V. Mannucci, *Curr. Org. Synth.* **2005**, 2, 479; c) P. V. Ramachandran, T. E. Burghardt, *Pure Appl. Chem.* **2006**, 78, 1397.
- [11] For the preparation of alkyl zinc halides, see: a) S. Huo, *Org. Lett.* **2003**, 5, 423; b) F. O. Arp, G. C. Fu, *J. Am. Chem. Soc.* **2005**, 127, 10482; c) A. Krasovskiy, V. Malakhov, A. Gavryushin, P. Knochel, *Angew. Chem.* **2006**, 118, 6186; *Angew. Chem. Int. Ed.* **2006**, 45, 6040.
- [12] For a review on the reactivity and utility of organozinc compounds, see: *The Chemistry of Organozinc Compounds: R-Zn* (Eds.: Z. Rappoport, I. Marek), Wiley, Chichester, **2006**; for very recent examples, see: a) A. Whitehead, J. P. McParland, P. R. Hanson, *Org. Lett.* **2006**, 8, 5025; b) F. Denes, S. Cutri, A. Perez-Luna, F. Chemla, *Chem. Eur. J.* **2006**, 12, 6506; c) H. Ren, G. Dunet, P. Mayer, P. Knochel, *J. Am. Chem. Soc.* **2007**, 129, 5376; d) G. Sklute, C. Bolm, I. Marek, *Org. Lett.* **2007**, 9, 1259.
- [13] a) K. P. Chiev, S. Roland, P. Mangeney, *Tetrahedron: Asymmetry* **2002**, 13, 2205; b) T. Iwai, T. Mizuno, Y. Ishino, *Tetrahedron Lett.* **2004**, 45, 1083.
- [14] a) J. Esquivias, R. Gómez Arrayás, J. C. Carretero, *J. Org. Chem.* **2005**, 70, 7451; b) J. Esquivias, R. Gómez Arrayás, J. C. Carretero, *Angew. Chem.* **2006**, 118, 645; *Angew. Chem. Int. Ed.* **2006**, 45, 629; c) A. S. González, R. Gómez Arrayás, J. C. Carretero, *Org. Lett.* **2006**, 8, 2977; d) J. Esquivias, R. Gómez Arrayás, J. C. Carretero, *J. Am. Chem. Soc.* **2007**, 129, 1480.
- [15] a) S. Nakamura, H. Nakashima, H. Sugimoto, N. Shibata, T. Toru, *Tetrahedron Lett.* **2006**, 47, 7599; b) S. Nakamura, H. Sano, H. Nakashima, K. Kubo, N. Shibata, T. Toru, *Tetrahedron Lett.* **2007**, 48, 5565; see also: c) H. Morimoto, G. Lu, N. Aoyama, S. Matsunaga, M. Shibasaki, *J. Am. Chem. Soc.* **2007**, 129, 9588.
- [16] The same negative result was obtained with Cu^I catalysts, such as copper(I) thiophene-2-carboxylate (CuTC) and [Cu(CH₃CN)₄]ClO₄, and in the absence of a copper salt.
- [17] Cu^I salts, such as [Cu(CH₃CN)₄]PF₆ or CuTC, were also effective catalysts in a concentration of 10 mol % for the reaction of **1f**: Compound **2f** was formed in 93 or 94 % yield within 15 min.
- [18] In this model reaction, a similar high reactivity was observed in the presence of Cu(OTf)₂ (5 mol %) and a phosphine ligand (5 mol %), such as 2,2'-bis(diphenylphosphanyl)-1,1'-binaphthyl (binap). Unfortunately, however, the racemic product **3f** was obtained when (*R*)-binap was used as the ligand.
- [19] Although aliphatic *N*-(2-pyridylsulfonyl)imines appear to be formed quantitatively in 2–3 h upon the direct condensation of an aldehyde with 2-pyridylsulfonamide in the presence of Ti(OEt)₄ (1 equiv) in CH₂Cl₂ at room temperature (TLC monitoring), all attempts to isolate the pure aldimine resulted in the recovery of the hydrolysis products.
- [20] For examples of the Michael-type addition of organometallic reagents to α,β-unsaturated imines, see: a) K. Tomioka, Y. Shioya, Y. Nagaoka, K. Yamada, *J. Org. Chem.* **2001**, 66, 7051; b) J. P. McMahon, J. A. Ellman, *Org. Lett.* **2005**, 7, 5393; c) T. Soeta, M. Kuriyama, K. Tomioka, *J. Org. Chem.* **2005**, 70, 297; d) F. Palacios, J. Vicario, *Org. Lett.* **2006**, 8, 5405; ref. [13a].
- [21] The *Z* configuration was assigned to the major *N*-sulfonyl enamide by analogy with the corresponding product of the addition of Et₂Zn (see ref. [13a]). The configuration of the latter product was in turn established by NMR spectroscopy.
- [22] [Cu(CH₃CN)₄]PF₆ was chosen as the stoichiometric copper source as a result of its much higher solubility in CH₂Cl₂ relative to that of Cu(OTf)₂.
- [23] See the Supporting Information for experimental details and crystallographic data. CCDC-654016 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [24] For an N,N,N,N-tetracoordinated Cu^{II} complex of bis(2-pyridyl)sulfonamido ligands, see: A. Nanthakumar, J. Miura, S. Diltz, C.-K. Lee, G. Aguirre, F. Ortega, J. W. Ziller, P. J. Walsh, *Inorg. Chem.* **1999**, 38, 3010.
- [25] This N,N coordination the imine to copper contrasts with the N,O-type metal complexes of (2-pyridylsulfonyl)aldimines proposed by Toru and co-workers on the basis of theoretical calculations, whereby the pyridyl nitrogen atom and one of the sulfonyl oxygen atoms were proposed to form a five-membered chelate ring with the metal center; see refs. [3f, 14a].
- [26] This mode of coordination could also facilitate pseudointramolecular delivery of the organocopper nucleophile following the Zn/Cu transmetalation; see, for example: P. Knochel, R. D. Singer, *Chem. Rev.* **1993**, 93, 2117.
- [27] C. S. Pak, D. S. Lim, *Synth. Commun.* **2001**, 31, 2209.