


Tetrasubstituted Furans by a Pd^{II}-Catalyzed Three-Component Michael Addition/Cyclization/Cross-Coupling Reaction **

Yuanjing Xiao and Junliang Zhang*

The design and discovery of new reactions for the synthesis of highly substituted furans has been stimulated by their appearance in many bioactive natural products and important pharmaceuticals.^[1,2] There has been recent focus on the development of metal-catalyzed one- or two-component reactions for synthesis of these types of compounds, including the cyclization of allenyl ketones,^[3] 3-alkyn-1-ones,^[3k,4] 1-(1-alkynyl)-cyclopropyl ketones,^[5] (Z)-2-en-4-yn-1-ols,^[6] and 2-(1-alkynyl)-2-alken-1-ones,^[7] and cycloisomerization of cyclopropyl ketones^[8]/cyclopropenyl ketones.^[9] Recently, the design of multicomponent reactions^[10] that preserve atom economy in a one-pot reaction has attracted attention because of the application to efficient construction of molecular structures. Herein we report results of a Pd^{II}-catalyzed three-component Michael addition/cyclization/cross-coupling reaction to afford highly functionalized tetrasubstituted furans.^[11]

As a continuation of our interest in the design and discovery of new reactions for the synthesis of highly substituted furans,^[5,8,9b] we envisioned that 2-(1-alkynyl)-2-alken-1-ones might react with nucleophiles and an allyl halide in the presence of a catalytic Pd^{II} species that might exhibit dual roles, serving simultaneously as a Lewis acid and a transition metal (Scheme 1). First, Pd^{II} acts as a Lewis acid and a transition metal to facilitate both the nucleophilic addition step and the cyclization step^[12] to afford a furanyl-palladium intermediate, which then reacts with the allyl halide by insertion and subsequent β -halide elimination to provide product furans and regenerate the PdX₂ catalyst. To the best of our knowledge, only one example of Pd(OAc)₂ serving as a dual role catalyst exists.^[13] Herein we present our recent results on the [PdCl₂(CH₃CN)₂]-catalyzed three-component Michael addition/cyclization/cross-coupling of 2-(1-alkynyl)-2-alken-1-ones **1** with various nucleophiles and allyl chlorides.



Scheme 1. a) General reaction scheme for the Pd^{II}-catalyzed three-component reaction to afford highly substituted furans. b) Proposed reaction pathway for the reaction in (a). In the first step the Pd^{II} serves as a Lewis acid and a transition metal, and in subsequent steps the Pd^{II} serves as a transition metal.

First, we examined the reaction of 2-phenylethynyl-3-phenyl-2-buten-1-one (**1a**) with MeOH and allyl chloride (**3a**) under different reaction conditions. After numerous attempts, we isolated the desired product **4aaa** in 75% yield after running the reaction for 24 hours at room temperature in CH₃CN with [PdCl₂(CH₃CN)₂] (5 mol%) as the catalyst, methanol (4.0 equiv) as the nucleophile, allyl chloride (4.0 equiv) as the allylating reagent, and K₂CO₃ (4.0 equiv) as the base. Only trace amounts (<1%) of the furan generated by the competitive side reaction from protonation could be detected by ¹H NMR spectra analysis (Table 1, entry 6). Alternate conditions, such as lowering the equivalents of the base, MeOH, or allylic chloride or changing the solvent or the base lead to a lower yield of the product (Table 1, entries 1–12). Surprisingly, when allyl bromide (even 10 equiv) was used instead of allyl chloride, the reaction proceeded slowly to give the desired furan in 37% yield (determined by ¹H NMR spectroscopy) after 48 hours. We deduced that relative to [PdCl₂(CH₃CN)₂], the weak Lewis acid [PdBr₂(CH₃CN)₂] that is generated in the reaction is not strong enough to activate the substrate. To support this deduction, two control reactions were carried out under the standard conditions with the exception that 20 mol% of Ph₃P (Table 1, entry 14) or 1 equivalent of KBr (Table 1, entry 15) was added; neither of the reactions worked well and the yield of the product was not more than 20%, thus, further verifying that [PdCl₂(CH₃CN)₂] is a stronger Lewis acid than [PdBr₂(CH₃CN)₂].

[*] Dr. Y. Xiao, Prof. Dr. J. Zhang
Shanghai Key Laboratory of Green Chemistry and Chemical Processes
Department of Chemistry
East China Normal University
3663 N. Zhongshan Road, Shanghai 200062 (P.R. China)
Fax: (+86) 21-6223-3213
E-mail: jlzhang@chem.ecnu.edu.cn

[**] Financial support from the National Science Foundation of China and East China Normal University is greatly appreciated. This work was sponsored by the Shanghai Pujiang Program (07pj14039) and the Shanghai Leading Academic Discipline Project (B409).

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

Table 1: Screening of reaction conditions for the three-component Michael addition/cyclization/cross-coupling reaction.^[a]

Entry	Base (equiv)	2a/3a [equiv]	Solvent	t [h]	Yield [%] ^[b]
1	none	4.0/4.0	CH ₃ CN	36	0
2	Na ₂ CO ₃ (4.0)	4.0/4.0	CH ₃ CN	36	41
3	NaHCO ₃ (4.0)	4.0/4.0	CH ₃ CN	36	50
4	K ₃ PO ₄ (4.0)	4.0/4.0	CH ₃ CN	36	62
5	CS ₂ CO ₃ (4.0)	4.0/4.0	CH ₃ CN	24	42
6	K ₂ CO ₃ (4.0)	4.0/4.0	CH ₃ CN	24	88 (75)
7	K ₂ CO ₃ (2.0)	4.0/4.0	CH ₃ CN	24	80
8	K ₂ CO ₃ (2.0)	2.0/2.0	CH ₃ CN	24	53
9	K ₂ CO ₃ (2.0)	2.0/4.0	CH ₃ CN	24	62
10	K ₂ CO ₃ (4.0)	4.0/4.0	THF	24	52
11	K ₂ CO ₃ (4.0)	4.0/4.0	1,4-dioxane	24	68
12	K ₂ CO ₃ (4.0)	4.0/4.0	DMF	24	69
13 ^[c]	K ₂ CO ₃ (4.0)	4.0/4.0	CH ₃ CN	24	0
14 ^[d]	K ₂ CO ₃ (4.0)	4.0/4.0	CH ₃ CN	24	0
15 ^[e]	K ₂ CO ₃ (4.0)	4.0/4.0	CH ₃ CN	24	20
16 ^[f]	K ₂ CO ₃ (4.0)	4.0/4.0	CH ₃ CN	48	37
17 ^[g]	K ₂ CO ₃ (4.0)	4.0/4.0	CH ₃ CN	48	78









[a] Reactions run with **1a** (0.5 mmol) at room temperature. [b] Yields were determined by ¹H NMR spectroscopy using CH₂Br₂ as an internal standard; yields of isolated product are shown in parentheses. [c] 5 mol % of [Pd(PPh₃)₄] was used. [d] 20 mol % of Ph₃P was added. [e] 1.0 equiv of KBr was added. [f] 10 equiv of allyl bromide used instead of allyl chloride. [g] 5 mol % of allylpalladium chloride dimer was used.

(CH₃CN)₂ and [PdCl₂(PPh₃)₂] (generated by adding Ph₃P to the mixture). In addition, no reaction occurred when [Pd-(PPh₃)₄] was employed as the catalyst instead of [PdCl₂-(CH₃CN)₂]. All these observations indicate that [PdCl₂-(CH₃CN)₂], acting simultaneously as a Lewis acid and a transition-metal catalyst, is the key to making this transformation work well.^[14]

To determine the scope of this transformation, we studied the reaction of **1a** with various nucleophiles **2** and allyl chlorides **3**, and the results are listed in Table 2. Notably: 1) Alcohols other than methanol, such as isopropanol, benzyl alcohol, and phenol, can act as O nucleophiles to give tetrasubstituted furans in moderate to excellent yields (Table 2, entries 1–6); 2) Dimethyl malonate can act as a nucleophilic component to allow the introduction of more functional groups (Table 1, entries 7–11); 3) Substituted allyl chlorides, such as **3b–d**, can be used as allylating reagents to give more functionalized furans in good yields (Table 2, entries 1–3, 8–11); 4) The coupling occurs exclusively at the less substituted terminus of the allyl chloride and the configuration of the C=C bond in 3-phenyl-2-propenyl chloride **3e** remained intact when it was introduced as the allylating agent (Table 2, entry 10), a result that was consistent with previous findings.^[3g]

This transformation can be successfully extended to a variety of 2-(1-alkynyl)-2-alken-1-ones **1** leading to the generation of the corresponding tetrasubstituted furan in 40–91 % yield (Table 3, entries 1–7). The substituents on the

Table 2: Scope of the [PdCl₂(CH₃CN)₂]-catalyzed three-component Michael addition/cyclization/cross-coupling of **1a**.^[a]

				
Entry	NuH (2)	3	t [h]	Yield (4) ^[a]
1	MeOH		14	 R = Me, 4aab , ^[f] 66 %
2 ^[b]	2a		36	R = CO ₂ Me, 4aac , 70 %
3	2a		48	 4aad , 89 %
4 ^[b,c]	<i>i</i> PrOH	3a	72	 R = <i>i</i> Pr, 4aba , 42 %
5	BnOH 2c	3a	24	R = Bn, 4aca , 63 %
6	PhOH 2d	3a	24	R = Ph, 4ada , 66 %
7 ^[d]		3a	10	 R = H, R' = H, 4aea , 91 %
8 ^[d]	2e	3b	12	R = Me, R' = H, 4aeb , 62 %
9 ^[d]	2e	3c	20	R = CO ₂ Me, R' = H, 4aec , 54 %
10 ^[d,e]	2e		24	R = H, R' = Ph, 4aee , 79 %
11 ^[d]	2e	3d	24	 4aed , 85 %

[a] Unless otherwise specified, reactions were carried out under standard conditions. Reported yields are of the isolated product. [b] Additional [PdCl₂(CH₃CN)₂] (5 mol %) was added after the reaction was stirred for 24 h. [c] Used *i*PrOH (6.0 equiv) and allyl chloride (6.0 equiv). [d] 2.0 equiv of dimethyl malonate were used. [e] Additional 0.5 equiv of dimethyl malonate was added after the reaction was stirred for 12 h. [f] The designation **4aab** for the product indicates that the reactants used were **1a**, **2a**, and **3b**, respectively.

Table 3: [PdCl₂(CH₃CN)₂]-catalyzed three-component Michael addition/cyclization/cross-coupling of various 2-(1-alkynyl)-2-alken-1-ones **1** with MeOH and allyl chloride.^[a]

Entry	1	<i>t</i> [h]	Yield 4 ^[b]
1	 1b	30	 4baa , 77%
2	 1c	24	 4caa , 80%
3	 1d	24	 4daa , 91%
4 ^[c]	 1e	43	 4eaa , 40%
5	 1f	36	 4faa , 73%
6	 1g	18	 4gaa , 65%
7	 1h	24	 4haa , 57%
8	 1i	72	 4iaa , 63%

[a] Unless otherwise noted the reactions were performed under standard conditions. [b] Yield of isolated product. [c] Reaction conditions: MeOH (8.0 equiv), allyl chloride (8.0 equiv), K₂CO₃ (6.0 equiv), and [PdCl₂(CH₃CN)₂] (10 mol%).

alkynes can affect the reaction; for example, alkynes bearing aryl groups afford relatively higher yields than those bearing alkyl groups (Table 3, compare entry 1 to entry 4). The type of nucleophile also affects the yield and the rate of the reaction (compare entries 4–7 in Table 2). Cyclic substrate **1h** also affords a fused furan in 57% yield under these reaction conditions (Table 3, entry 7). The results in Tables 2 and 3 clearly show that our new approach to tetrasubstituted furans allows easy introduction of various functional groups, such as alkoxy groups, aryloxy groups, and dimethyl malonate groups, as well as various types of allylic groups directly incorporated into the furan ring. The new method provides opportunities for further elaboration of the functional groups on the polysubstituted furans.

In conclusion, we have developed a [PdCl₂(CH₃CN)₂]-catalyzed three-component Michael addition/cyclization/cross-coupling reaction that provides an efficient route to functionalized tetrasubstituted furans. We found that [PdCl₂(CH₃CN)₂] plays a dual role in this transformation. This finding will be helpful to other cases in which activation of a carbonyl group or related functional group, such as enone (Lewis acid role), and activation of a double or triple bond (transition-metal role) are desired to occur simultaneously. As all three starting materials are readily available, this method may allow the synthesis of more complex furans. Further studies on the scope and synthetic applications of this methodology are currently being carried out in our laboratory.

Experimental Section

Typical procedure for the synthesis of **4aaa** (Table 1, entry 6): [PdCl₂(CH₃CN)₂] (6.0 mg, 0.025 mmol) was added to a mixture of **1a** (123.0 mg, 0.5 mmol), MeOH (64.0 mg, 2.0 mmol), allyl chloride (153.0 mg, 2.0 mmol), and K₂CO₃ (276.0 mg, 2.0 mmol) in CH₃CN (2.0 mL) at room temperature. The reaction mixture was stirred for 24 h, at which time **1a** was consumed completely according to TLC analysis, and then concentrated in vacuo after filtration. The residue was purified by column chromatography on silica gel (petroleum ether/Et₂O = 10:1) to give desired product **4aaa** (119 mg) in 75% yield as a colorless oil. ¹H NMR (500 MHz, CDCl₃): δ = 7.66–7.59 (m, 2H), 7.40–7.34 (m, 6H), 7.28–7.20 (m, 2H), 5.90–5.78 (m, 1H), 5.29 (s, 1H), 5.04–4.96 (m, 2H), 3.41 (s, 3H) 3.29–3.25 (m, 2H), 2.30 ppm (s, 3H); ¹³C NMR (125 MHz, CDCl₃): δ = 149.04, 148.00, 141.30, 135.86, 131.43, 128.43, 128.20, 127.19, 126.72, 126.64, 125.36, 120.86, 118.36, 115.45, 77.38, 56.78, 28.29, 12.58 ppm; MS (70 eV): *m/z* (%): 318 (3.10) [M⁺], 105 (100) [C₆H₅CO]⁺, HRMS calcd for C₂₂H₂₂O₂: 318.1620, found: 318.1620.

Received: October 2, 2007

Revised: December 6, 2007

Published online: January 29, 2008

Keywords: cyclization · domino reactions · furans · multicomponent reactions · palladium

- [1] For reviews, see: a) X. L. Hou, H. Y. Cheung, T. Y. Hon, P. L. Kwan, T. H. Lo, S. Y. T. Tong, H. N. C. Wong, *Tetrahedron* **1998**, 54, 1955; b) B. A. Keay, *Chem. Soc. Rev.* **1999**, 28, 209; c) T. L. Gilchrist, *J. Chem. Soc. Perkin Trans. 1* **1999**, 2849; d) X. L. Hou, Z. Yang, H. N. C. Wong in *Progress in Heterocyclic Chemistry*,

- Vol. 14 (Eds.: G. W. Gribble, T. L. Gilchrist), Pergamon, Oxford, **2002**, p. 139.
- [2] a) R. J. Sundberg in *Comprehensive Heterocyclic Chemistry*, Vol. 5 (Eds.: A. R. Katritzky, C. W. Rees), Pergamon, New York, **1984**, p. 313; b) H. Heaney in *Natural Products Chemistry* (Ed.: K. Nakanishi), Kodansha, Tokyo, **1974**, p. 297; c) B. H. Lipshutz, *Chem. Rev.* **1986**, 86, 795; d) T. H. Brown, M. A. Armitage, R. C. Blakemore, P. Blurton, G. J. Durant, C. R. Ganellin, R. G. Iffe, M. E. Parsons, D. A. Rawlings, B. P. Slingsby, *Eur. J. Med. Chem.* **1990**, 25, 217; e) M. Shipman, *Contemp. Org. Synth.* **1995**, 2, 1; f) L. A. Paquette, *Org. Lett.* **2000**, 2, 4095.
- [3] a) J. A. Marshall, E. D. Robinson, *J. Org. Chem.* **1990**, 55, 3450; b) J. A. Marshall, G. S. Bartley, *J. Org. Chem.* **1994**, 59, 7169; c) J. A. Marshall, E. M. Wallace, *J. Org. Chem.* **1995**, 60, 796; d) A. S. K. Hashmi, *Angew. Chem.* **1995**, 107, 1749; *Angew. Chem. Int. Ed. Engl.* **1995**, 34, 1581; e) A. S. K. Hashmi, T. L. Ruppert, T. Knofel, J. W. Bats, *J. Org. Chem.* **1997**, 62, 7295; f) A. S. K. Hashmi, L. Schwarz, J. H. Choi, T. M. Frost, *Angew. Chem.* **2000**, 112, 2382; *Angew. Chem. Int. Ed.* **2000**, 39, 2285; g) S. Ma, L. Li, *Org. Lett.* **2000**, 2, 941; h) S. Ma, Z. Yu, *Angew. Chem.* **2002**, 114, 1853; *Angew. Chem. Int. Ed.* **2002**, 41, 1775; i) S. Ma, J. Zhang, L. Lu, *Chem. Eur. J.* **2003**, 9, 2447; j) J. T. Kim, A. V. Kel'in, V. Gevorgyan, *Angew. Chem.* **2003**, 115, 102; *Angew. Chem. Int. Ed.* **2003**, 42, 98; k) A. W. Sromek, A. V. Kel'in, V. Gevorgyan, *Angew. Chem.* **2004**, 116, 2330; *Angew. Chem. Int. Ed.* **2004**, 43, 2280; l) M. H. Suhre, M. Reif, S. F. Kirsch, *Org. Lett.* **2005**, 7, 3925; m) A. W. Sromek, M. Rubina, V. Gevorgyan, *J. Am. Chem. Soc.* **2005**, 127, 10500; n) A. Dudnik, V. Gevorgyan, *Angew. Chem.* **2007**, 119, 5287; *Angew. Chem. Int. Ed.* **2007**, 46, 5195; o) T. Schwier, A. W. Sromek, D. M. L. Yap, D. Chernyak, V. Gevorgyan, *J. Am. Chem. Soc.* **2007**, 129, 9868.
- [4] a) Y. Fukuda, H. Shiragami, K. Utimoto, H. Nozaki, *J. Org. Chem.* **1991**, 56, 5816; b) A. V. Kel'in, V. Gevorgyan, *J. Org. Chem.* **2002**, 67, 95.
- [5] J. Zhang, H. G. Schmalz, *Angew. Chem.* **2006**, 118, 6856; *Angew. Chem. Int. Ed.* **2006**, 45, 6704.
- [6] Y. H. Liu, F. Song, Z. Song, M. Liu, B. Yan, *Org. Lett.* **2005**, 7, 5409.
- [7] a) T. Yao, X. Zhang, R. C. Larock, *J. Am. Chem. Soc.* **2004**, 126, 11164; b) T. Yao, X. Zhang, R. C. Larock, *J. Org. Chem.* **2005**, 70, 7679; c) Y. H. Liu, S. Zhou, *Org. Lett.* **2005**, 7, 4609; d) N. T. Patil, H. Wu, Y. Yamamoto, *J. Org. Chem.* **2005**, 70, 4531.
- [8] a) S. Ma, J. Zhang, *Angew. Chem.* **2003**, 115, 193; *Angew. Chem. Int. Ed.* **2003**, 42, 183; b) S. Ma, L. Lu, J. Zhang, *J. Am. Chem. Soc.* **2004**, 126, 9645.
- [9] a) A. Padwa, J. M. Kassir, S. L. Xu, *J. Org. Chem.* **1991**, 56, 6971; b) S. Ma, J. Zhang, *J. Am. Chem. Soc.* **2003**, 125, 12386.
- [10] For recent reviews, see: a) A. Dömling, *Curr. Opin. Chem. Biol.* **2002**, 6, 306; b) R. V. A. Orru, M. de Greef, *Synthesis* **2003**, 1471; c) C. Hulme, V. Gore, *Curr. Med. Chem.* **2003**, 10, 51; d) J. Zhu, *Eur. J. Org. Chem.* **2003**, 1133; e) V. Nair, C. Rajesh, A. U. Vinod, S. Bindu, A. R. Sreekanth, J. S. Mathen, L. Balagopal, *Acc. Chem. Res.* **2003**, 36, 899; f) D. J. Ramón, M. Yus, *Angew. Chem.* **2005**, 117, 1628; *Angew. Chem. Int. Ed.* **2005**, 44, 1602; g) A. Dondoni, A. Massi, *Acc. Chem. Res.* **2006**, 39, 451; h) A. Dömling, *Chem. Rev.* **2006**, 106, 17; i) D. M. D'Souza, T. J. J. Müller, *Chem. Soc. Rev.* **2007**, 36, 1095; j) T. J. J. Müller in *Topics in Organometallic Chemistry* (Ed. T. J. J. Müller), Springer, Berlin, **2006**, Vol. 19, pp. 149; k) *Multicomponent Reactions* (Eds.: J. Zhu, H. Bienaymé), Wiley-VCH, Weinheim, **2005**.
- [11] For recent examples of syntheses of tetrasubstituted furans, see: a) J. M. Aurrecoechea, E. Pérez, *Tetrahedron Lett.* **2001**, 42, 3839; b) J. Okauchi, T. Tanaka, T. Minami, *J. Org. Chem.* **2001**, 66, 3924; c) R. Sanz, D. Miguel, A. Martinez, J. M. Álvarez-Gutiérrez, F. Rodríguez, *Org. Lett.* **2007**, 9, 727; d) See references [3i,3k-o] and [7b and c].
- [12] For examples of Lewis acid catalyzed cyclization of similar systems, see references [5 and 6a].
- [13] Pd(OAc)₂ acts as both a Lewis acid and a transition metal, see: N. Asao, T. Nogami, K. Takahashi, Y. Yamamoto, *J. Am. Chem. Soc.* **2002**, 124, 764.
- [14] As suggested by one reviewer, the present data also support an alternative reaction mechanism that is depicted below. We want to thank the reviewer.

