


Synthetic Methods

Nickel-Catalyzed Reductive Cyclization of Unactivated 1,6-Enynes in the Presence of Organozinc Reagents**

Mao Chen, Yue Weng, Mian Guo, Hua Zhang, and Aiwen Lei*

The transition-metal-catalyzed cyclization of 1,6-enynes is one of the most convenient and efficient methods for the synthesis of a variety of carbocyclic and heterocyclic compounds.^[1–4] By this approach, enynes can be transformed into cyclic skeletons in a one-pot fashion with impressive regio- and stereoselectivity.^[3,5,6] The readily available and inexpensive metal nickel stands out for its remarkable power in catalyzing cycloisomerization and cycloaddition reactions of enynes.^[7–9] Important examples include a Ni-catalyzed [4+2] cycloaddition developed by Wender and co-workers^[10,11] and a series of cyclization reactions of activated 1,6-enynes described by Montgomery and co-workers [Scheme 1,




Scheme 1. Ni-mediated cyclization of enynes. binap = 2,2'-bis(diphenylphosphanyl)-1,1'-binaphthyl.

Eq. (1,2)).^[7,12–15] These strategies have established new approaches for the highly stereoselective formation of ring systems decorated with exocyclic trisubstituted or tetrasubstituted C=C bonds. Viable substrates for these processes, however, usually bear electron-withdrawing substituents. To our knowledge, very few studies have been described to date in which the potential of Ni catalysis in direct cyclizations of

electronically unactivated enynes has been explored [Scheme 1, Eq. (3)]. Such unactivated enynes have historically been used in combination with catalytic systems based on Pd,^[16–19] Au,^[20] Ru,^[21] and Rh.^[19,22–24] We and others have described highly enantioselective Rh-mediated cycloisomerization reactions of 1,6-enynes [Scheme 1, Eq. (3), path A].^[25–29] To further explore this cyclization chemistry, we have now investigated a Ni-based system with the aim of uncovering new reactivity and selectivity in a Ni-mediated Pauson–Khand-type cyclization under a reductive atmosphere, such as CO gas.^[30–34]

Unlike Ni^{II} species, most Ni⁰ complexes are extremely air-sensitive and are thus rather difficult to manipulate.^[35] Consequently, many known processes have resorted to the generation of these compounds in situ from air-stable Ni^{II} precursors in the presence of an appropriate reducing reagent, such as diisobutylaluminum hydride or a dialkyl zinc reagent.^[7,36] We initially set out to probe the feasibility of the Pauson–Khand reaction (PKR) of enyne **1a** in the presence of [Ni(PPh₃)₂Cl₂] (5 mol %) and *i*Pr₂Zn (30 mol %) under a CO atmosphere (balloon pressure).^[36,37] Despite extensive experimentation, this system failed to deliver any of the desired PKR product. However, a new compound was detected by serendipity when 3.0 equivalents of *i*Pr₂Zn were used. The compound was isolated and identified as **2a**, that is, a compound into which a molecule of dihydrogen has formally been incorporated following cyclization, rather than a molecule of CO as intended (Scheme 2).



Scheme 2. Ni-catalyzed cyclization of the 1,6-enyne **1a**.

Compound **3a** was established to be a superior substrate to **1a** upon subsequent optimization of the reaction and was therefore chosen as the substrate for the further screening of catalyst systems. The reactions were carried out at 40 °C for 18 h at a concentration of 0.1 M in THF. It was found that a combination of [Ni(acac)₂] (10 mol %) and *i*Pr₂Zn (3.0 equiv) furnished the product of reductive cyclization **4a** in 95 % yield with exclusive formation of the *Z* alkene (Table 1, entry 3). A decrease in the temperature, the catalyst loading, or the

[*] M. Chen, Y. Weng, M. Guo, H. Zhang, Prof. A. Lei
College of Chemistry and Molecular Sciences
Wuhan University
Wuhan, Hubei, 430072 (China)
Fax: (+86) 27-68754067
E-mail: aiwenlei@whu.edu.cn
Homepage: http://www.chem.whu.edu.cn/GCI_WebSite/main.htm

[**] This research was supported by the National Natural Science Foundation of China (20502020) and a startup fund from Wuhan University.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

Table 1: Ni-catalyzed reductive cyclization of **3a** under different conditions.^[a]

Entry	Catalyst (mol %)	Ligand (mol %)	<i>i</i> Pr ₂ Zn [equiv]	Yield [%]
1 ^[b]	[Ni(acac) ₂] (10)	—	3.0	64
2	[Ni(acac) ₂] (5)	—	3.0	69 ^[c]
3	[Ni(acac) ₂] (10)	—	3.0	95 ^[c]
4	[Ni(acac) ₂] (10)	—	2.0	64
5	[Ni(acac) ₂] (10)	—	1.0	40
6	[Ni(PPh ₃) ₂ Cl ₂] (10)	—	3.0	94 ^[c]
7	[Ni(PPh ₃) ₄] (10)	—	3.0	45
8	[Ni(acac) ₂] (10)	PPh ₃ (10)	3.0	94
9	[Ni(acac) ₂] (10)	PPh ₃ (20)	3.0	90
10	[Ni(acac) ₂] (10)	PPh ₃ (40)	3.0	85

[a] All reactions were carried out at a substrate concentration of 0.1 M for 18 h. The yields and the selectivities were determined by GC. [b] The reaction was carried out at 28 °C. [c] Yield of the isolated product. acac = acetylacetonate, Ts = *p*-toluenesulfonyl.

amount of *i*Pr₂Zn used resulted in a lower yield of **4a** (Table 1, entries 1, 2, 4, and 5). When [Ni(PPh₃)₄] was used as the catalyst, the product was formed in 45% yield. However, an almost identical result was observed with [Ni(PPh₃)₂Cl₂] (Table 1, entry 6) to that with [Ni(acac)₂]. Further experiments revealed that the presence of PPh₃ as a ligand in a 1:1 ratio with [Ni(acac)₂] did not effect the outcome of the reaction; at higher ratios of 2:1 or 4:1, PPh₃ inhibited the reaction to a certain extent (Table 1, entries 8–10).^[12,13,38]

A variety of N-tethered enynes were tested as substrates under the optimized reaction conditions (Table 2). In most cases, the desired cyclized product was generated with exceedingly high selectivity for the *Z*-configured alkene,^[39] as confirmed unambiguously by an NOE experiment. The steric and electronic properties of the substituent on the aryl group attached to the alkyne affected the reaction. Compound **3b** with a *p*-MeO group was transformed into the desired product in 89% yield with high selectivity for the *Z* alkene, whereas lower yields were observed for the reactions of **3c** and **3d**; the position of the methoxy substituent did not have an influence on the stereoselectivity (Table 2, entries 2–4). When an electron-withdrawing ester substituent was present, the product was obtained in 65% yield, and the selectivity dropped to *Z/E* 74:24 (Table 2, entry 5).^[40] When the aromatic substituent on the alkyne was exchanged for an alkyl group, the product was furnished in moderate yield but with a selectivity of greater than 99:1 (Table 2, entry 6). A substrate with an internal alkene in the side chain was transformed into **4g** in 66% yield. Compound **4j** was also obtained in 24% yield, and no cyclic product with an isomerized alkene side chain was detected (Table 2, entry 7). When the protecting group on the N tether was changed from a tosyl group to Boc (**3h**) or Cbz (**3i**), the catalyst system still afforded the cyclized products in satisfactory yields with high selectivity.

Table 2: Ni-catalyzed reductive cyclization reactions of compounds **3** to give pyrrolidine derivatives.^[a]

Entry	3	4	Yield [%] ^[b]	<i>Z/E</i> ^[c]
1	3a R ¹ = Ph	4a	95	> 99:1
2	3b R ¹ = Ph-4-OMe	4b	89	> 99:1
3	3c R ¹ = Ph-3-OMe	4c	56	> 99:1
4	3d R ¹ = Ph-2-OMe	4d	36	> 99:1
5	3e R ¹ = Ph-4-COOEt	4e	65	74:26
6	3f R ¹ = C ₅ H ₁₁	4f	62	> 99:1
7	3g	4g	66	> 99:1
8	3h	4h	72	> 99:1
9	3i	4i	66	> 99:1

[a] All reactions were carried out with [Ni(acac)₂] (10 mol %) and *i*Pr₂Zn (3.0 equiv) in THF at 40 °C for 18 h. [b] Yield of the isolated product. [c] The *Z/E* ratio was determined by GC. Boc = *tert*-butoxycarbonyl, Cbz = carbobenzyloxy.


To further explore the scope of application of this method, we tested the use of O-tethered enynes as substrates (Table 3). Good selectivity for the *Z* isomer and moderate yields were observed when substrates with an aryl or an alkyl group attached to the alkyne were used. Again, only when a *p*-COOEt group was present on the aromatic ring did the yield and selectivity drop (to 39% and 75:25; Table 3, entry 3). This result is similar to that observed with **3e**. Poor diastereoselectivity was observed for the reaction of **1a**, with a methyl substituent at the 5-position, but the *Z/E* ratio of the product was still greater than 99:1 (Table 3, entry 5).

Table 3: Ni-catalyzed reductive cyclization reactions of compounds **1** to give tetrahydrofuran derivatives **2**.^[a]

Entry	1	2	Yield [%] ^[b]	<i>Z/E</i> ^[c]
1	1b R ¹ = Ph	2b	66 (84)	> 99:1
2	1c R ¹ = Ph-4-OMe	2b	65 (77)	> 99:1
3 ^[d]	1d R ¹ = Ph-4-COOEt	2b	39	75:25
4	1e R ¹ = C ₅ H ₁₁	2b	65	> 99:1
5	1a	2b	72	> 99:1

[a] Reactions were carried out in the presence of *i*Pr₂Zn (3.0 equiv) and [Ni(acac)₂] (10 mol %) at 40 °C for 12 h. [b] Yield of the isolated product. Yields determined by GC are reported in parentheses. [c] The *Z/E* ratio was determined by GC. [d] Reaction time: 18 h.

A speculative catalytic cycle is shown in Scheme 3. Oxidative cyclization of the enyne to Ni^0 , derived from $[\text{Ni}(\text{acac})_2]$ through reduction by $i\text{Pr}_2\text{Zn}$,^[41] would afford a nickel metallocyclopentene intermediate **I**,^[42–44] which could undergo transmetalation with $i\text{Pr}_2\text{Zn}$ through two possible pathways to form intermediate **II** or **III**. Subsequent β -H elimination and reductive elimination would generate **VI** and




Scheme 3. Proposed mechanism for the reductive cyclization promoted by $[\text{Ni}(\text{acac})_2]$.

VII, respectively, both of which would be converted into the product of reductive cyclization upon hydrolysis. Montgomery et al. suggested the formation of five- or seven-membered nickel metallacycle intermediates through the oxidative cyclization of an alkynal or an alkynone to a Ni^0 species and subsequent generation of ring-opened intermediates by transmetalation between the nickel metallacycle and an organozinc reagent, in direct analogy to the formation of **II** and **III** in Scheme 3.^[7] Furthermore, Knochel and co-workers have demonstrated such an exchange between organonickel intermediates and Et_2Zn .^[45–49] The palladium-catalyzed reductive cyclization of carbon-tethered 1,6-enynes in the presence of a silicon hydride and HOAc provided similar products, but the mechanism was different from our hypothesis.^[50]

To gain more information about the reaction mechanism, we quenched the reaction of **3a** with CD_3COOD .^[12] Compound **5**, with the deuterium label on the $\text{C}=\text{C}$ bond, and compound **6**, with a deuterium-labeled methyl group, were obtained as 53 and 19% of the product mixture, respectively (Scheme 4), a result that indicates the existence of both **VI** and **VII** as intermediates. Moreover, the high selectivity for the formation of products with the *Z* configuration is also consistent with the formation of a Ni metallacycle. It is still unclear why we have never observed alkylated cyclization products. Such products were reported to be dominant when electron-deficient enynes were used as substrates.

In conclusion, we have developed an inexpensive $[\text{Ni}(\text{acac})_2]$ -catalyzed reductive cyclization of unactivated 1,6-enynes in the presence of $i\text{Pr}_2\text{Zn}$ under mild conditions. Through a simple procedure, functionalized pyrrolidine and tetrahydrofuran derivatives were produced in a generally



Scheme 4. Confirmation of the proposed mechanism by a quenching experiment with CD_3COOD .

highly stereoselective manner in moderate to high yields. Our proposed mechanism was confirmed through a quenching experiment with CD_3COOD . Studies into the scope of the reaction and mechanistic studies are under way.

Experimental Section

Typical procedure: $i\text{Pr}_2\text{Zn}$ (6 mL, 0.5 M in THF) was added to a solution of **3b** (355 mg, 1.0 mmol) and $[\text{Ni}(\text{acac})_2]$ (25.6 mg, 0.1 mmol) in anhydrous THF (4 mL) at -78°C . The reaction mixture was stirred at 40°C for 18 h under N_2 , and then 2 M HCl (25 mL) was added to quench the reaction. The mixture was extracted with dichloromethane, and the combined organic layers were washed with saturated NaHCO_3 and brine, dried (Na_2SO_4), and then concentrated in vacuo. The residue was purified by column chromatography over silica gel (ethyl acetate/petroleum 1:20, v/v) to give **4b** (318 mg, 89%) as a colorless oil. ^1H NMR (300 MHz, CDCl_3): δ = 7.63 (d, J = 8.1 Hz, 2H), 7.20 (d, J = 7.8 Hz, 2H), 6.98 (d, J = 8.4 Hz, 2H), 6.78 (d, J = 8.7 Hz, 2H), 6.06 (s, 1H), 4.12 (d, J = 14.7 Hz, 1H), 3.93 (d, J = 14.4 Hz, 1H), 3.70 (s, 3H), 3.44 (t, J = 7.8 Hz, 1H), 2.76 (m, 1H), 2.62 (t, J = 8.4 Hz, 1H), 2.30 (s, 3H), 1.05 ppm (d, J = 6.6 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3): δ = 158.7, 143.8, 139.6, 133.0, 129.9, 129.7, 129.5, 127.9, 121.6, 114.1, 55.4, 54.1, 51.0, 39.1, 21.7, 17.1 ppm; HRMS (APCI): calcd for $\text{C}_{20}\text{H}_{23}\text{NO}_3\text{S}$: 357.1399 $[M]^+$; found: 357.1404.

Received: September 27, 2007

Revised: December 21, 2007

Published online: February 8, 2008

Keywords: cyclization · enynes · nickel · organozinc reagents · synthetic methods

- [1] B. M. Trost, *Science* **1991**, 254, 1471–1477.
- [2] B. M. Trost, *Angew. Chem.* **1995**, 107, 285–307; *Angew. Chem. Int. Ed. Engl.* **1995**, 34, 259–281.
- [3] M. Lautens, W. Klute, W. Tam, *Chem. Rev.* **1996**, 96, 49–92.
- [4] E.-i. Negishi, C. Copéret, S. Ma, S.-Y. Liou, F. Liu, *Chem. Rev.* **1996**, 96, 365–393.
- [5] C. Aubert, O. Buisine, M. Malacria, *Chem. Rev.* **2002**, 102, 813–834.
- [6] A. B. Dounay, L. E. Overman, *Chem. Rev.* **2003**, 103, 2945–2964.
- [7] J. Montgomery, *Angew. Chem.* **2004**, 116, 3980–3998; *Angew. Chem. Int. Ed.* **2004**, 43, 3890–3908.
- [8] K. Fischer, K. Jonas, P. Misbach, R. Stabba, G. Wilke, *Angew. Chem.* **1973**, 85, 1001–1012; *Angew. Chem. Int. Ed. Engl.* **1973**, 12, 943–953.
- [9] G. Wilke, *Angew. Chem.* **1988**, 100, 189–211; *Angew. Chem. Int. Ed. Engl.* **1988**, 27, 185–206.
- [10] P. A. Wender, T. E. Jenkins, *J. Am. Chem. Soc.* **1989**, 111, 6432–6434.

- [11] P. A. Wender, T. E. Smith, *J. Org. Chem.* **1996**, *61*, 824–825.
- [12] J. Montgomery, A. V. Savchenko, *J. Am. Chem. Soc.* **1996**, *118*, 2099–2100.
- [13] J. Montgomery, E. Oblinger, A. V. Savchenko, *J. Am. Chem. Soc.* **1997**, *119*, 4911–4920.
- [14] J. Seo, H. Fain, J.-B. Blanc, J. Montgomery, *J. Org. Chem.* **1999**, *64*, 6060–6065.
- [15] X.-Q. Tang, J. Montgomery, *J. Am. Chem. Soc.* **1999**, *121*, 6098–6099.
- [16] B. Radetich, T. V. RajanBabu, *J. Am. Chem. Soc.* **1998**, *120*, 8007–8008.
- [17] C. H. Oh, H. H. Jung, J. S. Kim, S. W. Cho, *Angew. Chem.* **2000**, *112*, 768–771; *Angew. Chem. Int. Ed.* **2000**, *39*, 752–755.
- [18] B. M. Trost, Y. Shi, *J. Am. Chem. Soc.* **1993**, *115*, 9421–9438.
- [19] I. J. S. Fairlamb, *Angew. Chem.* **2004**, *116*, 1066–1070; *Angew. Chem. Int. Ed.* **2004**, *43*, 1048–1052.
- [20] S. Wang, L. Zhang, *J. Am. Chem. Soc.* **2006**, *128*, 14274–14275.
- [21] B. Schmidt, *Angew. Chem.* **2003**, *115*, 5146–5149; *Angew. Chem. Int. Ed.* **2003**, *42*, 4996–4999.
- [22] H.-Y. Jang, F. W. Hughes, H. Gong, J. Zhang, J. S. Brodbelt, M. J. Krische, *J. Am. Chem. Soc.* **2005**, *127*, 6174–6175.
- [23] B. M. Trost, A. McClory, *Angew. Chem.* **2007**, *119*, 2120–2123; *Angew. Chem. Int. Ed.* **2007**, *46*, 2074–2077.
- [24] Y. Oonishi, A. Taniuchi, M. Mori, Y. Sato, *Tetrahedron Lett.* **2006**, *47*, 5617–5621.
- [25] A. Lei, M. He, X. Zhang, *J. Am. Chem. Soc.* **2003**, *125*, 11472–11473.
- [26] X. Tong, D. Li, Z. Zhang, X. Zhang, *J. Am. Chem. Soc.* **2004**, *126*, 7601–7607.
- [27] P. Cao, B. Wang, X. Zhang, *J. Am. Chem. Soc.* **2000**, *122*, 6490–6491.
- [28] A. Lei, M. He, S. Wu, X. Zhang, *Angew. Chem.* **2002**, *114*, 3607–3610; *Angew. Chem. Int. Ed.* **2002**, *41*, 3457–3460.
- [29] A. Lei, J. P. Waldkirch, M. He, X. Zhang, *Angew. Chem.* **2002**, *114*, 4708–4711; *Angew. Chem. Int. Ed.* **2002**, *41*, 4526–4529.
- [30] S. E. Gibson (née Thomas), A. Stevenazzi, *Angew. Chem.* **2003**, *115*, 1844–1854; *Angew. Chem. Int. Ed.* **2003**, *42*, 1800–1810.
- [31] O. Geis, H.-G. Schmalz, *Angew. Chem.* **1998**, *110*, 955–958; *Angew. Chem. Int. Ed.* **1998**, *37*, 911–914.
- [32] M. Rodríguez Rivero, J. Adrio, J. C. Carretero, *Eur. J. Org. Chem.* **2002**, 2881–2889.
- [33] K. H. Park, Y. K. Chung, *Synlett* **2005**, 545–559.
- [34] G. Garcia-Gomez, J. M. Moreto, *J. Am. Chem. Soc.* **1999**, *121*, 878–879.
- [35] D. J. Krysan, P. B. Mackenzie, *J. Org. Chem.* **1990**, *55*, 4229–4230.
- [36] P. Knochel, A. Boudier, L. O. Bromm, E. Hupe, J. A. Varela, A. Rodriguez, C. Koradin, T. Bunlaksananusorn, H. Laaziri, F. Lhermitte, *Pure Appl. Chem.* **2000**, *72*, 1699–1703.
- [37] E. Negishi, *Pure Appl. Chem.* **1992**, *64*, 323–334.
- [38] Y. Sato, M. Takimoto, M. Mori, *J. Am. Chem. Soc.* **2000**, *122*, 1624–1634.
- [39] When the reactions were monitored by TLC, we observed only the spot of the desired product and a spot at the baseline, which we suspect to be an oligomeric product. Oligomerization may be the reason behind the isolation of the desired product in low yields in entries 3 and 4 of Table 2.
- [40] We do not have enough data as yet to explain the difference between the selectivity of the reaction of **3e** and that of the reactions of substrates **3a–3d** and **3f–3h**.
- [41] When [Ni(cod)₂] (10 mol%) was employed as the catalyst precursor (cod = 1,5-cyclooctadiene), the reductive cyclization of **3a** was equally efficient. The desired product was isolated in 85% yield.
- [42] R. H. Grubbs, A. Miyashita, *J. Am. Chem. Soc.* **1978**, *100*, 1300–1302.
- [43] R. H. Grubbs, A. Miyashita, M. Liu, P. Burk, *J. Am. Chem. Soc.* **1978**, *100*, 2418–2425.
- [44] R. J. McKinney, D. L. Thorn, R. Hoffmann, A. Stockis, *J. Am. Chem. Soc.* **1981**, *103*, 2595–2603.
- [45] A. Vaupel, P. Knochel, *Tetrahedron Lett.* **1994**, *35*, 8349–8352.
- [46] S. Vettel, A. Vaupel, P. Knochel, *Tetrahedron Lett.* **1995**, *36*, 1023–1026.
- [47] H. Stadtmüller, A. Vaupel, C. E. Tucker, T. Stuedemann, P. Knochel, *Chem. Eur. J.* **1996**, *2*, 1204–1220.
- [48] A. Vaupel, P. Knochel, *J. Org. Chem.* **1996**, *61*, 5743–5753.
- [49] S. Vettel, A. Vaupel, P. Knochel, *J. Org. Chem.* **1996**, *61*, 7473–7481.
- [50] B. M. Trost, F. Rise, *J. Am. Chem. Soc.* **1987**, *109*, 3161–3163.