


Asymmetric Iminium Ion Catalysis: An Efficient Enantioselective Synthesis of Pyranonaphthoquinones and β -Lapachones**

Magnus Rueping,* Erli Sugiono, and Estíbaliz Merino

Naphthoquinones are common in the plant kingdom. As a result of their molecular structure they exhibit redox properties that influence various regulatory cellular processes. Their cytotoxic activity is a result of the quinone core structure, which is an important pharmacophore.^[1] Furthermore, a number of naturally occurring 1,4- and 1,2-naphthoquinones are associated with diverse biological activities^[2] and are components of antibacterial, fungicidal, antimalarial, anti-parasitic, and antitumoral agents.^[3] Based on their biological and structural properties, the 1,2- and 1,4-naphthoquinones are considered privileged structures in medicinal chemistry.

Following previous studies on the activation of aldimines and carbonyl functionalities with chiral Brønsted acids,^[4] we decided to examine an enantioselective Brønsted acid catalyzed synthesis of 1,2-pyranonaphthoquinones starting from hydroxynaphthoquinone and α,β -unsaturated aldehydes [Eq. (1)]. However, in initial experiments we did not obtain satisfactory results with regard to reactivity and selectivity.



Therefore, we decided to investigate an alternative reaction with a chiral secondary amine as the organocatalyst. In contrast to primary amines previously employed, which are known to lead to the condensation product,^[5] we assumed that it should be feasible to activate α,β -unsaturated aldehydes **2** using secondary amines through the formation of an intermediary iminium ion.^[6] The subsequent 1,4-addition reaction to 2-hydroxy-1,4-naphthoquinone **1**, followed by an acetalization should give the corresponding enantiomerically enriched 1,4-pyranonaphthoquinones **3**. The initial experiments showed that the reaction of **1** with the α,β -unsaturated

aldehyde **2a** could be carried out in the presence of diarylprolinol ethers^[7] **4a** and **4b** in DMSO to give the corresponding 1,2-pyranonaphthoquinone **3a** in good yields and excellent enantioselectivities (Table 1, entries 1 and 2).^[8] This new

Table 1: Solvent survey of the diarylprolinol ether catalyzed enantioselective synthesis of 1,4-naphthoquinones.^[a]

Entry ^[b]	T [°C]	Catalyst	Solvent	Yield [%] ^[d]	ee [%] ^[e]
1	RT	4a	DMSO	86	82
2	RT	4b	DMSO	78	98
3	0	4a	Et ₂ O	48	90
4	0	4a	Bu ₂ O	24	85
5	0	4a	CH ₂ Cl ₂	71	90
6	0	4a	toluene	54	66
7 ^[c]	−20	4a	CH ₂ Cl ₂	76	92
8 ^[c]	−20	4b	CH ₂ Cl ₂	77	99
9	−20	4a	toluene	34	92
10 ^[c]	−20	4b	toluene	69	98

[a] TMS = trimethylsilyl. [b] Reactions were performed with hydroxyquinone **1**, aldehyde **2a** (1.5 equiv) and 20 mol% **4** for 20 h. [c] 40 h. [d] Yield of isolated product after column chromatography. [e] Enantiomeric excess was determined by HPLC.

asymmetric addition–cyclization cascade^[9] can also be performed in other solvents, whereby the best enantioselectivities (up to 99% ee) and yields were obtained with dichloromethane as the solvent in combination with catalyst **4b** (Table 1, entry 8).

Further reaction optimization focused on the solvent concentration and catalyst loading of **4a** and **4b** (Table 2). While lower catalyst loadings resulted in slightly reduced enantioselectivities, the variation of the concentration did not have any significant influence (Table 2, entries 1–5). Again catalyst **4b** gave the best results when the reaction was carried out in dichloromethane (0.2 M **1**, 0.2 M **2a**) (Table 2, entries 6–8).


Using the optimized reaction conditions we investigated the scope of the diarylprolinol ether catalyzed enantioselective addition–cyclization reaction cascade using various α,β -unsaturated aldehydes **2** (Table 3). In general, aliphatic (Table 3, entries 1–6) as well as aromatic (Table 3, entries 7–15) α,β -unsaturated aldehydes could be employed successfully in this new transformation, and a diverse set of 1,4-

[*] Prof. Dr. M. Rueping, Dr. E. Sugiono, Dr. E. Merino
Degussa Endowed Professorship
Institute for Organic Chemistry and Chemical Biology
Johann Wolfgang Goethe-University Frankfurt am Main
Max-von-Laue Strasse 7, 60438 Frankfurt am Main (Germany)
Fax: (+49) 69-798-29248
E-mail: M.rueping@chemie.uni-frankfurt.de

[**] The authors acknowledge Degussa GmbH and the DFG (Priority Programme Organocatalysis) for financial support as well the Ministerio de Educación y Ciencia, Spain, for a stipend given to E.M. and Dr. M. Bolte for the X-ray crystal structure analysis.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.


Table 2: Effect of catalyst loading and concentration of **1**.



Entry ^[a]	Catalyst (mol %)	Concentration	Yield [%] ^[b]	ee [%] ^[c]
1	4a (5)	0.2 M	47	87
2	4a (10)	0.2 M	76	88
3	4a (20)	0.1 M	53	90
4	4a (20)	0.2 M	76	92
5	4a (20)	0.3 M	69	91
6	4b (5)	0.2 M	40	99
7	4b (10)	0.2 M	41	99
8	4b (20)	0.2 M	77	99

[a] Reactions were performed with hydroxyquinone **1** and aldehyde **2a** (1.5 equiv). [b] Yield of isolated product after column chromatography. [c] Enantiomeric excess was determined by HPLC using a Chiralpak AD-H column.

Table 3: Scope of the diarylprolinol-catalyzed addition–cyclization cascade reaction.




Entry ^[a]	Product	R	T [°C]	Yield [%] ^[e]	ee [%] ^[f]
1	3a ^[b]	CH(CH ₃) ₂	−20	77	99
2	3b	C ₂ H ₅	−20	55	98
3	3c	C ₃ H ₇	−20	75	99
4	3d	C ₄ H ₉	−20	52	98
5	3e	C ₇ H ₁₅	−20	46	98
6	3f	C ₁₀ H ₂₁	−20	43	99
7	3g	Ph	−20	56	91
8	3h ^[c]	2-ClC ₆ H ₄	RT	50	99
9	3h ^[d]	2-ClC ₆ H ₄	RT	57	99
10	3i ^[d]	2-BrC ₆ H ₄	RT	84	99
11	3j ^[d]	2-CH ₃ C ₆ H ₄	RT	57	96
12	3k ^[d]	2-CF ₃ C ₆ H ₄	RT	49	95
13	3l	3-BrC ₆ H ₄	0	82	90
14	3m	4-BrC ₆ H ₄	0	87	92
15	3n	4-CH ₃ OC ₆ H ₄	0	51	93

[a] Reactions were performed with hydroxyquinone **1**, aldehyde **2** (1.3 equiv), and 20 mol % **4b** for 3 d. [b] 40 h. [c] 24 h. [d] Performed with 20 mol % **4a** at RT for 24 h. [e] Yield of isolated product after column chromatography. [f] Enantiomeric excess was determined by HPLC on a chiral stationary phase.

pyranonaphthoquinones **3a–n** was isolated in good yields and with excellent enantioselectivities (90–99 % ee).

Having developed an efficient and highly enantioselective method for the synthesis of 1,4-pyranonaphthoquinones **3**, we decided to transform the new products to the desired biologically active 1,2-pyranonaphthoquinones **5** (Table 4). It was thereby shown that application of sodium borohydride and concentrated hydrochloric or sulfuric acid resulted in the corresponding β -lapachone derivatives **5** without loss of enantiopurity.^[10] The 1,4-naphthoquinone **3** can also be


Table 4: Transformation of 1,4-naphthoquinones to 1,2-naphthoquinones.




Entry	Product	R	Yield [%]	ee [%]
5b	5b	C ₂ H ₅	98%	62%
5c	5c	C ₃ H ₇	98%	75%
5d	5d	C ₄ H ₉	98%	49%
5e	5e	C ₇ H ₁₅	98%	85%
5f	5f	C ₁₀ H ₂₁	99%	55%
5g	5g	Ph	91%	58%
5h	5h	2-ClC ₆ H ₄	99%	90%
5j	5j	2-CH ₃ C ₆ H ₄	94%	43%
5m	5m	3-BrC ₆ H ₄	92%	66%
5n	5n	4-CH ₃ OC ₆ H ₄	90%	61%

transformed to further interesting derivatives. For instance, oxidation of **3a** with PCC provides the corresponding lactone **6a**, which again can be isolated without loss of enantiopurity (Scheme 1).^[10]

With regard to the mechanism, we assume that the reaction of the diphenylprolinol ether **4** with the α,β -unsaturated aldehyde **2** results in the intermediary iminium ion **A** (Scheme 2). Subsequent 1,4-addition to 2-hydroxy-1,4-naphthoquinone (**1**) followed by an isomerization gives rise to adduct **B**, which after hydrolysis and acetylation yields the desired 1,4-naphthoquinone **3** with regeneration of the catalyst.


Scheme 1. Oxidation of 1,4-pyranonaphthoquinone **3a**. PCC = pyridinium chlorochromate.



Scheme 2. Diarylprolinol ether catalyzed reaction of 2-hydroxy-1,4-naphthoquinone (**1**) with α,β -unsaturated aldehyde **2**.

The constitution and absolute configuration of the new products was proven by X-ray crystal structure analysis, as we were able to obtain suitable single crystals of the 1,4-naphthoquinone **3j** as well as the 1,2-naphthoquinone **5c** (Figure 1).




Figure 1. X-ray crystal structures of 1,4-naphthoquinone **3j** and 1,2-naphthoquinone **5c** (ellipsoids drawn at the 50% probability level).

In summary we have developed a Lewis base catalyzed enantioselective addition cyclization cascade which results in biologically interesting quinones. In this efficient transformation not only aliphatic but also aromatic α,β -unsaturated aldehydes can be applied to provide 1,4-pyranonaphthoquinones in good yields and with excellent enantioselectivities (90–99% *ee*). The 1,4-naphthoquinones obtained can be further reacted to provide the valuable 1,2-pyranonaphthoquinones and lapachones as well as useful 1,4-naphthoquinonolactones. These transformations proceed without loss of enantiopurity. In current work we are subjecting further substrates to this efficient transformation.

Received: November 5, 2007
Published online: March 10, 2008

Keywords: iminium ions · lapachones · Lewis base catalysis · naphthoquinones · organocatalysis

- [1] J. S. Driscoll, G. F. Hazard, H. B. Wood, A. Goldin, *Cancer Chemother. Rep. Part 2* **1974**, 4, 1–362.
- [2] A. Ravelo, A. Estévez-Braun, H. Chávez-Orellana, E. Pérez-Sacau, D. Mesa-Silverio, *Curr. Top. Med. Chem.* **2004**, 4, 241.
- [3] a) P. Guiraud, R. Steiman, G. M. Campos-Takaki, F. Seigle-Murandi, M. S. de Buochberg, *Planta Med.* **1994**, 60, 373; b) V. F. de Andrade-Neto, M. O. F. Goulart, J. F. da Silva Filho, M. J. da Silva, M. C. F. R. Pinto, A. V. Pinto, M. G. Zalis, L. H. Carvalho, A. U. Krettli, *Bioorg. Med. Chem. Lett.* **2004**, 14, 1145; c) E. Pérez-Sacau, A. Estévez-Braun, A. G. Ravelo, D. Gutiérrez Yapu, A. Giménez Turba, *Chem. Biodiversity* **2005**, 2, 264; d) M. O. Goulart, C. I. Zani, J. Tonholo, F. C. Freitas de Abreu, A. B. Oliveira, D. S. Raaslan, S. Starling, E. Chiari, *Bioorg. Med. Chem. Lett.* **1997**, 7, 2043; e) A. Fujiwara, T. Mori, A. Lida, S. Ueda, T. Nomura, H. Tokuda, H. Nishino, *J. Nat. Prod.* **1998**, 61, 629; f) P. Krishnan, K. F. Bastow, *Biochem. Pharmacol.* **2000**, 60, 1367; g) E. Pérez-Sacau, A. Estévez-Braun, A. G. Ravelo, E. A. Ferro, H. Tokuda, T. Mukainaka, H. Nishino, *Bioorg. Med. Chem.* **2003**, 11, 483; h) M. Suzuki, M. Amano, J. Choi, H. J. Park, B. W. Williams, K. Ono, C. W. Song, *Radiat. Res.* **2006**, 165, 525; i) J. I. Lee, D. Y. Choi, H. S. Chung, H. G. Seo, H. J. Woo, B. T. Choi, Y. H. Choi, *Exp. Oncol.* **2006**, 28, 30; j) H. J. Woo, K.-Y. Park, C.-H. Rhu, W. H. Lee, B. T. Choi, G. Y. Kim, Y.-M. Park, Y. H. Choi, *J. Med. Food* **2006**, 9, 161.
- [4] a) M. Rueping, E. Sugiono, C. Azap, T. Theissmann, M. Bolte, *Org. Lett.* **2005**, 7, 3781; b) M. Rueping, A. P. Antonchick, T. Theissmann, *Angew. Chem.* **2006**, 118, 3765; *Angew. Chem. Int. Ed.* **2006**, 45, 3683; c) M. Rueping, A. P. Antonchick, T. Theissmann, *Angew. Chem.* **2006**, 118, 6903; *Angew. Chem. Int. Ed.* **2006**, 45, 6751; d) M. Rueping, E. Sugiono, C. Azap, *Angew. Chem.* **2006**, 118, 2679; *Angew. Chem. Int. Ed.* **2006**, 45, 2617; e) M. Rueping, C. Azap, *Angew. Chem.* **2006**, 118, 7996; *Angew. Chem. Int. Ed.* **2006**, 45, 7832; f) M. Rueping, E. Sugiono, F. R. Schoepke, *Synlett* **2007**, 144; g) M. Rueping, W. Ieawsuwan, A. P. Antonchick, B. J. Nachtsheim, *Angew. Chem.* **2007**, 119, 2143; *Angew. Chem. Int. Ed.* **2007**, 46, 2097; h) M. Rueping, A. P. Antonchick, *Angew. Chem.* **2007**, 119, 4646; *Angew. Chem. Int. Ed.* **2007**, 46, 4562; *Angew. Chem. Int. Ed.* **2007**, 46, 4562; i) M. Rueping, A. P. Antonchick, C. Brinkmann, *Angew. Chem.* **2007**, 119, 7027; *Angew. Chem. Int. Ed.* **2007**, 46, 6903.
- [5] Ethylendiamine diacetate and β -alanine were successfully applied in the condensation reaction of 2-hydroxy-1,4-naphthoquinone with α,β -unsaturated aldehydes: a) J. P. Malerich, T. J. Maimone, G. I. Elliott, D. Trauner, *J. Am. Chem. Soc.* **2005**, 127, 6276; b) Y. R. Lee, J. H. Choi, D. T. L. Trinh, N. W. Kim, *Synthesis* **2005**, 3026.
- [6] Review: G. Lelais, D. W. C. MacMillan, *Aldrichimica Acta* **2006**, 39, 79.
- [7] Examples for the use of diarylprolinol ethers in asymmetric catalysis: a) M. Marigo, T. C. Wabnitz, D. Fielenbach, K. A. Jørgensen, *Angew. Chem.* **2005**, 117, 804; *Angew. Chem. Int. Ed.* **2005**, 44, 794; b) M. Marigo, D. Fielenbach, A. Branton, A. Kjærsgaard, K. A. Jørgensen, *Angew. Chem.* **2005**, 117, 3769; *Angew. Chem. Int. Ed.* **2005**, 44, 3703; c) J. Franzen, M. Marigo, D. Fielenbach, T. C. Wabnitz, A. Kjærsgaard, K. A. Jørgensen, *J. Am. Chem. Soc.* **2005**, 127, 18296; d) Y. Hayashi, H. Gotoh, T. Hayashi, M. Shoji, *Angew. Chem.* **2005**, 117, 4284; *Angew. Chem. Int. Ed.* **2005**, 44, 4212; e) M. Marigo, J. Franzen, T. B. Poulsen, W. Zhuang, K. A. Jørgensen, *J. Am. Chem. Soc.* **2005**, 127, 6964; f) Y. G. Chi, S. H. Gellman, *Org. Lett.* **2005**, 7, 4253; g) D. Enders, M. R. M. Hüttl, C. Grondal, G. Raabe, *Nature* **2006**, 441, 861; h) W. Wang, H. Li, J. Wang, L. Zu, *J. Am. Chem. Soc.* **2006**, 128, 10354; i) A. Carlone, M. Marigo, C. North, A. Landa, K. A. Jørgensen, *Chem. Commun.* **2006**, 4928; j) I. Ibrahim, A. Córdova, *Chem. Commun.* **2006**, 1760; k) S. Bertelsen, P. Diner, R. L. Johansen, K. A. Jørgensen, *J. Am. Chem. Soc.*

- 2007, 129, 1536; l) J. Vesely, I. Ibrahim, G. L. Zhao, R. Rios, A. Córdova, *Angew. Chem.* **2007**, 119, 792; *Angew. Chem. Int. Ed.* **2007**, 46, 778; m) D. Enders, M. R. M. Hüttl, J. Runsink, G. Raabe, B. Wendt, *Angew. Chem.* **2007**, 119, 471; *Angew. Chem. Int. Ed.* **2007**, 46, 467; n) L. Zu, H. Li, H. Xie, J. Wang, W. Jiang, Y. Tang, W. Wang, *Angew. Chem.* **2007**, 119, 3806; *Angew. Chem. Int. Ed.* **2007**, 46, 3732; o) A. Carlone, S. Cabrera, M. Marigo, K. A. Jørgensen, *Angew. Chem.* **2007**, 119, 1119; *Angew. Chem. Int. Ed.* **2007**, 46, 1101; p) Y. Hayashi, T. Okano, S. Aratake, D. Hazeldard, *Angew. Chem.* **2007**, 119, 5010; *Angew. Chem. Int. Ed.* **2007**, 46, 4922; q) J. Alemán, S. Cabrera, E. Maerten, J. Overgaard, K. A. Jørgensen, *Angew. Chem.* **2007**, 119, 5616; *Angew. Chem. Int. Ed.* **2007**, 46, 5520. Review on the use of diarylprolinol ethers: r) C. Palomo, A. Mielgo, *Angew. Chem.* **2006**, 118, 8042; *Angew. Chem. Int. Ed.* **2006**, 45, 7876.
- [8] For an interesting review on the application of organocatalysis in natural product synthesis, see: R. M. de Figueiredo, M. Christmann, *Eur. J. Org. Chem.* **2007**, 2575.
- [9] For an excellent review on cascade reactions, see: D. Enders, C. Grondal, M. R. M. Hüttl, *Angew. Chem.* **2007**, 119, 1590; *Angew. Chem. Int. Ed.* **2007**, 46, 1570.
- [10] A detailed description is given in the Supporting Information.