

Chiral Phosphoric Acid Catalyzed Enantioselective Friedel–Crafts Alkylation of Indoles with Nitroalkenes: Cooperative Effect of 3 Å Molecular Sieves**


Junji Itoh, Kohei Fuchibe, and Takahiko Akiyama*

The enantioselective Friedel–Crafts alkylation^[1] of indoles with nitroalkenes is one of the most important carbon–carbon bond-forming reactions for the preparation of biologically active compounds such as indole alkaloids.^[2] Although several chiral catalysts, including salen/AlCl complexes,^[3a] bis(sulfonamide) compounds,^[3b] thiourea-containing structures,^[3c,f] and Zn^{II} or Cu^{II}–bisoxazoline complexes,^[3d,e,g] have been reported for the reaction, the enantioselectivities are not always satisfactory. The development of more efficient catalysts is required.


Recently, novel Brønsted acid catalysts were developed and used in a number of asymmetric reactions,^[4] and the chiral phosphoric acids derived from (*R*)-binol (binol = 2,2'-dihydroxy-1,1'-binaphthyl) have been extensively studied as versatile organocatalysts for enantioselective reactions.^[5–7] Most of the phosphoric acid catalyzed reactions reported so far involve imine or iminium ion electrophiles, however, there are scattered examples that include carbonyl compounds^[8a,b,9h,j] aziridines,^[8c] or nitrones as electrophiles.^[8d] The application of chiral phosphoric acid catalysis to other electrophiles is desirable, and the enantioselective addition of a nucleophile to a nitroalkene by using a chiral phosphoric acid has not previously been reported. We describe herein the first chiral phosphoric acid catalyzed Friedel–Crafts alkylation of indoles^[9] with nitroalkenes to afford Friedel–Crafts adducts in excellent enantioselectivities (Scheme 1).

Chiral phosphoric acid (*R*)-**3** provided the best enantioselectivity in the Friedel–Crafts alkylation of indole **1a** with nitroalkene **2a** (2 equiv) in benzene/1,2-dichloroethane (1:1) at –35 °C. Although the system resulted in the formation of Friedel–Crafts adduct **4a** with good enantioselectivity, the

Previous Studies



This Work



Scheme 1. Comparison of reactions from previous studies and this work.

yield was low (Table 1, entry 1).^[10] To our delight, we found that the addition of activated, powdered molecular sieves (M.S.) significantly improved both the chemical yield and the enantioselectivity. Among the molecular sieves examined, the 3 Å molecular sieves gave the best result (Table 1, entry 2).^[11] Since the addition of a small amount of water to the reaction mixture deteriorates the chemical yield (Table 1, entry 5),^[12,13] we suppose that the molecular sieves absorb the small amount of water present in the reaction medium.

We explored the scope and limitations of the reaction by using various indoles and nitroalkenes under the optimized reaction conditions, and the results are summarized in Table 2. A wide range of nitroalkenes bearing electron-donating (**2b–c**), electron-withdrawing (**2d–e**), and hetero-

Table 1: Effect of the 3 Å molecular sieves.^[a]

Entry	M.S.	Yield [%] ^[b]	ee [%] ^[c]
1		5	75
2	3 Å	76	91
3	4 Å	69	83
4	5 Å	59	88
5	3 Å ^[d]	1	60

[a] Reactions were carried out with **1a** (0.2 mmol) and **2a** (0.4 mmol; 2 equiv) in benzene (0.5 mL)/1,2-dichloroethane (0.5 mL). [b] Yield of isolated product. [c] Enantiomeric excess was determined by HPLC analysis. [d] Used 10 equiv of H₂O.

[*] Dr. J. Itoh, Dr. K. Fuchibe,^[†] Prof. Dr. T. Akiyama
Department of Chemistry, Faculty of Science
Gakushuin University
1-5-1 Mejiro, Toshima-ku, Tokyo 171-8588 (Japan)
Fax: (+81) 3-5992-1029
E-mail: takahiko.akiyama@gakushuin.ac.jp

[†] Present address: Department of Chemistry
Graduate School of Pure and Applied Sciences
University of Tsukuba
Tsukuba (Japan)

[**] This work was partially supported by a Grant-in Aid for Scientific Research on Priority Areas “Advanced Molecular Transformation of Carbon Resources” from the Ministry of Education, Science, Sports, Culture, and Technology (Japan). J.I. thanks the JSPS Research Fellowships for Young Scientists.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

aromatic (**2f**) groups underwent the Friedel–Crafts alkylation reaction to afford Friedel–Crafts adducts **4** with excellent enantioselectivities (Table 2, entries 1–6). Aliphatic nitroalkenes (**2g–j**) gave the corresponding adducts with high

Table 2: Chiral Brønsted acid catalyzed Friedel–Crafts alkylation of indoles with nitroalkenes.^[a]

Entry	R ¹	R ²	t [h]	Yield [%] ^[b]	ee [%] ^[c]
1	H (1a)	Ph (2a)	48	76 (4a)	91
2	H (1a)	4-MeC ₆ H ₄ (2b)	116	64 (4b)	90
3 ^[d,e]	H (1a)	4-MeOC ₆ H ₄ (2c)	119	74 (4c)	91
4 ^[e]	H (1a)	4-ClC ₆ H ₄ (2d)	115	73 (4d)	91
5	H (1a)	4-CF ₃ C ₆ H ₄ (2e)	72	84 (4e)	91
6 ^[e]	H (1a)	2-thienyl (2f)	92	71 (4f)	90
7 ^[e,f]	H (1a)	Ph(CH ₂) ₂ (2g)	95	57 (4g)	88
8 ^[g]	H (1a)	CH ₃ (CH ₂) ₂ (2h)	234	70 (4h)	90
9 ^[g]	H (1a)	CH ₃ (CH ₂) ₄ (2i)	234	77 (4i)	90
10 ^[g]	H (1a)	CH ₃ (CH ₂) ₈ (2j)	235	62 (4j)	91
11 ^[e]	5-Cl (1b)	Ph (2a)	119	63 (4k)	90
12 ^[e]	5-Br (1c)	Ph (2a)	119	72 (4l)	90
13 ^[e]	7-Me (1d)	Ph (2a)	83	70 (4m)	94
14 ^[h]	H (1a)	Ph (2a)	95	> 99 (4a)	91


[a] Reactions were carried out with **1** (0.2 mmol) and of **2** (0.4 mmol; 2 equiv) in benzene (0.5 mL)/1,2-dichloroethane (0.5 mL). [b] Yield of isolated product. [c] Enantiomeric excess was determined by HPLC analysis. [d] The reaction was performed at –20 °C. [e] Employed 5 equiv of **2**. [f] Used 2 equiv of **1a** and 20 mol % of (*R*)-**3** at –20 °C. [g] Used 20 mol % of (*R*)-**3**. [h] The reaction was performed on a one-gram scale (8.56 mmol) in the presence of 850 mg of 3 Å molecular sieves and 5 equiv of **2a**.

enantioselectivities (Table 2, entries 7–10), but long reaction times were necessary to obtain good yields. The reaction tolerated a variety of different indoles (**1b–d**) and gave excellent results (Table 2, entries 11–13).

This reaction was carried out on a one-gram scale to demonstrate the synthetic utility of the present system. When 1.00 g of indole **1a** was treated with 5 equivalents of nitroalkene **2a**, 2.28 g of corresponding adduct **4a** was obtained in high chemical yield without any loss in the enantioselectivity (Table 2, entry 14).

Next, Friedel–Crafts adduct **4a** was transformed into triptamine **5**,^[2a] melatonin analogue **6**,^[2b] and 1,2,3,4-tetrahydro- β -carboline derivative **7**^[2c] (Scheme 2). These products were obtained in good yields without racemization.

To gain insight into the reaction mechanism we examined *N*-Me indole as a substrate under the optimized conditions. As expected, both the chemical yield and the enantioselectivity deteriorated significantly (11 % yield, 0 % ee). The presence of the N–H moiety of indole ring is essential for attaining high yield and enantioselectivity, therefore we assume that the phosphoric acid activates the nitro moiety and at the same time the phosphoryl oxygen atom forms a hydrogen bond with the hydrogen atom of the indole N–H moiety (Figure 1). This arrangement is in agreement with the



Scheme 2. Derivatization of the Friedel–Crafts adduct. Conditions:

a) NaBH₄ (5 equiv), NiCl₂·6 H₂O (1 equiv), MeOH, 30 min, 94 %; b) AcCl (1.5 equiv), Et₃N (1.5 equiv), CH₂Cl₂, 16 h, 91 %; c) PhCHO (1.2 equiv), CF₃CO₂H (2 equiv), MgSO₄, CH₂Cl₂, 68 h, 80 %, *anti*:-*syn* = 91:9; d) TsCl (1.5 equiv), Et₃N (1.5 equiv), CH₂Cl₂, 17 h, 97 %. Ts = *p*-toluenesulfonyl.

previously proposed nine-membered transition state,^[6a,b,g] wherein the phosphoric acid worked as a bifunctional catalyst.

In summary, we have developed a chiral phosphoric acid catalyzed Friedel–Crafts alkylation of indoles with nitroalkenes to generate Friedel–Crafts adducts with excellent enantioselectivities. We found that 3 Å molecular sieves lead to an efficient Friedel–Crafts alkylation in the presence of a chiral phosphoric acid. This reaction is the first example of nitroalkene activation catalyzed by a chiral phosphoric acid, and additional investigations to clarify the reaction mechanism and its application to other enantioselective reactions are underway.

Received: February 16, 2008

Published online: April 16, 2008

Keywords: asymmetric synthesis · Brønsted acids · enantioselectivity · Friedel–Crafts alkylation · indoles




Figure 1. Plausible transition state. The vinylic hydrogen of the nitroalkene lies behind the indole ring.

- [1] For reviews on Friedel–Crafts alkylation reactions, see: a) K. A. Jørgensen, *Synthesis* **2003**, 1117–1125; b) M. Bandini, A. Melloni, A. Umani-Ronchi, *Angew. Chem.* **2004**, *116*, 560–566; *Angew. Chem. Int. Ed.* **2004**, *43*, 550–556; c) M. Bandini, A. Melloni, S. Tommasi, A. Umani-Ronchi, *Synlett* **2005**, 1199–1222; d) S. B. Tsogoeva, *Eur. J. Org. Chem.* **2007**, 1701–1716.
- [2] a) K. S. Jandu, V. Barrett, M. Brockwell, D. Cambridge, D. R. Farrant, C. Foster, H. Giles, R. C. Glen, A. P. Hill, H. Hobbs, A. Honey, G. R. Martin, J. Salmon, D. Smith, P. Woollard, D. L. Selwood, *J. Med. Chem.* **2001**, *44*, 681–693; b) A. Finaru, A. Berthault, T. Besson, G. Guillaumet, S. Berteina-Raboin, *Tetrahedron Lett.* **2002**, *43*, 787–790; c) K. Paulvannan, R. Hale, R. Mesis, T. Chen, *Tetrahedron Lett.* **2002**, *43*, 203–207.
- [3] a) M. Bandini, A. Garelli, M. Rovinetti, M. Tommasi, A. Umani-Ronchi, *Chirality* **2005**, *17*, 522–529; b) W. Zhuang, R. G. Hazell, K. A. Jørgensen, *Org. Biomol. Chem.* **2005**, *3*, 2566–2571; c) R. P. Herrera, V. Sgarzani, L. Bernardi, A. Ricci, *Angew.*

- Chem.* **2005**, *117*, 6734–6737; *Angew. Chem. Int. Ed.* **2005**, *44*, 6576–6579; d) Y.-X. Jia, S.-F. Zhu, Y. Yang, Q.-L. Zhou, *J. Org. Chem.* **2006**, *71*, 75–80; e) S.-F. Lu, D.-M. Du, J. Xu, *Org. Lett.* **2006**, *8*, 2115–2118; f) E. M. Fleming, T. McCabe, S. J. Connon, *Tetrahedron Lett.* **2006**, *47*, 7037–7042; g) P. K. Singh, A. Bisai, V. K. Singh, *Tetrahedron Lett.* **2007**, *48*, 1127–1129.
- [4] For reviews on Brønsted acid catalysis, see: a) P. R. Schreiner, *Chem. Soc. Rev.* **2003**, *32*, 289–296; b) P. M. Pihko, *Angew. Chem.* **2004**, *116*, 2110–2113; *Angew. Chem. Int. Ed.* **2004**, *43*, 2062–2064; c) C. Bolm, T. Rantanen, I. Schiffrers, L. Zani, *Angew. Chem.* **2005**, *117*, 1788–1793; *Angew. Chem. Int. Ed.* **2005**, *44*, 1758–1763; d) P. M. Pihko, *Lett. Org. Chem.* **2005**, *2*, 398–403; e) Y. Takemoto, *Org. Biomol. Chem.* **2005**, *3*, 4299–4306; f) M. S. Taylor, E. N. Jacobsen, *Angew. Chem.* **2006**, *118*, 1550–1573; *Angew. Chem. Int. Ed.* **2006**, *45*, 1520–1543; g) T. Akiyama, J. Itoh, K. Fuchibe, *Adv. Synth. Catal.* **2006**, *348*, 999–1010; h) X. Yu, W. Wang, *Chem. Asian J.* **2008**, *3*, 516–532.
- [5] For reviews on phosphoric acid catalysis, see: a) S. J. Connon, *Angew. Chem.* **2006**, *118*, 4013–4016; *Angew. Chem. Int. Ed.* **2006**, *45*, 3909–3912; b) T. Akiyama, *Chem. Rev.* **2007**, *107*, 5744–5758.
- [6] Our reports, see: a) T. Akiyama, J. Itoh, K. Yokota, K. Fuchibe, *Angew. Chem.* **2004**, *116*, 1592–1594; *Angew. Chem. Int. Ed.* **2004**, *43*, 1566–1568; b) T. Akiyama, H. Morita, J. Itoh, K. Fuchibe, *Org. Lett.* **2005**, *7*, 2583–2585; c) T. Akiyama, Y. Saitoh, H. Morita, K. Fuchibe, *Adv. Synth. Catal.* **2005**, *347*, 1523–1526; d) T. Akiyama, Y. Tamura, J. Itoh, H. Morita, K. Fuchibe, *Synlett* **2006**, 141–143; e) J. Itoh, T. Akiyama, K. Fuchibe, *Angew. Chem.* **2006**, *118*, 4914–4916; *Angew. Chem. Int. Ed.* **2006**, *45*, 4796–4798; f) T. Akiyama, H. Morita, K. Fuchibe, *J. Am. Chem. Soc.* **2006**, *128*, 13070–13071; g) M. Yamanaka, J. Itoh, K. Fuchibe, T. Akiyama, *J. Am. Chem. Soc.* **2007**, *129*, 6756–6764; h) T. Akiyama, Y. Honma, J. Itoh, K. Fuchibe, *Adv. Synth. Catal.* **2008**, *350*, 399–402.
- [7] For selected examples of the phosphoric acid catalyzed reactions, see: a) D. Uraguchi, M. Terada, *J. Am. Chem. Soc.* **2004**, *126*, 5356–5357; b) G. B. Rowland, H. Zhang, E. B. Rowland, S. Chennamadhavuni, Y. Wang, J. C. Antilla, *J. Am. Chem. Soc.* **2005**, *127*, 15696–15697; c) R. I. Storer, D. E. Carrera, Y. Ni, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2006**, *128*, 84–86; d) J. Seayad, A. M. Seayad, B. List, *J. Am. Chem. Soc.* **2006**, *128*, 1086–1087; e) M. Rueping, E. Sugiono, C. Azap, *Angew. Chem.* **2006**, *118*, 2679–2681; *Angew. Chem. Int. Ed.* **2006**, *45*, 2617–2619; f) X.-H. Chen, X.-Y. Xu, H. Liu, L.-F. Cun, L.-Z. Gong, *J. Am. Chem. Soc.* **2006**, *128*, 14802–14803; g) M. Rueping, C. Azap, *Angew. Chem.* **2006**, *118*, 7996–7999; *Angew. Chem. Int. Ed.* **2006**, *45*, 7832–7835; h) G. Li, Y. Liang, J. C. Antilla, *J. Am. Chem. Soc.* **2007**, *129*, 5830–5831; i) J. Zhou, B. List, *J. Am. Chem. Soc.* **2007**, *129*, 7498–7499; j) M. Terada, K. Machioka, K. Sorimachi, *J. Am. Chem. Soc.* **2007**, *129*, 10336–10337; k) M. Rueping, A. P. Antonchick, C. Brinkmann, *Angew. Chem.* **2007**, *119*, 7027–7030; *Angew. Chem. Int. Ed.* **2007**, *46*, 6903–6906.
- [8] a) D. Nakashima, H. Yamamoto, *J. Am. Chem. Soc.* **2006**, *128*, 9626–9627; b) M. Rueping, W. Ieawsuwan, A. P. Antonchick, B. J. Nachtsheim, *Angew. Chem.* **2007**, *119*, 2143–2146; *Angew. Chem. Int. Ed.* **2007**, *46*, 2097–2100; c) E. B. Rowland, G. B. Rowland, E. Rivera-Otero, J. C. Antilla, *J. Am. Chem. Soc.* **2007**, *129*, 12084–12085; d) P. Jiao, D. Nakashima, H. Yamamoto, *Angew. Chem.* **2008**, *120*, 2445–2447; *Angew. Chem. Int. Ed.* **2008**, *47*, 2411–2413.
- [9] Phosphoric acid catalyzed enantioselective addition reactions of furan, indole, and pyrrole to imines and α , β -unsaturated ketones have been reported: a) D. Uraguchi, K. Sorimachi, M. Terada, *J. Am. Chem. Soc.* **2004**, *126*, 11804–11805; b) M. Terada, K. Sorimachi, *J. Am. Chem. Soc.* **2007**, *129*, 292–293; c) M. Terada, S. Yokoyama, K. Sorimachi, D. Uraguchi, *Adv. Synth. Catal.* **2007**, *349*, 1863–1867; d) Q. Kang, Z.-A. Zhao, S.-L. You, *J. Am. Chem. Soc.* **2007**, *129*, 1484–1485; e) G. B. Rowland, E. B. Rowland, Y. Liang, J. A. Perman, J. C. Antilla, *Org. Lett.* **2007**, *9*, 2609–2611; f) G. Li, G. B. Rowland, E. B. Rowland, J. C. Antilla, *Org. Lett.* **2007**, *9*, 4065–4068; g) Y.-X. Jia, J. Zhong, S.-F. Zhu, C.-M. Zhang, Q.-L. Zhou, *Angew. Chem.* **2007**, *119*, 5661–5663; *Angew. Chem. Int. Ed.* **2007**, *46*, 5565–5567; h) M. Rueping, B. J. Nachtsheim, S. A. Moreth, M. Bolte, *Angew. Chem.* **2008**, *120*, 603–606; *Angew. Chem. Int. Ed.* **2008**, *47*, 593–596; i) M. J. Wanner, P. Hauwert, H. E. Schoemaker, R. D. Gelder, J. H. van Maarseveen, H. Hiemstra, *Eur. J. Org. Chem.* **2008**, 180–185; j) H.-Y. Tang, A.-D. Lu, Z.-H. Zhou, G.-F. Zhao, L.-N. He, C.-C. Tang, *Eur. J. Org. Chem.* **2008**, 1406–1410; See also: k) Y.-Q. Wang, J. Song, R. Hong, H. Li, L. Deng, *J. Am. Chem. Soc.* **2006**, *128*, 8156–8157.
- [10] The absolute stereochemistry of **4a** was determined by chiral HPLC analysis by comparison to the retention time described in the reference [3c].
- [11] The powdered 3 Å molecular sieves were activated by placing the powder under vacuum and heating with a heat gun for 10 minutes.
- [12] For leading references on molecular sieves, see: a) J. K. Thomas, *Chem. Rev.* **1993**, *93*, 301–320; b) M. E. Davis, *Acc. Chem. Res.* **1993**, *26*, 111–115.
- [13] When **1a** and **2a** were treated with 3 Å M.S. alone, **4a** was obtained in 3% yield.