


# Self-Assembly of a Cyclic $\text{Zn}_4\text{O}_4$ Tetramer by Aerobic Oxidation of a Bisoxazoline: A Molecular “Nest” for Nucleophilic $\text{OH}^-$ \*\*

Béatrice Jacques, Clémence Dro, Stéphane Bellemin-Laponnaz,\* Hubert Wadepohl, and Lutz H. Gade\*

The bewildering complexity of biological structures and functions is based on a limited number of building blocks. Further transformation of these primary structural elements provides the key to the assembly of functional units of great complexity. The oxidative functionalization of biological monomers, in particular amino acids, is an essential step in the generation of the binding sites for metal atoms in the form of catecholates, hydroxamates, and related ligating functions.<sup>[1]</sup> Their complexation to substitutionally labile metal ions gives rise to stable highly ordered mono or polynuclear coordination units.<sup>[2]</sup> These may play a role in the sequestering of metal ions, the stabilization of biopolymer structures or as active sites in metalloenzymes.<sup>[3]</sup>

There is an ever growing number of synthetic host molecules for the binding of charged or neutral molecules which may be employed as catalysts, building blocks for extended solids, and sensors.<sup>[4]</sup> Many of these are metallamacrocycles<sup>[5]</sup> or have cage structures based on pre-conceived ligand architectures which




**Scheme 1.** Synthesis of the tetranuclear complex  $[\{(\text{Me}_2\text{box})\text{O}\}_4\text{Zn}_4(\text{OH})][\text{ClO}_4]_3$  **2** and its conversion into  $[\{(\text{Me}_2\text{box})\text{O}\}_4\text{Zn}_4][\text{ClO}_4]_4$  (**4**).

assemble to give the polynuclear aggregate.<sup>[6]</sup> Herein we report the combination of a selective functionalization step, the aerobic oxidation of a bisoxazoline, with the formation of a highly ordered metallamacrocyclic complex which itself is shown to be a functional catalytic system.

Stirring equimolar amounts of the bisoxazoline  $\text{Me}_2\text{box}$ <sup>[7]</sup> (**1**) and  $\text{Zn}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$  in methanol for 24 h led to the precipitation of a white crystalline solid, which was characterized as the tetranuclear complex  $[\{(\text{Me}_2\text{box})\text{O}\}_4\text{Zn}_4(\text{OH})][\text{ClO}_4]_3$  (**2**; Scheme 1).

The mass balance in Equation (1) implies the generation



of perchloric acid during the course of the oxygenation, which in turn explains the formation of protonated bisoxazoline as a by-product and the incomplete conversion (isolated yield of

[\*] Dr. B. Jacques, Prof. Dr. H. Wadepohl, Prof. Dr. L. H. Gade  
Anorganisch-Chemisches Institut  
Universität Heidelberg  
Im Neuenheimer Feld 270, 69120 Heidelberg (Germany)  
Fax: (+49) 6221-545-609  
E-mail: lutz.gade@uni-hd.de


Dr. C. Dro, Dr. S. Bellemin-Laponnaz  
Institut de Chimie  
Université Louis Pasteur Strasbourg  
4 rue Blaise Pascal, 67000 Strasbourg (France)  
E-mail: bellemin@chimie.u-strasbg.fr

[\*\*] We thank the Deutsche Forschungsgemeinschaft (SFB 623) and the CNRS for funding, the Alexander von Humboldt-Stiftung for a research fellowship (B.J.) and the MENRT (France) for a Ph.D. grant (C.D.). We are also grateful to Prof. R. Welter and Dr. A. De Cian for one of the X-ray diffraction analyses.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.


$2 \leq 40\%$ ). Carrying out the reaction in the presence of  $\text{NaHCO}_3$  gave the tetranuclear complex  $[(\text{Me}_2\text{box})\text{O}]_4\text{Zn}_4(\text{OH})[\text{ClO}_4]_3$  (**2**) in 88 % yield with respect to  $\text{Zn}(\text{ClO}_4)_2$ .

A single-crystal X-ray structure analysis of **2**<sup>[8]</sup> established the eight-membered ring core consisting of four zinc and four oxygen atoms (Figure 1).<sup>[9]</sup> Each zinc atom is coordinated by two nitrogen atoms from two different oxazoline ligands as well as two alkoxy oxygen atoms in a  $\mu_2$ -O mode. The hydroxide ion bridges three of the four zinc atoms [Zn(1)-O(15) 2.144(1), Zn(3)-O(15) 2.170(1), and Zn(4)-O(15) 2.113(1) Å].<sup>[10]</sup>




**Figure 1.** Side view of the molecular structure of the  $\{\text{Zn}_4\text{-OH}\}$  complex **2**. Two of the three perchlorate ions are shown. All hydrogen atoms (except for the OH hydrogen atom) are omitted for clarity. Principal bond lengths [Å] and angles [°]: O(15)-Zn(4) 2.113(1), O(15)-Zn(1) 2.144(1), O(15)-Zn(3) 2.170(1), O(15)-Zn(2) 3.020(1), O(15)-H(15) 0.77(3); O(14)-Zn(4)-O(13) 163.57(5), O(13)-Zn(3)-O(12) 171.77(5), O(11)-Zn(2)-O(12) 139.76(5), O(14)-Zn(1)-O(11) 173.40(5), Zn(2)-O(11)-Zn(1) 113.82(5), Zn(2)-O(12)-Zn(3) 118.67(6), Zn(3)-O(13)-Zn(4) 96.19(5), Zn(4)-O(14)-Zn(1) 99.70(5).

The formation of the tetranuclear complex **2**, in which the bisoxazolines are oxygenated to give alkoxy ligands,<sup>[11]</sup> occurred upon conducting this transformation in the presence of air. Under a rigorous anaerobic atmosphere of purified argon, the reaction did not give a similar oxygenation of the ligand and self-assembly of tetrameric aggregates, but afforded the mononuclear complex  $[(\text{Me}_2\text{box})\text{Zn}(\text{H}_2\text{O})_3](\text{ClO}_4)_2$  (**3**), the triflate salt **3a** of which has been structurally characterized.<sup>[8,12]</sup> Performing the reaction under an atmosphere of  $^{18}\text{O}_2$  provided further proof for a clean and selective aerobic oxidation of the ligand [Eq. (2)].



The formation of the  $^{18}\text{O}$ -labeled complex **2a** was confirmed by the ESI mass spectrum, which indicates, at  $m/z$  1435.1918, the presence of a tetranuclear complex cation


corresponding to the formula  $[(\text{Me}_2\text{box})^{18}\text{O}]_4\text{Zn}_4(\text{OH})[\text{ClO}_4]_2^+$  (Figure 2). IR spectroscopic data are also consistent with the introduction of an  $^{18}\text{O}$  atom on the bridging carbon of the bisoxazoline: the C-O stretching mode observed at  $1151 \text{ cm}^{-1}$  in complex **2** is shifted to a lower frequency at  $1144 \text{ cm}^{-1}$  in  $^{18}\text{O}$ -labeled **2a** (see the Supporting Information).



**Figure 2.** ESI mass spectra of: a) the complex  $[(\text{Me}_2\text{box})\text{O}]_4\text{Zn}_4(\text{OH})[\text{ClO}_4]_3$  **2** and b) the complex  $[(\text{Me}_2\text{box})^{18}\text{O}]_4\text{Zn}_4(\text{OH})[\text{ClO}_4]_3$  **2a**.

Whereas the slow aerobic autoxidation of uncoordinated bisbenzoxazoles has previously been noted,<sup>[13]</sup> in the case at hand, no oxidation of the  $\text{Me}_2\text{box}$  ligand occurred in the absence of the metal salt,<sup>[14]</sup> suggesting a significant role of the metal in this selective oxidation of a box derivative.<sup>[15]</sup> Zinc is thought to activate the neutral box, which is deprotonated giving the bisoxazolinato species.<sup>[16]</sup> In turn, this anionic bisoxazolinato will react with oxygen<sup>[17]</sup> (acting as an electrophilic diradical) in a similar way as observed by Itoh and co-workers for  $\beta$ -ketiminates.<sup>[18]</sup> In the presence of radical scavengers, such as TEMPO (2,2,6,6-tetramethylpiperidine-*N*-oxide) and BHT (2,6-di-*tert*-butyl-4-methylphenol), the formation of **2/2a** was not observed.

Upon reaction of  $[(\text{Me}_2\text{box})\text{O}]_4\text{Zn}_4(\text{OH})^{3+}$  (**2**) with one equivalent of  $\text{HBF}_4$  in diethyl ether, quantitative conversion into the corresponding  $\text{OH}^-$ -free tetracationic complex  $[(\text{Me}_2\text{box})\text{O}]_4\text{Zn}_4^{4+}$  (**4**) took place immediately (Scheme 1). This complex was obtained in various crystalline forms, invariably cocrystallizing with solvent (dichloromethane and/or methanol, hexane). Its molecular structure, determined by X-ray diffraction of a hexane-containing crystal, is depicted in Figure 3a.<sup>[8]</sup> The molecular cation has crystallographic  $C_2$  symmetry, and the alternating arrangement of the bisoxazoline ligands confers an overall virtual  $D_{2d}$



**Figure 3.** Molecular structure of the complex tetracation in **4**: a) view along the virtual fourfold improper axis, and b) the adduct of the cation with two  $\text{ClO}_4^-$  counterions occupying the molecular “nest” structure. Space-filling models of: c) the complex cation and d) the  $\text{ClO}_4^-$  adduct, showing the excellent fit between the  $\{\text{Zn}_4\text{O}_4\}$  host and the guest anions. Zn light blue, N dark blue, O red, C gray, Cl green. Principal bond lengths [Å] and angles [°]: Zn–N 2.038(3)–2.043(3), Zn–O(11,12) 1.982(2)–2.009(2), Cl(1)–O(21)/O(22) 1.445(3)/1.442(3), Cl(1)–O(23)/O(24) 1.429(3)/1.423(3); N–Zn–N 111.9(2)–114.8(2), O–Zn–O 141.7(1)–149.1(1), Zn(1)–O(11)–Zn(2) 124.1(1), Zn(3)–O(12)–Zn(2) 121.1(1).

symmetry to the cationic tetramer, which may be viewed as a molecular “nest” for anionic or Lewis basic guest molecules (Figure 3c). Four perchlorate anions balance the charge of the complex cation. In the crystal, two of them are arranged as molecular guests above and below the  $\{\text{Zn}_4\text{O}_4\}$  core of the molecule, with the eight-membered ring being perfectly suited to accommodate the bridging O–Cl–O moiety of the anions (Figure 3b and d). The  $\text{Zn}\cdots\text{O}_{\text{perchlorate}}$  distances are 2.54–2.63 Å, and this bonding interaction is also reflected by the elongation of the Cl–O bond lengths within the perchlo-

rate ions ( $\text{Cl–O}_{\text{terminal}} = 1.42\text{--}1.43$  Å vs.  $1.45\text{--}1.46$  Å).<sup>[19]</sup> The exact fit of anions and cations is apparent in the space-filling model depicted in Figure 3d and consistent with the observation of the mass peaks in the ESI MS spectra corresponding to  $\{\text{cation} + 2\text{ClO}_4\}^{+2+}$ .

The hydroxo complex **2** is formed from **4** under neutral conditions in aqueous solution. This suggests that the  $\text{p}K_{\text{a}}$  of zinc(II)-bound water is close to physiological pH. A potentiometric titration experiment gave a  $\text{p}K_{\text{a}}$  value of 8.2. Given this “activation of water”, we probed the ability of **2** to

catalyze the hydrolysis of phosphates. The reactivity of phosphotriester substrate tris(4-nitrophenyl)phosphate (TNP) was evaluated in buffered EtOH/H<sub>2</sub>O solutions.<sup>[20,21]</sup> The reaction was found to be first order in both tetranuclear zinc complex and substrate, and the second-order rate constants of hydrolysis were evaluated from pH 7.0 to 9.2. The pH rate profile revealed an inflection point at 8.15 which corresponds to the pK<sub>a</sub> obtained in the potentiometric titration. We may therefore conclude that the Zn<sub>3</sub>-bridging hydroxide is most likely the base that induces the TNP hydrolysis. The relative reactivity of metal-promoted hydrolysis versus free OH<sup>−</sup> provides some insight in the mechanism of the hydrolysis.<sup>[22]</sup> The second-order rate constant (*k'*<sub>max</sub>)<sup>[23]</sup> of the zinc-OH catalyzed hydrolysis was found to be 2.2 M<sup>−1</sup> s<sup>−1</sup>, and is slightly lower than that observed for the free OH<sup>−</sup> ion (10.7 M<sup>−1</sup> s<sup>−1</sup>).<sup>[23]</sup> The nucleophilicity of a metal-bound hydroxide must be lower than free OH<sup>−</sup> on the basis of simple steric and charge considerations. Therefore, the fact that the rate hydrolysis with the zinc complex is of approximately the same order of magnitude of free OH<sup>−</sup> appears to be consistent with a mechanism involving a nucleophilic attack of the metal-bound hydroxide upon the phosphorus atom aided by a certain degree of zinc–phosphate affinity.

In summary, a simple molecule is activated towards a highly selective aerobic oxygenation by metal complexation. This is relevant for the use of bisoxazolines and bisoxazolinates as stereodirecting ligands in enantioselective catalysis.<sup>[24]</sup> In the case at hand, the oxidation is followed by the assembly in the form of a highly ordered metallamacrocyclic which displays catalytic phosphatase activity. The combination of these different levels of structural assembly and associated function may be viewed as a simple model for the systems of far greater complexity encountered in nature.

Received: February 22, 2008  
Published online: May 5, 2008

**Keywords:** autoxidation · metallacycles · oxazoline · phosphatase · zinc

- [1] *Microbial Transport Systems* (Ed.: G. Winkelmann), Wiley-VCH, Weinheim, 2002.
- [2] a) *Concepts and Models in Bioinorganic Chemistry* (Eds.: H.-B. Kraatz, N. Metzler-Nolte), Wiley, New York, 2006; b) I. Bertini, H. B. Gray, E. I. Stiefel, J. S. Valentine, *Biological Inorganic Chemistry*, University Science Books, Mill Valley, 2007; c) W. Kaim, B. Schwederski, *Bioinorganic Chemistry: Inorganic Elements in the Chemistry of Life*, Wiley, New York, 1994; d) S. J. Lippard, J. M. Berg, *Principles of Bioinorganic Chemistry*, University Science Books, Mill Valley, 1994.
- [3] R. H. Holm, P. Kennepohl, E. I. Solomon, *Chem. Rev.* 1996, 96, 2239.
- [4] a) J. M. Lehn, *Supramolecular Chemistry: Concepts and Perspectives*, VCH, Weinheim, 1995; b) *Transition Metals in Supramolecular Chemistry, Vol. 5* (Ed.: J.-P. Sauvage), Wiley, New York, 1999.
- [5] For a comprehensive overview of the field of metallamacrocycles, see: G. Mezei, C. M. Zaleski, V. L. Peccoraro, *Chem. Rev.* 2007, 107, 4933.
- [6] Selected reviews: a) D. L. Caulder, K. N. Raymond, *Acc. Chem. Res.* 1999, 32, 975; b) M. Albrecht, *Chem. Rev.* 2001, 101, 3457; c) M. Fujita, M. Tominaga, A. Hori, B. Therrien, *Acc. Chem. Res.* 2005, 38, 369; d) D. Fiedler, D. H. Leung, R. G. Bergman, K. N. Raymond, *Acc. Chem. Res.* 2005, 38, 349; e) C. J. Johnson, *Chem. Soc. Rev.* 1998, 27, 289.
- [7] a) S. Bellemin-Laponnaz, L. H. Gade, *Chem. Commun.* 2002, 1286; b) S. Dagorne, S. Bellemin-Laponnaz, R. Welter, *Organometallics* 2004, 23, 3053.
- [8] Crystal data for complex **2**: C<sub>49</sub>H<sub>79</sub>Cl<sub>5</sub>N<sub>8</sub>O<sub>25</sub>Zn<sub>4</sub>, monoclinic, space group *P*2<sub>1</sub>/*c*, *a* = 13.9925(7), *b* = 18.0171(8), *c* = 25.6993(12) Å, β = 91.185(1)°, *V* = 6477.5(5) Å<sup>3</sup>, *Z* = 4, μ = 1.755 mm<sup>−1</sup>, *F*<sub>000</sub> = 3336, *T* = 100(2) K; θ range 1.8 to 32.3°. Index ranges *h,k,l* (indep. set): −20–20, 0–26, 0–38. Reflections coll 139 158, ind 21 752 [*R*<sub>int</sub> = 0.0379], obs [*I* > 2σ(*I*): 17 382. Final *R* indices [*F*<sub>o</sub> > 4σ(*F*<sub>o</sub>): *R*(*F*) = 0.0344, *wR*(*F*<sup>2</sup>) = 0.0822, *GooF* = 1.029. Crystal data for complex **3a**: C<sub>14</sub>H<sub>25</sub>F<sub>6</sub>N<sub>2</sub>O<sub>11</sub>S<sub>2</sub>Zn, orthorhombic, space group *Pbcn*, *a* = 12.1460(3), *b* = 13.0680(3), *c* = 16.5790(4) Å, *V* = 2631.5(1) Å<sup>3</sup>, *Z* = 4, μ = 1.187 mm<sup>−1</sup>, *F*<sub>000</sub> = 1308. *T* = 173(2) K; *q* range 2.3 to 29.1°. Index ranges *h,k,l*: −16–16, −17–17, −22–22. Reflections coll 6527, ind 3543 [*R*<sub>int</sub> = 0.0315], obs [*I* > 2σ(*I*): 2281. Final *R* indices [*F*<sub>o</sub> > 4σ(*F*<sub>o</sub>): *R*(*F*) = 0.0464, *wR*(*F*<sup>2</sup>) = 0.1266, *GooF* = 0.966. Crystal data for complex **4**: C<sub>48</sub>H<sub>76</sub>Cl<sub>4</sub>N<sub>8</sub>O<sub>28</sub>Zn<sub>4</sub>·*n* hexane: orthorhombic, space group *Pbcn*, *a* = 21.341(5), *b* = 15.190(3), *c* = 23.750(5) Å, *V* = 7699(3) Å<sup>3</sup>, *Z* = 4, μ = 1.445 mm<sup>−1</sup>, *F*<sub>000</sub> = 3328. *T* = 100(2) K; θ range 1.9 to 29.6°. Index ranges *h,k,l* (ind set): −29–0, 0–21, −32–0. Reflections coll 175 724, ind 10 805 [*R*<sub>int</sub> = 0.0831], obs [*I* > 2σ(*I*): 6979. Final *R* indices [*F*<sub>o</sub> > 4σ(*F*<sub>o</sub>): *R*(*F*) = 0.0656, *wR*(*F*<sup>2</sup>) = 0.2044, *GooF* = 1.130. Data collection at low temperature (Bruker AXS Smart 1000 CCD (**2**, **4**) or Enraf Nonius Kappa CCD (**3a**) diffractometer, Mo Kα radiation, graphite monochromator, λ = 0.71073 Å), semiempirical absorption correction,<sup>[8a]</sup> Structure solution: heavy atom method combined with structure expansion by direct methods (**2**),<sup>[8b]</sup> direct methods (**3a**),<sup>[8c]</sup> or direct methods with dual-space recycling (**4**).<sup>[8d]</sup> Refinement: full-matrix least squares on *F*<sup>2</sup>; all non-hydrogen atoms anisotropic, hydrogen atoms at calculated positions (refined riding).<sup>[8e]</sup> Electron density attributed to heavily disordered *n*-hexane was removed from the structure (and the corresponding *F*<sub>o</sub>) of **4** with the SQUEEZE procedure.<sup>[8f,g]</sup> CCDC-675278 (**2**), CCDC-675279 (**3a**), and CCDC-675280 (**4**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data\_request/cif; a) G. M. Sheldrick, *SADABS*, Bruker AXS, 2004–2007; b) P. T. Beurskens in *Crystallographic Computing 3* (Eds.: G. M. Sheldrick, C. Krüger, R. Goddard), Clarendon Press, Oxford, UK, 1985, p. 216; P. T. Beurskens, G. Beurskens, R. de Gelder, J. M. M. Smits, S. Garcia-Granda, R. O. Gould, *DIRDIF-2007*, Raboud University Nijmegen, The Netherlands, 2007; c) G. M. Sheldrick, *SHELXS-86*, University of Göttingen, 1986; G. M. Sheldrick, *Acta Crystallogr. Sect. A* 1990, 46, 467; d) G. M. Sheldrick, H. A. Hauptman, C. M. Weeks, R. Miller, I. Usón, *Ab initio phasing in International Tables for Crystallography, Vol. F* (Eds.: M. G. Rossmann, E. Arnold), IUCr and Kluwer Academic Publishers, Dordrecht, 2001, pp. 333–351; G. M. Sheldrick, *SHELXD*, University of Göttingen 2000–4; e) G. M. Sheldrick, *SHELXL-97*, University of Göttingen, 1997; G. M. Sheldrick, *Acta Crystallogr. Sect. A* 2008, 64, 112; f) P. van der Sluis, A. L. Spek, *Acta Crystallogr. Sect. A* 1990, 46, 194; g) A. L. Spek, *PLATON*, Utrecht University, The Netherlands; A. L. Spek, *J. Appl. Crystallogr.* 2003, 36, 7.
- [9] Structurally characterized organozinc {Zn<sub>4</sub>O<sub>4</sub>} tetramers: a) F. H. van der Steen, J. Boersma, A. L. Spek, G. van Koten, *Organometallics* 1991, 10, 2467; b) P. A. van der Schaaf, E. Wissing, J. Boersma, W. J. J. Smeets, A. L. Spek, G. van Koten, *Organometallics* 1993, 12, 3624; c) J. Hunger, S. Blaurock, J. Sieler, *Z. Anorg. Allg. Chem.* 2005, 631, 472; d) E. Asato, H.

- Furutachi, T. Kawahashi, M. Mikuriya, *J. Chem. Soc. Dalton Trans.* **1995**, 3897; other {Zn<sub>4</sub>} assemblies: e) B. Müller, A. Schneider, M. Tesmer, H. Vahrenkamp, *Inorg. Chem.* **1999**, 38, 1900; f) C. J. Matthews, L. K. Thompson, S. R. Parsons, Z. Xu, D. O. Miller, S. L. Heath, *Inorg. Chem.* **2001**, 40, 4448; g) A. Demšar, J. Košmrlj, S. Petriček, *J. Am. Chem. Soc.* **2002**, 124, 3951.
- [10] For macrocyclic {Zn<sub>4</sub>} complexes stabilizing {Zn<sub>4</sub>(μ-OH)} units, see: a) M. J. Grannas, B. F. Hoskins, R. Robson, *Inorg. Chem.* **1994**, 33, 1071; b) M. Bell, A. J. Edwards, B. F. Hoskins, E. H. Kachab, R. Robson, *J. Am. Chem. Soc.* **1989**, 111, 3603.
- [11] Lippard et al. reported a N<sub>2</sub>O<sup>-</sup> ligand that may be seen as the bis(imidazole)analogue. Investigation of coordination chemistry with zinc revealed a different behavior; see: a) F. Chu, J. Smith, V. M. Lynch, E. V. Anslyn, *Inorg. Chem.* **1995**, 34, 5689; b) W. B. Tolman, S. Liu, J. G. Bentsen, S. J. Lippard, *J. Am. Chem. Soc.* **1991**, 113, 152. For selected examples of mononuclear zinc model complexes with tripodal N,N,O ligands, see: c) F. Gross, H. Vahrenkamp, *Inorg. Chem.* **2005**, 44, 3321; d) P. Ghosh, G. Parkin, *J. Chem. Soc. Dalton Trans.* **1998**, 2281; e) B. S. Hammes, C. J. Carrano, *Inorg. Chem.* **1999**, 38, 4593; f) I. Hegelmann, A. Beck, C. Eichhorn, B. Weibert, N. Burzlaff, *Eur. J. Inorg. Chem.* **2003**, 339.
- [12] Structurally characterized bisoxazoline zinc complexes: a) S. Crosignani, G. Desimoni, G. Faita, S. Filippone, A. Mortoni, P. Rigetti, M. Zema, *Tetrahedron Lett.* **1999**, 40, 7007; b) J. Thorhauge, M. Roberson, R. G. Hazell, K. A. Jørgensen, *Chem. Eur. J.* **2002**, 8, 1888.
- [13] L. Forlani, C. Boga, E. Del Vecchio, M. Padovani, *Arkivoc* **2003**, xv, 75.
- [14] On the other hand, mixing bisoxazoline **1** and Zn(ClO<sub>4</sub>)<sub>2</sub>·6H<sub>2</sub>O in methanol led, after a few minutes, to the formation of a white precipitate. Isolated straight away, it was identified as the mononuclear complex [(Me<sub>2</sub>box)Zn(H<sub>2</sub>O)<sub>3</sub>](ClO<sub>4</sub>)<sub>2</sub> (**3**). However, when this reaction mixture was stirred in air over longer periods of time, it slowly converted into the tetrameric complex [(Me<sub>2</sub>box)O]<sub>4</sub>Zn<sub>4</sub>(OH)[ClO<sub>4</sub>]<sub>3</sub> (**2**).
- [15] V. Čaplar, Z. Roje, V. Tomišić, G. Horvat, J. Požar, I. Piantanida, M. Žinić, *Tetrahedron* **2004**, 60, 8079.
- [16] S. Dagorne, S. Bellemin-Laponnaz, A. Maise-François, *Eur. J. Inorg. Chem.* **2007**, 913.
- [17] For the reaction of zinc alkyls with oxygen, see: a) J. Lewiński, W. Marciniak, J. Lipkowski, I. Justyniak, *J. Am. Chem. Soc.* **2003**, 125, 12698; b) J. Lewiński, Z. Ochal, E. Bojarski, E. Tratkiewicz, I. Justyniak, J. Lipkowski, *Angew. Chem.* **2003**, 115, 4791; *Angew. Chem. Int. Ed.* **2003**, 42, 4643; c) J. Lewiński, W. Śliwiński, M. Dranka, I. Justyniak, J. Lipkowski, *Angew. Chem.* **2006**, 118, 4944; *Angew. Chem. Int. Ed.* **2006**, 45, 4826; d) J. Lewiński, W. Bury, M. Dutkiewicz, M. Maurin, I. Justyniak, J. Lipkowski, *Angew. Chem.* **2008**, 120, 583; *Angew. Chem. Int. Ed.* **2008**, 47, 573.
- [18] For the oxidative degradation of β-diketiminato ligands, see: S. Yokota, Y. Tachi, S. Itoh, *Inorg. Chem.* **2002**, 41, 1342.
- [19] Perchlorate structures: a) P. K. Coughlin, S. J. Lippard, *J. Am. Chem. Soc.* **1981**, 103, 3228; b) P. K. Coughlin, S. J. Lippard, *J. Am. Chem. Soc.* **1984**, 106, 2328; c) Effendy, C. di Nicola, M. Nitiatmodjo, C. Pettinari, B. W. Skelton, A. H. White, *Inorg. Chim. Acta* **2005**, 358, 735; d) K. B. Jensen, E. Johansen, F. B. Larsen, C. J. McKenzie, *Inorg. Chem.* **2004**, 43, 3801; e) S. K. Mandal, L. K. Thompson, M. J. Newlands, E. J. Gabe, *Inorg. Chem.* **1989**, 28, 3707; f) H. Adams, S. Clunas, D. E. Fenton, *Chem. Commun.* **2002**, 418; g) O. Mamula, A. von Zelewsky, T. Bark, H. Stoeckli-Evans, A. Neels, G. Bernadinelli, *Chem. Eur. J.* **2000**, 6, 3575.
- [20] The hydrolysis of TNP was monitored by UV analysis (405 nm); see the Supporting Information for experimental details. See also: D. P. Goldberg, R. C. di Targiani, F. Namuswe, E. C. Minnihan, S. Chang, L. N. Zakharov, A. L. Rheingold, *Inorg. Chem.* **2005**, 44, 7559.
- [21] Selected reviews covering the field of artificial phosphatases/nucleases: a) F. Mancin, P. Scrimin, P. Tecilla, U. Tonellato, *Chem. Commun.* **2005**, 2540; b) R. Ott, R. Krämer, *Appl. Microbiol. Biotechnol.* **1999**, 52, 761; c) E. L. Hegg, J. N. Burstyn, *Coord. Chem. Rev.* **1998**, 173, 133; d) L. M. Berreau, *Adv. Phys. Org. Chem.* **2006**, 41, 79; e) F. Meyer, *Eur. J. Inorg. Chem.* **2006**, 3789; f) D. E. Wilcox, *Chem. Rev.* **1996**, 96, 2435.
- [22] a) S. H. Gellman, R. Petter, R. Breslow, *J. Am. Chem. Soc.* **1986**, 108, 2388; b) T. Koike, E. Kimura, *J. Am. Chem. Soc.* **1991**, 113, 8935.
- [23] *k*<sub>max</sub> represents the pH-independent rate constant. See the Supporting Information for details.
- [24] Reviews: a) F. Fache, E. Schulz, M. Lorraine Tommasino, M. Lemaire, *Chem. Rev.* **2000**, 100, 2159; b) H. A. McManus, P. J. Guiry, *Chem. Rev.* **2004**, 104, 4151.