A Journal of the Gesellschaft Deutscher Chemiker IS Chemiker Chemiker Chemiker Chemiker Chemiker Chemiker Www.angewandte.org International Edition 2008-47/31 K Rb K^{+} Na Nat Rb K Nat Rb

A synthetic ion channel ...

... based on a metal–organic polyhedron (MOP) is described by K. Kim and co-workers in the Communication on page 5755 ff. The MOP transports protons and alkali metal ions across lipid membranes with cation selectivity in the order of $Li^+ \gg Na^+ > K^+ > Rb^+ > Cs^+$. Cation– π interactions between cations and the aromatic rings lining the windows of the MOP seem to play a crucial role in ion transport.



Inside Cover

Minseon Jung, Hyunuk Kim, Kangkyun Baek, and Kimoon Kim*

A synthetic ion channel based on a metal–organic polyhedron (MOP) is described by K. Kim and co-workers in the Communication on page 5755 ff. The MOP transports protons and alkali metal ions across lipid membranes with cation selectivity in the order of $Li^+ \! \! > \! \! Na^+ \! \! > \! \! K^+ \! \! > \! Rb^+ \! \! > \! \! \! Cs^+\! \! .$ Cation– π interactions between cations and the aromatic rings lining the windows of the MOP seem to play a crucial role in ion transport.

