


Conclusive Evidence for an S_N2 -Si Mechanism in the $B(C_6F_5)_3$ -Catalyzed Hydrosilylation of Carbonyl Compounds: Implications for the Related Hydrogenation**

Sebastian Rendler and Martin Oestreich*

The precise mechanistic understanding of chemical transformations is an urgent challenge in synthetic chemistry as it guides the targeted design of improved or even novel processes. For example, transition-metal-free reduction of $C=X$ bonds ($X = O$ and NR) catalyzed by boron-based Lewis acids is currently attracting considerable attention.^[1] Illuminating its mechanism(s) of action might open the door for the development of yet unknown enantioselective variants. In this context, commercially available tris(pentafluorophenyl)borane (**1**)^[2] is a particularly effective catalyst for both hydrosilylation and hydrogenation.^[3] In a series of seminal papers, Piers et al. had reported a protocol that is based on triorganosilanes as stoichiometric reducing reagents.^[4] Moreover, Stephan et al. reported that dihydrogen—clearly the most desirable reducing agent—also facilitates smooth turnover in $C=NR$ and $C\equiv N$ reductions.^[5,6] In this scenario, an unconventional $B(C_6F_5)_3$ -catalyzed activation of dihydrogen is operative.^[5] The direct investigation of this dihydrogen activation^[8] by experimentally straightforward techniques is certainly demanding, though.^[9,10] In turn, examination of the closely related silane activation^[11,12] might provide a solid foundation for the delineation of the basic mechanistic principles of both processes.^[13] We report herein a simple yet conclusive investigation of the transition state operative in the $B(C_6F_5)_3$ -catalyzed hydrosilylation of prochiral acetophenone using a silane with a stereogenic silicon center^[14] as a stereochemical probe.^[15,16] We then discuss implications for the related hydrogenation.

Based on comprehensive experimental data, Piers et al. suggested a seemingly counterintuitive three-step mechanism for the hydrosilylation of carbonyl compounds (Scheme 1).^[14a,b] The catalysis commences with the activation of silane **2** by the strong Lewis acid **1** through reversible coordination to the hydridic Si–H bond (**1**→**3**). The two



Scheme 1. The Piers mechanism of the $B(C_6F_5)_3$ -catalyzed hydrosilylation of carbonyl compounds.

resonance structures **3a** and **3b** of the thus-formed intermediate rationalize the capability of **1** to abstract a hydride from silicon in the subsequent step. Silyl transfer to the Lewis basic carbonyl oxygen of **4** thereby produces ion pair **5** (**3**→**5**).


As implied by the complete absence of any crossover when two different mass-labeled silanes are used,^[4b] the ion pair is apparently not solvent-separated and undergoes rapid hydride transfer from the borohydride to the electrophilic carbon of the silylcarboxonium ion to give **6** along with regenerated catalyst **1** (**5**→**1**). The fate of intermediate **3** has remained vague: Concerted S_N2 -type displacement at silicon (S_N2 -Si^[18,19]) of a boron-coordinated hydride by the carbonyl oxygen of **4** has been postulated. Another intriguing piece of information emerges from rapid H/D exchange when **2** and its deuterated congener are used in the absence of Lewis bases.^[4b]

Inspired by remarkable achievements utilizing stereogenicity at silicon as a chiral probe,^[15,20] we decided to examine the nature and consecutive reaction of intermediate **3** by applying our previously developed family of asymmetrically substituted silanes.^[14] We first assessed silanes *rac*-**2a–d** in the hydrosilylation of acetophenone (**4**) in the presence of catalytic amounts of catalyst **1** (5.0 mol %) in order to identify a sufficiently reactive stereogenic silane (**4**→*rac*-**6**, Scheme 2). It is interesting to note that, without exception, all silanes decorated with a *t*Bu group—cyclic *rac*-**2a**,^[21a] cyclic and strained *rac*-**2c**,^[21b] and acyclic *rac*-**2d**^[21c]—were to completely unreactive. In contrast, cyclic *rac*-**2b**^[15b] equipped with an *i*Pr group readily delivered the desired product *rac*-**6b** in good yield and with notable diastereoselection (vide infra). The difference in steric hindrance between *i*Pr-

[*] Dr. S. Rendler, Prof. Dr. M. Oestreich
Organisch-Chemisches Institut
Westfälische Wilhelms-Universität Münster
Corrensstrasse 40, 48149 Münster (Germany)
Fax: (+49) 251-83-36501
E-mail: martin.oestreich@uni-muenster.de
Homepage: http://www.uni-muenster.de/Chemie.oc/research/oestreich/oe_welcome.html

[**] The authors thank Marion Emmert (group of Prof. Dr. Gerhard Erker, Münster) for a sample of $B(C_6F_5)_3$. This research was supported by the Fonds der Chemischen Industrie and the Aventis Foundation (Karl-Winnacker-Stipendium to M.O., 2006–2008).


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200801675>.



Scheme 2. Screening of suitable chiral probes.

substituted *rac*-**2b** and *t*Bu-substituted *rac*-**2a** might account for this dramatic reactivity difference in an assumed S_N2 -Si displacement. However, this argument is inconclusive since Reed et al. had demonstrated that *i*Pr-substituted silanes display a higher tendency to form silylium-ion-type intermediates as a result of α -C-H hyperconjugative stabilization;^[22] these silanes would thus show superior reactivity in an S_N1 -Si-type mechanism.

Strong evidence against a free silylium ion intermediate^[23] would come from the classical reaction setup for a Walden inversion at silicon,^[18] which required repeating the reduction with enantioenriched silane (^{Si}R)-**2b** [**4** $\rightarrow$ ($^{Si}R,R$)-**6b**, Scheme 3].^[24] Prochiral ketone **4** was treated with (^{Si}R)-**2b** (90 % *ee*) in the presence of catalytic amounts of $B(C_6F_5)_3$ to



Scheme 3. Two-step stereochemical analysis: An inversion-retention pathway.


yield optically active ($^{Si}R,R$)-**6b** as a mixture of diastereomers (d.r. 74:26). After racemization-free reductive cleavage of ($^{Si}R,R$)-**6b** with diisobutylaluminum hydride (DIBAL-H) under standard reaction conditions,^[21a,25] silane (^{Si}S)-**2b** was recovered in high chemical yield and with inverted absolute configuration [($^{Si}R,R$)-**6b** $\rightarrow$ (^{Si}S)-**2b**]. With 90 % *ee* for (^{Si}R)-**2b** and 84 % *ee* for (^{Si}S)-**2b**, inversion at the silicon atom is virtually immaculate (97 % inversion and marginal 3 % racemization). Absolute configurations were unambiguously assigned by HPLC analysis on a chiral stationary phase as well as by analysis of the optical rotation. The relative config-

uration of ($^{Si}R,R$)-**6b** was deduced from the configuration of the isolated alcohol (*R*)-**8** (38 % *ee*).

This result clearly supports a concerted S_N2 -Si mechanism for the reduction step. Spontaneous heterolytic dissociation of intermediate **3** (Scheme 1) would liberate a free (or toluene-stabilized^[26]) achiral silylium ion, thereby producing racemic material. Nucleophilic attack at silicon by the Lewis basic carbonyl oxygen of **4** must occur *anti* to the quasi-linear B-H-Si array^[4b,27] via transition state **7** (Scheme 3). Moreover, we emphasize that **7** should be regarded as a transition state because any lifetime on the reaction timescale would likely result in racemization by pseudorotational processes like those observed with hypervalent cyclic silicon intermediates.^[28]

The origin of diastereoselection is noteworthy as it is induced by the single-point-bound stereogenic silicon atom coordinated to the Lewis basic carbonyl oxygen (**5**, Scheme 1). Piers et al. had already presented solid data that it is the borohydride and not another molecule of the silane that functions as the reducing agent.^[4b] Overall, the reaction studied here represents an example of chirality transfer from silicon to carbon,^[29] which follows a “one-silicon” as opposed to a “two-silicon” cycle.^[15]

To further exclude involvement of free silylium ions, we performed another rigorous control experiment. An equimolar mixture of enantioenriched silane (^{Si}R)-**2b** (90 % *ee*) and deuterium-labeled achiral silane [2H]-**9** was exposed to the borane catalyst **1** in the absence of a Lewis base (Scheme 4).^[30] Mass spectrometric analysis of the isotopic



Scheme 4. Racemization-free scrambling in the absence of a Lewis base: No support for free silylium ions [$B = B(C_6F_5)_3$ and Si or $Si^* = R_3Si$].

distribution after 2 h at room temperature showed complete scrambling: (^{Si}R)-[$^1H/^2H$]-**2b** (89 % *ee*, H/D 54:46) and [2H]-**9** (H/D 50:50). The preservation of the stereochemical integrity in isolated (^{Si}R)-[$^1H/^2H$]-**2b** excludes: 1) a mechanism through (achiral) silylium ion intermediates and 2) an S_N2 -Si displacement at activated **3** with a silane as the attacking nucleophile as both of these would bring about racemization. It is therefore plausible to suggest a σ -bond metathesis involving a four-centered cyclic transition state. Related transition-metal-catalyzed processes are known to proceed with stereoretention at silicon.^[14,15,31] Transition-state **10** (Scheme 4) consisting of two borane-activated silanes fulfills the stereochemical requirements; alternatively, a complex of

type **3** (Scheme 1) might be sufficiently reactive to directly undergo σ -bond metathesis with a free silane.

Based on these insights into the $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed hydrosilylation, conclusions might be drawn, to a certain extent, for the closely related hydrogenation.^[10] A comparison of the bond energies of $\text{R}_3\text{Si-H}$ (90 kcal mol⁻¹^[32]) and H-H (108 kcal mol⁻¹^[8]) corroborates that $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed heterolytic dissociation of H_2 into the contact ion pair " $\text{H}^+[\text{HB}(\text{C}_6\text{F}_5)_3]^-$ " is, as is the case for $\text{R}_3\text{Si-H}$, an unfavorable event in the absence of a Lewis base. However, an $\text{S}_{\text{N}}2$ -type process at hydrogen similar to that at silicon is implicated:^[10] η^1 (end-on) coordination^[27] of $\text{B}(\text{C}_6\text{F}_5)_3$ to dihydrogen could activate the H-H bond for nucleophilic attack by the Lewis basic substrate. Again, a tight contact ion pair consisting of an iminium ion (with imines as substrates^[5]) and the borohydride is formed. Liberation of the amine after subsequent hydride transfer then completes the catalytic cycle.

In summary, the refined mechanistic picture of the $\text{B}(\text{C}_6\text{F}_5)_3$ -catalyzed hydrosilylation bodes well for the development of catalytic asymmetric approaches. It seems that their efficiency will highly depend on asymmetric induction of a chiral nonracemic borane.^[3,33] A proof of principle was included in a very recent report, which might also stimulate further research in this area.^[5b]

Our mechanistic investigation and proof of an $\text{S}_{\text{N}}2$ -Si transition state in the $\text{Si} \rightarrow \text{B}$ hydride-transfer step is predicated on a simple Walden-type analysis employing a silane having a silicon stereocenter as a stereochemical probe. It might well be worth applying this technique to systems relying on other Lewis bases, e.g., phosphines,^[9] and to the recent dihydrogen-splitting molecules introduced by Stephan et al.^[6,7,34]

Received: April 9, 2008

Published online: July 9, 2008

Keywords: hydrosilylation · Lewis acids · nucleophilic substitution · reaction mechanisms · silicon

- [1] A. L. Kenward, W. E. Piers, *Angew. Chem.* **2008**, *120*, 38–42; *Angew. Chem. Int. Ed.* **2008**, *47*, 38–41.
- [2] A. G. Massey, A. J. Park, *J. Organomet. Chem.* **1964**, *2*, 245–250.
- [3] a) G. Erker, *Dalton Trans.* **2005**, 1883–1890; b) W. E. Piers, *Adv. Organomet. Chem.* **2005**, *52*, 1–76; c) W. E. Piers, T. Chivers, *Chem. Soc. Rev.* **1997**, *26*, 345–354.
- [4] a) D. J. Parks, W. E. Piers, *J. Am. Chem. Soc.* **1996**, *118*, 9440–9441; b) D. J. Parks, J. M. Blackwell, W. E. Piers, *J. Org. Chem.* **2000**, *65*, 3090–3098; c) J. M. Blackwell, E. R. Sonmor, T. Scoccitti, W. E. Piers, *Org. Lett.* **2000**, *2*, 3921–3923.
- [5] a) P. A. Chase, T. Jurca, D. W. Stephan, *Chem. Commun.* **2008**, 1701–1703; b) D. Chen, J. Klankermayer, *Chem. Commun.* **2008**, 2130–2131.
- [6] For the metal-free hydrogenation of imines and nitriles using the Stephan catalyst^[7] ($\text{R}_2\text{PC}_6\text{F}_4\text{B}(\text{C}_6\text{F}_5)_2$ systems, $\text{R} = 2,4,6\text{-Me}_3\text{C}_6\text{H}_2$ or $t\text{Bu}$), see: P. A. Chase, G. C. Welch, T. Jurca, D. W. Stephan, *Angew. Chem.* **2007**, *119*, 8196–8199; *Angew. Chem. Int. Ed.* **2007**, *46*, 8050–8053.
- [7] For the landmark paper, see: G. C. Welch, R. R. San Juan, J. D. Masuda, D. W. Stephan, *Science* **2006**, *314*, 1124–1126.
- [8] a) G. J. Kubas, *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 6901–6907; b) J. W. Tye, M. B. Hall in *Activation of Small Molecules* (Ed.: W. B. Tolman), Wiley-VCH, Weinheim, **2006**, pp. 126–158.
- [9] Attempts to spectroscopically observe $\text{B}(\text{C}_6\text{F}_5)_3$ -dihydrogen adducts have been unfruitful so far: G. C. Welch, D. W. Stephan, *J. Am. Chem. Soc.* **2007**, *129*, 1880–1881.
- [10] For an excellent quantum-chemical investigation including a plausible mechanism, see: T. A. Rokob, A. Hamza, A. Stirling, T. Soós, I. Pápai, *Angew. Chem.* **2008**, *120*, 2469–2472; *Angew. Chem. Int. Ed.* **2008**, *47*, 2435–2438.
- [11] M. S. Eisen in *The Chemistry of Organic Silicon Compounds*, Vol. 3 (Eds.: Z. Rappoport, Y. Apeloig), Wiley, New York, **2001**, pp. 2037–2128.
- [12] S. Rendler, M. Oestreich in *Modern Reduction Methods* (Eds.: P. G. Andersson, I. J. Munslow), Wiley-VCH, Weinheim, **2008**, pp. 183–207.
- [13] A relationship of that is also observed in Cu-H-catalyzed reductions: a) S. Rendler, M. Oestreich, *Angew. Chem.* **2007**, *119*, 504–510; *Angew. Chem. Int. Ed.* **2007**, *46*, 498–504; b) B. H. Lipshutz in *Modern Organocopper Chemistry* (Ed.: N. Krause), Wiley-VCH, Weinheim, **2002**, pp. 167–187; c) J. M. Stryker, W. S. Mahoney, J. F. Daeuble, D. M. Bestrensky in *Catalysis of Organic Reactions* (Ed.: W. E. Pascoe), Marcel Dekker, New York, **1992**, pp. 29–44.
- [14] M. Oestreich, *Synlett* **2007**, 1629–1643.
- [15] a) M. Oestreich, S. Rendler, *Angew. Chem.* **2005**, *117*, 1688–1691; *Angew. Chem. Int. Ed.* **2005**, *44*, 1661–1664; b) S. Rendler, M. Oestreich, C. P. Butts, G. C. Lloyd-Jones, *J. Am. Chem. Soc.* **2007**, *129*, 502–503.
- [16] We explicitly note that a recent report on the supposedly mechanistically related dehydrogenative silylation of alcohols^[17] also included an inconclusive observation—substantial racemization at silicon—for the hydrosilylation of carbonyl compounds with a silicon-stereogenic silane: S. Shinke, T. Tsuchimoto, Y. Kawakami, *Silicon Chem.* **2007**, *3*, 243–249.
- [17] J. M. Blackwell, K. L. Foster, V. H. Beck, W. E. Piers, *J. Org. Chem.* **1999**, *64*, 4887–4892.
- [18] a) L. H. Sommer, *Stereochemistry, Mechanism, and Silicon*, McGraw-Hill, New York, **1965**; b) L. H. Sommer, *Intra-Sci. Chem. Rep.* **1973**, *7*, 1–44; c) R. J. P. Corriu, C. Guerin, J. J. E. Moreau in *Topics in Stereochemistry*, Vol. 15 (Ed.: E. L. Eliel), Wiley, New York, **1984**, pp. 355–374.
- [19] A. P. Bento, F. M. Bickelhaupt, *J. Org. Chem.* **2007**, *72*, 2201–2207.
- [20] Selected examples: a) A. G. Brook, *Acc. Chem. Res.* **1974**, *7*, 77–84; b) V. Bagheri, M. P. Doyle, J. Taunton, E. E. Claxton, *J. Org. Chem.* **1988**, *53*, 6158–6150; c) M. P. Doyle, K. G. High, V. Bagheri, R. J. Pieters, P. J. Lewis, M. M. Pearson, *J. Org. Chem.* **1990**, *55*, 6082–6086; d) K. Tomooka, A. Nakazaki, T. Nakai, *J. Am. Chem. Soc.* **2000**, *122*, 408–409; e) A. Nakazaki, T. Nakai, K. Tomooka, *Angew. Chem.* **2006**, *118*, 2293–2296; *Angew. Chem. Int. Ed.* **2006**, *45*, 2235–2238.
- [21] a) S. Rendler, G. Auer, M. Keller, M. Oestreich, *Adv. Synth. Catal.* **2006**, *348*, 1171–1182; b) S. Rendler, O. Plefka, B. Karatas, G. Auer, R. Fröhlich, C. Mück-Lichtenfeld, S. Grimme, M. Oestreich, unpublished results; c) G. L. Larson, E. Torres, *J. Organomet. Chem.* **1985**, *293*, 19–27.
- [22] Z. Xie, R. Bau, A. Benesi, C. A. Reed, *Organometallics* **1995**, *14*, 3933–3941.
- [23] For leading references on this controversial debate, see: a) R. J. P. Corriu, M. Henner, *J. Organomet. Chem.* **1974**, *74*, 1–28; b) J. B. Lambert, L. Kania, S. Zhang, *Chem. Rev.* **1995**, *95*, 1191–1201; c) J. B. Lambert, Y. Zhao, *Angew. Chem.* **1997**, *109*, 389–391; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 400–401; d) C. A. Reed, *Acc. Chem. Res.* **1998**, *31*, 325–332; e) K.-C. Kim, C. A. Reed, D. W. Elliott, L. J. Mueller, F. Tham, L. Lin, J. B. Lambert, *Science* **2002**, *297*, 825–827; f) P. P. Gaspar, *Science*

- 2002**, 297, 785–786; g) T. Müller, *Adv. Organomet. Chem.* **2005**, 53, 155–216.
- [24] For an early debate on silylium intermediates and stereogenic silicon, see: a) J. D. Austin, C. Eaborn, *J. Chem. Soc.* **1964**, 2279–2280; b) L. H. Sommer, D. L. Bauman, *J. Am. Chem. Soc.* **1969**, 91, 7076–7078.
- [25] Stereospecificity of reduction with DIBAL-H is secured: M. Oestreich, G. Auer, M. Keller, *Eur. J. Org. Chem.* **2005**, 184–195.
- [26] G. A. Olah, G. Rasul, X.-Y. Li, H. A. Buchholz, G. Sanford, G. K. S. Prakash, *Science* **1994**, 263, 983–984.
- [27] For a transition-metal η^1 -silane complex, see: J. Yang, P. S. White, C. K. Schauer, M. Brookhart, *Angew. Chem.* **2008**, 120, 4209–4211; *Angew. Chem. Int. Ed.* **2008**, 47, 4141–4143.
- [28] a) E. P. A. Couzijn, M. Schakel, F. J. J. de Kanter, A. W. Ehlers, M. Lutz, A. L. Spek, K. Lammertsma, *Angew. Chem.* **2004**, 116, 3522–3524; *Angew. Chem. Int. Ed.* **2004**, 43, 3440–3442; b) E. P. A. Couzijn, A. W. Ehlers, M. Schakel, K. Lammertsma, *J. Am. Chem. Soc.* **2006**, 128, 13634–13639.
- [29] M. Oestreich, *Chem. Eur. J.* **2006**, 12, 30–37.
- [30] Treatment of (^{Si}R)-**2b** (90% ee) with **1** in toluene resulted in no racemization as well. See the Supporting Information for details.
- [31] S. Rendler, G. Auer, M. Oestreich, *Angew. Chem.* **2005**, 117, 7793–7797; *Angew. Chem. Int. Ed.* **2005**, 44, 7620–7624.
- [32] J. Y. Corey in *The Chemistry of Organic Silicon Compounds* (Eds.: S. Patai, Z. Rappoport), Wiley, New York, **1989**, pp. 1–56.
- [33] K. Ishihara, H. Yamamoto, *Eur. J. Org. Chem.* **1999**, 527–538.
- [34] a) G. C. Welch, L. Cabrera, P. A. Chase, E. Hollink, J. D. Masuda, P. Wei, D. W. Stephan, *Dalton Trans.* **2007**, 3407–3414; b) P. Spies, G. Erker, G. Kehr, K. Bergander, R. Fröhlich, S. Grimme, D. W. Stephan, *Chem. Commun.* **2007**, 5072–5074.