


Divergent Titanium-Mediated Allylations with Modulation by Nickel or Palladium**

Araceli G. Campaña, Btissam Bazdi, Noelia Fuentes, Rafael Robles, Juan M. Cuerva,*
J. Enrique Oltra,* Susana Porcel, and Antonio M. Echavarren*

Titanocene(III)-mediated radical processes are important tools for the formation of C–C bonds under mild conditions, and are compatible with many functional groups.^[1,2] Moreover, titanium(III) complexes can be used substoichiometrically,^[3] which has allowed the development of enantioselective versions of these reactions.^[4]

A serious limitation of these radical processes, however, derives from the fact that titanium(III)-mediated radical generation requires reactive substrates, such as allylic halides, which are often cumbersome in introduction and manipulation. Allylic carbonates and carboxylates, in contrast, are easily prepared and handled but are inert against titanocene(III) complexes. Nevertheless, it is known that nickel and palladium complexes can readily activate allylic carbonates and carboxylates **I** (Scheme 1) to form η^3 -allylmethyl complexes (**II**). On the basis of these results, we deemed that the combination of palladium or nickel derivatives with titanocene(III) complexes would facilitate the development of novel allylation processes using accessible allyl carbonates or carboxylates.

In the case of palladium catalysis, it is known that the Oppolzer-type cyclization of organometallic species (**II**, M = Pd, Scheme 1) to cyclic derivatives (**VI**) is relatively slow at room temperature.^[5] Thus, reduction of **II** by a single-electron-transfer reagent, such as $[\text{Cp}_2\text{TiCl}]$,^[6] could lead to the allylic radical **III**, which might be eventually trapped by a second $[\text{Cp}_2\text{TiCl}]$ species to give an allylic titanium(IV) complex **IV**. Finally, nucleophilic attack of the organometallic derivative **IV** on an aldehyde or other electrophilic reagent would provide the corresponding allylation product **V**.



Scheme 1. Mechanism for palladium-catalyzed, titanocene-mediated allylation of carbonyl compounds and nickel-catalyzed, titanocene-mediated carbocyclization of allylic carboxylates. [M] = transition metal catalyst; Cp = cyclopentadienyl; E⁺ = electrophilic reagent (e.g. aldehyde). R = OEt, Me, or Ph.

On the other hand, nickel-catalyzed carbocyclizations, via intermediates such as **II** (M = Ni, Scheme 1) to cyclic derivatives **VI**, are relatively fast at room temperature.^[7] Once formed, **VI** might be reduced by $[\text{Cp}_2\text{TiCl}]$ to a primary radical **VII**, which could be trapped by a second $[\text{Cp}_2\text{TiCl}]$ species to give an alkyl titanium(IV) complex **VIII**. Hydrolysis of the organometallic derivative **VIII** would yield carbocycles **IX**. Thus, we anticipated that the use of palladium or nickel catalysts could modulate titanium(III) to drive allylation reactions with allyl carboxylates by two different pathways, either through intermolecular coupling with electrophilic reagents or to give carbocyclic derivatives by an intramolecular allylation.


To check our hypothesis, we chose allylic carbonate **E-1** as a model allylation reagent. Thus, reaction of decanal with carbonate **E-1** and an excess of $[\text{Cp}_2\text{TiCl}]$ (2.0 equiv),^[8] in the presence of PdCl_2 (20 mol %) and triphenylphosphine,^[9] gave the expected coupling product **2** as a single stereoisomer in 76 % yield (Scheme 2).^[10] In contrast, when carbonate **E-1** was treated with an excess of $[\text{Cp}_2\text{TiCl}]$ (2.0 equiv) in the presence of $[\text{NiCl}_2(\text{PPh}_3)_2]$ (20 mol %), a mixture of carbocycles **3** and **4** (4:1 ratio) was obtained in almost quantitative

[*] A. G. Campaña, B. Bazdi, N. Fuentes, Dr. R. Robles, Dr. J. M. Cuerva, Prof. J. E. Oltra
Department of Organic Chemistry, Faculty of Sciences
University of Granada
18071 Granada (Spain)
E-mail: jmcuerva@ugr.es
joltra@ugr.es

S. Porcel, Prof. A. M. Echavarren
Institute of Chemical Research of Catalonia (ICIQ)
Av. Països Catalans 16, 43007 Tarragona (Spain)
Fax: (+34) 977-920-225
E-mail: aechavarren@iciq.es

[**] We thank the "Junta de Andalucía" (research group FQM339), the Spanish MEC (projects CTQ2005-08402/BQU, CTQ2007-60745/BQU, Consolider Ingenio 2010, and predoctoral fellowship to A.G.C.), the UAM (predoctoral fellowship to S.P.), and the ICIQ Foundation for financial support.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200802520>.



Scheme 2. Palladium-catalyzed formation of coupling product **2** and nickel-catalyzed formation of carbocycles **3** and **4** from **E-1**

yield. The *cis*-configuration of **3** was in accordance with that previously described for Oppolzer cyclizations mediated by nickel.^[11] When we repeated this experiment in the presence of decanal (2.0 equiv), we obtained similar results (4:1 mixture of **3** and **4**, 94% yield), whereas decanal was recovered unchanged. Moreover, control experiments showed that, in the absence of either the transition metal or the titanium complex, the above reactions did not occur, strongly supporting our hypothesis.

We subsequently assayed the palladium-catalyzed allylation of different carbonyl compounds with allylic carbonates and stoichiometric (method A) or substoichiometric (method B) amounts of titanium (Table 1).^[8] The titanocene-regenerating agent used in method B and developed in our laboratory, was a combination of manganese dust, 2,4,6-collidine and trimethylsilyl chloride.^[12] Thus, we achieved the allylation of different aldehydes and ketones with allyl carbonates, obtaining moderate to excellent yields of the expected homoallylic alcohols. It is worth noting that, although palladium-based methods for the allylation of aldehydes with allyl carboxylates are known,^[13] the allylation of ketones is less common and usually requires very long reaction times.^[13a,14] In this case, however, ketone allylation proceeded in a few hours at room temperature and even ketones prone to undergo pinacol coupling, such as acetophenone and tetralone, proved to be suitable substrates. Moreover, alkyl substitution at the allylic carbonate was

well-tolerated, yielding the corresponding γ -addition products (Table 1, entries 11–19). It should be noted that both isomers **E-1** and **Z-1**, as well as **E-22** and **Z-22**, led to the same stereoisomer (**2** and **23** respectively), thus demonstrating the stereoconvergent nature of the process. The reaction is also chemoselective: esters (ethyl benzoate) and nitriles (benzonitrile) were both inert under our conditions. Moreover, yields obtained from palladium/titanium-catalyzed allylations (method B) were similar and, in some cases, even better than those from the process using stoichiometric titanium.^[8] Stereoselectivity was also comparable for both methods, suggesting closely related reaction mechanisms for both procedures.

The results summarized in Table 1 are consistent with the mechanism depicted in Scheme 1, starting with the oxidative addition of the palladium complex to the allylic carbonate to give an η^3 -allyl palladium intermediate, which undergoes subsequent reduction to an allyl radical (such as **III**) by single-electron transfer from $[\text{Cp}_2\text{TiCl}]$.^[15] Another $[\text{Cp}_2\text{TiCl}]$ species then captures the allyl radical to form a nucleophilic allyl titanium(IV) complex, which reacts with the carbonyl compound.^[4b,16] This mechanistic hypothesis was strongly sup-

Table 1: Palladium/titanium-promoted (method A) and catalyzed (method B) allylation of carbonyl compounds with allylic carbonates.^[a]

Entry	Allyl carbonate	Carbonyl compound	Product	Method, Yield [%]
1		decanal		B, 75
2		3-phenyl propanal		B, 54
3		citronellal		B, 73 ^[b]
4		2-decanone		A, 85
5		adamantanone		B, 79
6		acetophenone		B, 74
7		tetralone		B, 74
8		acetyl ferrocene		A, 99

Table 1: (Continued)


Entry	Allyl carbonate	Carbonyl compound	Product	Method, Yield [%]
9		cyclododecanone		B, 78
10		<i>t</i> Butyl-cyclohexanone		B, 65 ^[c]
11		decanal		B, 58 ^[d]
12		2-decanone		B, 75 ^[e]
13		2-decanone		B, 55
14		decanal		B, 52
15		decanal		A, 63 ^[f]
16		—		A, 73 ^[b]
17		—		A, 74 ^[b]
18		—		B, 95
19		—		B, 92

[a] Method A: Carbonyl compound (1 mmol), allyl carbonate (4 mmol), [Cp₂TiCl₂] (2 mmol), PdCl₂ (0.2 mmol), PPh₃ (0.4 mmol), Mn (8 mmol). Method B: Carbonyl compound (1 mmol), allyl carbonate (4 mmol), [Cp₂TiCl₂] (0.4 mmol), PdCl₂ (0.2 mol), PPh₃ (0.4 mmol), Mn (8 mmol), Me₃SiCl (4 mmol), and 2,4,6-collidine (7 mmol). [b] 1:1 mixture of stereoisomers. [c] 2:1 mixture of *cis:trans* stereoisomers. [d] 3:7 mixture of *syn:anti* stereoisomers. [e] 3:2 mixture of stereoisomers. [f] single stereoisomer, tentatively assigned as the *anti* stereoisomer.

ported by the formation of product **2** when the previously prepared palladium complex **24**^[5b] was treated with decanal in the presence of PPh₃ and [Cp₂TiCl] (Scheme 3).^[17] Moreover, in the absence of either titanium or phosphines, the allylation reaction did not occur.

The nickel/titanium-based procedure afforded Oppolzer-type cyclizations (Table 2) at room temperature, under conditions substantially milder than those previously reported for this type of cyclization.^[5a] We also found that the ratio of carbocycles **3** and **4** obtained was dependent on the relative ratio of titanium and nickel catalysts employed.^[8] Cyclic compound **3** probably derives from the reduction of alkyl nickel **VI** by [Cp₂TiCl] (see Scheme 1), which is faster than any potential β-hydride elimination process. Thus, its production is maximized in the presence of high ratios of [Cp₂TiCl]. In fact, using 4 equivalents of [Cp₂TiCl] and only 0.1 equivalents of [NiCl₂(PPh₃)₂], we obtained almost exclusively cyclopentane **3** (**3:4** = 97:3) in good yield (75 %) (Table 2, entry 2). On the other hand, using substoichiometrically [Cp₂TiCl] (0.1 equiv) and [NiCl₂(PPh₃)₂] (0.2 equiv) substantially increased the proportion of exocyclic alkene **4** (**3:4**, 30:70; 60 % overall yield).^[8,18]

Different carbocycles and heterocycles were obtained in moderate to good yield by this method, with complete regiochemical control (Table 2). Excellent stereoselectivity was also obtained, affording exclusively products with *cis* configuration, with the only exception being **38** (9:1 *cis:trans*). Nickel/titanium-promoted cyclization of carbonates **37** and **39** stereoselectively led to **38** and **40** respectively, both products containing three stereogenic centers (Table 2, entries 9 and 10).^[19] Allylic acetates assayed under our conditions (Table 2, entries 3, 7, and 8) showed similar behavior to that of carbonates. In fact, nickel/titanium-induced cyclizations of acetate **27** (Table 2,



Scheme 3. Formation of adduct **2** from palladium complex **24** ($Z = C(CO_2Me)_2$, OTFA = trifluoroacetate).

Table 2: Nickel/titanium-promoted intramolecular allylation of alkenes (Oppolzer-type cyclization) with allylic carbonates.^[a]

Entry	Allylic substrate	Product	Yield [%]
1		3 + 4 (4:1)	98
2	Z-1 $Z = C(CO_2Me)$	3 + 4 (97:3)	75 ^[c]
3		3 + 4 (4:1)	98
4	26 $Z = C(CO_2Me)$		95
5	27 $Z = NSO_2Ph$	28	59
6	29 $Z = NSO_2Ph$	30	69 ^[d]
7	31 $Z = NTs$	32	75
8 ^[b]	33 $Z = C(CO_2Me)$	34	71
9	35 $Z = NSO_2Ph$	36	83 ^[e]
10	37 $Z = NSO_2Ph$	38	81
	39	40	

[a] Allyl carbonate or acetate (1 mmol), $[Cp_2TiCl_2]$ (2 mmol), $NiCl_2(PPh_3)_2$ (0.2 mmol), Mn (8 mmol). [b] **33** is a 5.4:1 mixture of *E*:*Z* diastereoisomers. [c] $[Cp_2TiCl_2]$ (4.0 mmol) was used. [d] Up to 50% of the related product derived from a β -hydride elimination process was obtained. [e] 10% of the product obtained was the stereoisomer with the opposite configuration of the vinyl group.

entry 3) and both stereoisomers *E*-**1** (Scheme 2) and *Z*-**1** (Table 2, entry 1) led to the same products, confirming the versatility of this method and revealing the stereoconvergent nature of the process. Additionally, the reaction proved to be compatible with different functional groups, including esters, ethers, and sulfonamides and permitted different substitution patterns in the involved alkenes.^[20] However, six-membered rings were not obtained under these reaction conditions.

In summary, we have demonstrated for the first time that late transition metals palladium and nickel are capable of modulating titanium(III) reactivity to achieve selective allylations of either carbonyl compounds or alkenes respectively, using allylic carbonates and carboxylates, instead of cumbersome allyl halides, as allylation reagents. The palladium/titanium-catalyzed allylation of ketones (method B) is an especially convenient procedure, as evidenced by the excellent diastereoselectivity obtained under mild conditions. The catalytic nature of this process should facilitate the development of enantioselective versions. Nickel/titanium-promoted Oppolzer-type cyclizations provided polyalkyl substituted carbo- and heterocycles at room temperature, under conditions compatible with several functional groups, and with excellent stereoselectivity. Moreover, by modifying the relative ratios of nickel and titanium employed, the final step of the cyclization process can be either a reductive (*cis*-substituted products)^[21] or a nonreductive (exocyclic alkenes) process. A more in-depth study of the reaction mechanisms involved in these processes and the development of enantioselective versions are underway.

Received: May 29, 2008

Published online: August 28, 2008

Keywords: allylation · carbocycles · nickel · palladium · titanium

- [1] For a seminal work on titanocene(III) chemistry see: T. V. Rajan Babu, W. A. Nugent, *J. Am. Chem. Soc.* **1994**, *116*, 986–997.
- [2] Recent reviews: a) J. M. Cuerva, J. Justicia, J. L. Oller-López, J. E. Oltra, *Top. Curr. Chem.* **2006**, *264*, 63–91; b) A. Gansäuer, J. Justicia, C.-A. Fan, D. Worgull, F. Piester, *Top. Curr. Chem.* **2007**, *279*, 25–52.
- [3] Selected references: a) A. Gansäuer, H. Bluhm, M. Pierobon, *J. Am. Chem. Soc.* **1998**, *120*, 12849–12859; b) J. Justicia, A. Rosales, E. Buñuel, J. L. Oller-López, M. Valdivia, A. Haïdour, J. E. Oltra, A. F. Barrero, D. J. Cárdenas, J. M. Cuerva, *Chem. Eur. J.* **2004**, *10*, 1778–1788; c) J. Justicia, J. L. Oller-López, A. G. Campaña, J. E. Oltra, J. M. Cuerva, E. Buñuel, D. J. Cárdenas, *J. Am. Chem. Soc.* **2005**, *127*, 14911–14921; d) J. M. Cuerva, A. G. Campaña, J. L. Oller-López, J. Justicia, A. Rosales, R. Robles, E. Buñuel, D. J. Cárdenas, J. E. Oltra, *Angew. Chem.* **2006**, *118*, 5648–5652; *Angew. Chem. Int. Ed.* **2006**, *45*, 5522–5526.
- [4] a) A. Gansäuer, H. Bluhm, B. Rinker, S. Narayan, M. Schick, T. Lauterbach, M. Pierobon, *Chem. Eur. J.* **2003**, *9*, 531–542; b) A. Rosales, J. L. Oller-López, J. Justicia, A. Gansäuer, J. E. Oltra, J. M. Cuerva, *Chem. Commun.* **2004**, 2628–2629; c) R. E. Estévez, J. L. Oller-López, R. Robles, C. R. Melgarejo, A. Gansäuer, J. M. Cuerva, J. E. Oltra, *Org. Lett.* **2006**, *8*, 5433–5436.

- [5] a) W. Oppolzer in *Comprehensive Organic Synthesis*, Vol. 5 (Eds.: B. M. Trost, I. Fleming), Pergamon, Oxford, **1991**, chap. 1.2; b) E. Gómez-Bengoa, J. M. Cuerva, A. M. Echavarrén, G. Martorell, *Angew. Chem.* **1997**, *109*, 795–797; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 767–769.
- [6] R. J. Enemærke, J. Larsen, T. Skrydstrup, K. Daasbjerg, *J. Am. Chem. Soc.* **2004**, *126*, 7853–7864.
- [7] W. Oppolzer, M. Bedoya-Zurita, C. Y. Switzer, *Tetrahedron Lett.* **1988**, *29*, 6433–6436.
- [8] For experimental details see Supporting Information.
- [9] Addition of triphenylphosphine considerably increased the yields, possibly stabilizing palladium intermediates.
- [10] The *anti* configuration of **2** was tentatively assigned on the basis of the stereochemical behavior evident in the closely related allylation of decanal with (*E*)-but-2-enyl ethyl carbonate promoted by Pd/Ti (Table 1, entry 11).
- [11] W. Oppolzer, T. H. Keller, M. Bedoya-Zurita, C. Stone, *Tetrahedron Lett.* **1989**, *30*, 5883–5886.
- [12] A. F. Barrero, A. Rosales, J. M. Cuerva, J. E. Oltra, *Org. Lett.* **2003**, *5*, 1935–1938.
- [13] Selected references: a) T. Tabuchi, J. Inanaga, M. Yamaguchi, *Tetrahedron Lett.* **1986**, *27*, 1195–1196; b) Y. Masuyama, K. Otake, Y. Kurusu, *Tetrahedron Lett.* **1988**, *29*, 3563–3566; c) S. Sebelius, O. A. Wallner, K. J. Szabó, *Org. Lett.* **2003**, *5*, 3065–3068; d) S.-F. Zhu, Y. Yang, L.-X. Wang, B. Liu, Q.-L. Zhou, *Org. Lett.* **2005**, *7*, 2333–2335.
- [14] a) K. Yasui, Y. Goto, T. Yajima, Y. Taniseki, K. Fugami, A. Tanaka, Y. Tamaru, *Tetrahedron Lett.* **1993**, *34*, 7619–7622; b) W. Qiu, Z. Wang, *J. Chem. Soc. Chem. Commun.* **1989**, 356–357; c) J. P. Takahara, Y. Masuyama, Y. Kurusu, *J. Am. Chem. Soc.* **1992**, *114*, 2577–2586; d) H. Nakamura, N. Asao, Y. Yamamoto, *Collect. Czech. Chem. Commun.* **1995**, 1273–1274.
- [15] We also noted that, in the absence of decanal, allylic carbonates can be dimerized, indicating that allyl radicals are involved in the palladium-mediated allylation process. Wurtz-type coupling products are the main reaction products if allyl radicals cannot be trapped, owing to a low concentration of [Cp₂TiCl]. Thus we carried out the Pd⁰/Ti^{III}-promoted Wurtz-type reaction of farnesyl ethyl carbonate, obtaining squalene (81 %) with acceptable regio- and stereoselectivity. For details, see Supporting Information.
- [16] For selected references of closely related nucleophilic allyltitanium complexes: a) A. Kasatkin, T. Nakagawa, S. Okamoto, F. Sato, *J. Am. Chem. Soc.* **1995**, *117*, 3881–3882; b) F. Sato, S. Iijima, M. Sato, *Tetrahedron Lett.* **1981**, *22*, 243–246.
- [17] In any case, alternative mechanisms based on nucleophilic allylpalladium complexes cannot be completely ruled out; see: a) H. Nakamura, N. Asao, Y. Yamamoto, *J. Chem. Soc. Chem. Commun.* **1995**, 1273–1274; b) O. A. Wallner, K. J. Szabó, *J. Org. Chem.* **2003**, *68*, 2934–2943; c) N. Solin, J. Kjellgren, K. Szabo, *J. Am. Chem. Soc.* **2004**, *126*, 7026–7033; d) Review: G. Zanoni, A. Pontiroli, A. Marchetti, G. Vidari, *Eur. J. Org. Chem.* **2007**, 3599–3611.
- [18] Exocyclic alkene **4** is reminiscent of products obtained in the Oppolzer cyclization. It should be noted, however, that nickel catalysts have seldom been used for this transformation, possibly because it is not easy to reintroduce them into a catalytic cycle; see: B.-L. Lin, L. Liu, Y. Fu, S.-W. Luo, Q. Chen, Q.-X. Guo, *Organometallics* **2004**, *23*, 2114–2123.
- [19] This excellent stereoselectivity is considerably higher than that reported for related cyclizations based on titanium(II) and zirconium(II)-complexes; see: a) T. Takahashi, D. Y. Kondakov, N. Suzuki, *Organometallics* **1994**, *13*, 3411–3412; b) Y. Takayama, Y. Gao, F. Sato, *Angew. Chem.* **1997**, *109*, 890–892; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 851–853; c) Y. Takayama, S. Okamoto, F. Sato, *Tetrahedron Lett.* **1997**, *38*, 8351–8354; d) A. D. Campbell, T. M. Raynham, R. J. K. Taylor, *Chem. Commun.* **1999**, 245–246.
- [20] Previously reported procedures to prepare similar carbocycles afforded lower yields and poorer chemical compatibility; see: a) W. Oppolzer, F. Schröder, *Tetrahedron Lett.* **1994**, *35*, 7939–7942; b) Y.-J. Song, I. G. Jung, H. Lee, Y. T. Lee, Y. K. Chung, H.-Y. Jang, *Tetrahedron Lett.* **2007**, *48*, 6142–6146.
- [21] It is worth noting that Bu₃SnH-induced radical cyclization of an allylic bromide closely related to carbonate **1** provided considerably worse regio- (91:9 *5-exo/6-endo*) and stereoselectivity (*cis/trans*, 65:35) than those afforded by the Ni/Ti system; see: G. Stork, M. E. Reynolds, *J. Am. Chem. Soc.* **1988**, *110*, 6911–6913.