


Isoxazole-Assisted Direct Substitution of the Hydroxy Group in α -Ketols: Introduction of Angular Substituents in a Polycyclic System**

Hiroshi Takikawa, Katsuyoshi Hikita, and Keisuke Suzuki*

The type-II polyketide biosynthetic pathway produces a vast array of polycyclic natural products such as **1**^[1] and **2**,^[2] attracting the attention of the biochemical and synthetic communities.^[3] In our continued synthetic endeavors towards these compounds, we searched for viable methods for introducing the characteristic angular substituents (Scheme 1).^[4] We envisaged α -ketol **A**^[5] as a key platform to achieve this goal, hoping that the isoxazole moiety would serve not only as a 1,3-dicarbonyl equivalent,^[6] but also as a




Scheme 1. Approach to polyketide-derived polycyclic natural products with angular substituents.

“guiding functionality” to install an angular substituent (**A** → **B**). This installation can hopefully be facilitated by the recently discovered ability of isoxazoles to stabilize the α -cationic species **C**.^[7]

By recognizing these aspects, we reasoned that ketol **3** would be amenable to the direct S_N1 -type substitution of the


hydroxy group (Scheme 2).^[8] Although formation of a cation next to a carbonyl group is disfavored,^[9] we hoped that the isoxazole unit would override the deficit. If viable, the process would become a straightforward approach to angularly substituted polycyclic natural products, such as **1** and **2**.



Scheme 2. Direct installation of an angular substituent. Bn = benzyl.

Herein, we report the affirmative answer to this challenge, that is, the isoxazole-assisted, direct substitution of α -ketol **3** with various nucleophiles to therefore allow the installation of angular substituents in polycyclic systems.

The initial feasibility study is illustrative [Eq. (1)]: Upon treatment of enantiomerically enriched α -ketol (*R*)-**3** (98% *ee*)^[5] with allyltrimethylsilane (3.0 equiv) in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ (1.0 equiv, CH_2Cl_2 , RT, 0.5 h), the desired reaction proceeded smoothly to give the α -allyl ketone **4a** in 94% yield. Notably, **4a** was completely racemic, thus suggesting the intermediacy of a well-developed cationic species next to the isoxazole.



Encouraged by the promising opportunities for installing various angular substituents by the S_N1 pathway, we examined the scope of this reaction, which led us to confirm its broad applicability (Table 1). Heteronucleophiles, such as alcohol **5** and thiol **6**, smoothly reacted at 0°C to give ether **4b** and sulfide **4c** in excellent yields (Table 1, entries 1 and 2). Furthermore, installation of angular aryl or (hetero)aryl groups by the Friedel–Crafts pathway proved to be achievable (Table 1, entries 3–9). Phloroglucinol derivative **7** cleanly gave ketone **4d** in 92% yield (Table 1, entry 3), and *N*-methylindole (**8**) reacted at ambient temperature, to give exclusively the 3-substituted product **4e** in 95% yield (Table 1, entry 4). While phenol (**9**) gave a regioisomeric

[*] Dr. H. Takikawa, K. Hikita, Prof. Dr. K. Suzuki
Department of Chemistry
Tokyo Institute of Technology, SORST-JST Agency
2-12-1 O-okayama, Meguro-ku, Tokyo 152-8551 (Japan)
Fax: (+81) 3-5734-2788
E-mail: ksuzuki@chem.titech.ac.jp
Homepage: <http://www.chemistry.titech.ac.jp/~org-synth/>

[**] We are grateful to Dr. Hidehiro Uekusa and Sachiyo Kubo for X-ray analyses. This work was partially supported by the Global COE program (Tokyo Institute of Technology) and a Grant-in-Aid for Scientific Research (JSPS). A JSPS Research Fellowship for Young Scientists to H.T. is also gratefully acknowledged.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200801577>.

Table 1: Reaction of α -ketol **3** with various nucleophiles.^[a]


Entry	Nucleophile	R ^[b]	T [°C]	t [h]	Yield [%]
1 ^[c]			0	10	87 (4b)
2			0	6	93 (4c)
3			0	9	92 (4d)
4			RT	8.5	95 (4e)
5			0 → RT	1	36 ^[d] (4f) 48 ^[d] (4g)
6 ^[e]			0	4	93 (4h)
7 ^[e]			0	8	90 (4i)
8 ^[e]			RT	1.5	90 (4j)
9 ^[e]			RT	16.5	94 (4k)

[a] Unless otherwise indicated, BF₃·OEt₂ (1.0 equiv) and nucleophile (3.0 equiv) in CH₂Cl₂ (0.1 M) at the temperature indicated. [b] Dots represent the position of the connectivity. [c] BF₃·OEt₂ (1.5 equiv). [d] Crude products were acetylated prior to the analysis. [e] Nucleophile (2.0 equiv).

mixture of *ortho*- and *para*-products, **4f** and **4g** (Table 1, entry 5), substituted phenols **10–13** selectively reacted at the *ortho* position to give **4h–4k** (Table 1, entries 6–9). Electron-rich phenols, **10** and **11**, reacted smoothly at 0 °C (Table 1, entries 6 and 7). Although the phenols **12** and **13** with electron-withdrawing groups were less reactive, the reactions smoothly proceeded at room temperature to afford the corresponding products **4j** and **4k** in excellent yields.

At this stage, we embarked on the advanced model studies, en route to isoprenoid hybrid **1** and bis(anthraquinone) **2** (Scheme 3). Associated with the regioselective installation of an angular prenyl group, we examined an indirect approach through cross metathesis.^[10] The angular allyl group in **4a** was converted into a prenyl group to give **4l** in 78 % yield by treatment with an excess amount of 2-methyl-2-butene in the presence of the second-generation Grubbs catalyst.


As a model for the stereoselective installation of an angular anthraquinone unit, which is characteristic to **2**,^[11] we examined the reactions of iodophenol **14**^[12] and naphthol



Scheme 3. Model studies toward **1** and **2**.

15.^[13] Gratifyingly, both substitution reactions smoothly proceeded in an *ortho*-selective manner to afford the corresponding products **4m** and **4n** in high yields.

Scheme 4 is an example for the diastereoselective version of the process, in which α -ketol **16**, having a methyl group at the β position to the reacting center, was used as the model substrate.^[14] Notably, the stereochemical complementarity to the pinacol-based approach was observed in the case of the allylation to ketol **16**. Upon treatment with allylsilane and BF₃·OEt₂, ketol **16** underwent a retentive allylation from the opposite side to the methyl group, to afford ketone **17a** in high yield. By contrast, the inverted installation of an angular allyl group to α -ketol **16** by the pinacol-based approach gave access to the isomeric ketone **19** as the sole product through the nucleophilic addition and the 1,2-shift.



Scheme 4. Complementary approaches for an angular allylation.


The diastereoselectivity was also observed with other nucleophiles (Table 2). Heteronucleophiles exclusively gave the corresponding *trans* products **17b** and **17c** in high yields (Table 2, entries 1 and 2). The reactions of phenol **13** and naphthol **15** occurred regio- and stereoselectively to afford **17d** and **17e** in excellent yields (Table 2, entries 3 and 4).

Table 2: Reaction of α -ketol **16**.^[a]

Entry	R	T [°C]	t [h]	Yield [%] (<i>trans/cis</i>) ^[b]
1 ^[c]		0 → RT	3	80 (17b) (>99:<1)
2		0 → RT	3.5	82 (17c) (>99:<1)
3		RT	18	81 (17d) (15:1)
4		0 → RT	2	93 (17e) (>99:<1)

[a] Unless otherwise indicated, $\text{BF}_3 \cdot \text{OEt}_2$ (1.0 equiv) and nucleophile (1.1 equiv) in CH_2Cl_2 (0.1 M) at the temperature indicated. [b] The stereochemical outcomes were determined by X-ray analysis (see the Supporting Information). [c] $\text{BF}_3 \cdot \text{OEt}_2$ (3.0 equiv).

Scheme 5 illustrates the unmasking of β -diketone **21** through the reductive conversion of ketone **4a**.^[15] The reductive opening of the isoxazole ring in **4a** with molybdenum hexacarbonyl^[16] gave vinylogous amide **20**, which turned out to be unexpectedly resistant at undergoing hydrolysis by usual acid hydrolysis.^[17] However, we were pleased to find that the desired hydrolysis could be executed with nitrosulfuric acid^[18] to give β -diketone **21** in 63% yield.



Scheme 5. Conversion of isoxazole **4a** into 1,3-diketone **21**.

In conclusion, the isoxazole-assisted $\text{S}_{\text{N}}1$ substitution of α -ketol allows facile introduction of angular substituents to polycyclic structures. The efficient ability of the isoxazole unit to stabilize the α -cation has promising potential in organic synthesis, and further studies are underway in our laboratories.

Experimental Section


Typical procedure for the $\text{S}_{\text{N}}1$ substitution of α -ketol **3** with allyltrimethylsilane: $\text{BF}_3 \cdot \text{OEt}_2$ (20 μL , 0.16 mmol) was added to a solution of α -ketol **3** (53 mg, 0.15 mmol) and allyltrimethylsilane (52 mg, 0.46 mmol) in CH_2Cl_2 (1.5 mL) at 0 °C. The reaction mixture was allowed to warm to room temperature. After stirring for 0.5 h, it was cooled to 0 °C and saturated NaHCO_3 was added. The products were extracted with EtOAc ($\times 3$) and the combined organic extracts were washed with brine, dried over Na_2SO_4 , and concentrated under reduced pressure. The residue was purified by PTLC with *n*-hexane/ EtOAc (2:1) to give ketone **4a** (53 mg, 94%) as a white solid.

Received: April 4, 2008

Published online: November 14, 2008

Keywords: Friedel–Crafts reactions · isoxazoles · polyketides · $\text{S}_{\text{N}}1$ reaction


- [1] a) H. Shigemori, K. Komatsu, Y. Mikami, J. Kobayashi, *Tetrahedron* **1999**, 55, 14925–14930; b) K. Komatsu, H. Shigemori, M. Shiro, J. Kobayashi, *Tetrahedron* **2000**, 56, 8841–8844.
- [2] a) H. Kushida, S. Nakajima, T. Koyama, H. Suzuki, K. Ojiri, H. Suda, JP 08143569, **1996**; b) A. M. Socha, D. Garcia, R. Sheffer, D. C. Rowley, *J. Nat. Prod.* **2006**, 69, 1070–1073.
- [3] For recent reviews, see: a) R. Thomas, *ChemBioChem* **2001**, 2, 612–627; b) C. Hertweck, A. Luzhetskyy, Y. Rebets, A. Bechtold, *Nat. Prod. Rep.* **2007**, 24, 162–190; c) U. Rix, C. Fischer, L. L. Remsing, J. Rohr, *Nat. Prod. Rep.* **2002**, 19, 542–580; d) J. Staunton, K. J. Weissman, *Nat. Prod. Rep.* **2001**, 18, 380–416; e) B. J. Rawlings, *Nat. Prod. Rep.* **1999**, 16, 425–484.
- [4] For recent reviews on stereoselective construction of quaternary stereocenters, see: a) *Quaternary Stereocenters: Challenges and Solutions for Organic Synthesis* (Eds.: J. Christoffers, A. Baro), Wiley-VCH, Weinheim, **2005**; b) J. Christoffers, A. Baro, *Adv. Synth. Catal.* **2005**, 347, 1473–1482; c) C. J. Douglas, L. E. Overman, *Proc. Natl. Acad. Sci. USA* **2004**, 101, 5363–5367; d) T. Denissova, L. Barriault, *Tetrahedron* **2003**, 59, 10105–10146; e) J. Christoffers, A. Mann, *Angew. Chem.* **2001**, 113, 4725–4732; *Angew. Chem. Int. Ed.* **2001**, 40, 4591–4597; f) E. J. Corey, A. Guzman-Perez, *Angew. Chem.* **1998**, 110, 402–415; *Angew. Chem. Int. Ed.* **1998**, 37, 388–401; g) K. Fuji, *Chem. Rev.* **1993**, 93, 2037–2066.
- [5] a) J. W. Bode, Y. Hachisu, T. Matsuura, K. Suzuki, *Tetrahedron Lett.* **2003**, 44, 3555–3558; b) T. Matsuura, J. W. Bode, Y. Hachisu, K. Suzuki, *Synlett* **2003**, 1746–1748; c) Y. Hachisu, J. W. Bode, K. Suzuki, *J. Am. Chem. Soc.* **2003**, 125, 8432–8433; d) H. Takikawa, Y. Hachisu, J. W. Bode, K. Suzuki, *Angew. Chem.* **2006**, 118, 3572–3574; *Angew. Chem. Int. Ed.* **2006**, 45, 3492–3494.
- [6] a) L. Claisen, *Ber. Dtsch. Chem. Ges.* **1891**, 24, 3900–3918; b) G. Stagno D'Alcontres, *Gazz. Chim. Ital.* **1950**, 80, 441–455; c) C. Kashima, *Heterocycles* **1979**, 12, 1343–1368; d) P. G. Baraldi, A. Barco, S. Benetti, G. P. Pollini, D. Simoni, *Synthesis* **1987**, 857–869; e) A. I. Kotyatkina, V. N. Zhabinsky, V. A. Khripach, *Russ. Chem. Rev.* **2001**, 70, 641–653.
- [7] This ability is exploited in our previously reported method for the stereocontrolled introduction of angular substituents: chiral, non-racemic α -ketol **3** undergoes stereoselective addition of a nucleophile, and a subsequent regioselective and stereospecific 1,2-shift of the resulting diol **D** affords the angularly substituted ketone **E**; K. Suzuki, H. Takikawa, Y. Hachisu, J. W. Bode, *Angew. Chem.* **2007**, 119, 3316–3318; *Angew. Chem. Int. Ed.* **2007**, 46, 3252–3254.
- [8] For recent studies on $\text{S}_{\text{N}}1$ -type displacements of an hydroxy group, see: a) A. Baba, M. Yasuda, Y. Nishimoto, T. Saito, Y.



Onishi, *Pure Appl. Chem.* **2008**, *80*, 845–854, and references therein; b) G. Kaur, M. Kaushik, S. Trehan, *Tetrahedron Lett.* **1997**, *38*, 2521–2524; c) S. H. Kim, C. Shin, A. N. Pae, H. Y. Koh, M. H. Chang, B. Y. Chung, Y. S. Cho, *Synthesis* **2004**, 1581–1584; d) S. K. De, R. A. Gibbs, *Tetrahedron Lett.* **2005**, *46*, 8345–8350; e) G. V. M. Sharma, K. L. Reddy, P. S. Lakshmi, R. Ravi, A. C. Kunwar, *J. Org. Chem.* **2006**, *71*, 3967–3969.

- [9] a) X. Creary, *Chem. Rev.* **1991**, *91*, 1625–1678; b) M. B. Smith, J. March, *Advanced Organic Chemistry*, 5th ed., Wiley, New York, **2001**, pp. 435–436, and references therein; c) N. Isaacs, *Physical Organic Chemistry*, 2nd ed., Longman Scientific Technical, England, **1995**, pp. 451–454.
- [10] A. K. Chatterjee, D. P. Sanders, R. H. Grubbs, *Org. Lett.* **2002**, *4*, 1939–1942.
- [11] For a benzyne-based approach toward anthraquinone, see: a) T. Matsumoto, T. Hosoya, M. Katsuki, K. Suzuki, *Tetrahedron Lett.* **1991**, *32*, 6735–6736; for a Diels–Alder approach toward anthraquinone, see: b) J. Savard, P. Brassard, *Tetrahedron* **1984**, *40*, 3455–3464; c) A. T. Khan, B. Blessing, R. R. Schmidt, *Synthesis* **1994**, 255–257; d) L. F. Tietze, K. M. Gericke, R. R. Singidi, *Angew. Chem.* **2006**, *118*, 7146–7150; *Angew. Chem. Int. Ed.* **2006**, *45*, 6990–6993.
- [12] a) T. Matsumoto, H. Yamaguchi, K. Suzuki, *Tetrahedron* **1997**, *53*, 16533–16544; b) T. Matsumoto, T. Sohma, H. Yamaguchi, K. Suzuki, *Chem. Lett.* **1995**, 677–678.
- [13] a) G. Matsuo, Y. Miki, M. Nakata, S. Matsumura, K. Toshima, *J. Org. Chem.* **1999**, *64*, 7101–7106; b) M. A. Brimble, T. J. Brenstrum, *J. Chem. Soc. Perkin Trans. 1* **2001**, 1624–1634.

- [14] For the preparation of chiral, nonracemic ketol **16**, see: Y. Koyama, R. Yamaguchi, K. Suzuki, *Angew. Chem.* **2008**, *120*, 1100–1103; *Angew. Chem. Int. Ed.* **2008**, *47*, 1084–1087.
- [15] For the oxidative conversion of isoxazoles into α -hydroxy- β -diketones, see: H. Takikawa, A. Takada, K. Hikita, K. Suzuki, *Angew. Chem.* **2008**, *120*, 7556–7559; *Angew. Chem. Int. Ed.* **2008**, *47*, 7446–7449.
- [16] M. Nitta, T. Kobayashi, *J. Chem. Soc. Perkin Trans. 1* **1985**, 1401–1406.
- [17] Upon exposure to H_2SO_4 (3M in THF, 40°C, 6 h), vinylogous amide **20** underwent slow decomposition to give acid **22** (78%) presumably by a retro-Claisen reaction of intermediary iminium **F** as shown in the scheme below. We confirmed that β -diketone **21** was most resistant to decomposition under the same reaction conditions (3M H_2SO_4 in THF, 40°C), further supporting our mechanistic postulate.



- [18] For hydrolysis of amides with nitrosonium salts, see: G. A. Olah, J. A. Olah, *J. Org. Chem.* **1965**, *30*, 2386–2387.