

Table 1: Catalytic formation of dienes from propargylic esters and TMS diazomethane.^[a]

Substrate 1	Reaction time	Diene 4	Yield [%] ^[b]	Stereoselectivity (<i>E</i> : <i>Z</i>)
	5 min		99	–
1a		4a		
	5 min		99	–
1b		4b		
	4 h		68 ^[c]	–
1c		4c		
	2 h		89	65:35
1d		4d + 4d'		
	2 h		99	65:35
1e		4e + 4e'		

[a] Reaction conditions: propargylic carboxylate (1.2 mmol, 1 equiv), styrene (5 equiv), trimethylsilyldiazomethane 2 M in diethyl ether (1.1 equiv), [RuCl(cod)Cp*] (5 mol %), 60 °C in 1 mL of dioxane.
 [b] Yield of compound isolated by chromatography on silica gel. [c] Conversion of 70%.

easily generates a ruthenium carbene on reaction with [RuCl(cod)Cp*].^[13,17]

The reaction of propargylic acetate **1a** (1.2 mmol) with N₂CHTMS (1.1 equiv 2 M in Et₂O, TMS = trimethylsilyl), styrene (5 equivalents) and [RuCl(cod)Cp*] (5 mol %) in dioxane for 5 min at 60 °C led to the formation of the conjugated diene **4a**, which was isolated in 99 % yield (Table 1).

The propargylic esters **1b–e**, with two substituents at the propargylic position, reacted under the same reaction conditions and gave the products **4b–e** in times of 5 min–2 h, in good yields (89–99 %) that were not influenced by the substituent bulkiness (Table 1).

As for the vinylcarbene dimerization [Eq. (3)], the presence of styrene was beneficial: without 5 equivalents of styrene, dienes **4d** and **4d'** were obtained in only 50 % yield. Coordination of styrene to the ruthenium intermediate may temporarily protect the coordinatively unsaturated ruthenium carbene species.




With disubstituted propargylic carboxylates **1a–e**, the formation of the TMS-substituted double bond in dienes **4** was highly stereoselective (100 % *Z*) Table 1. When the two substituents on **1** were different (**1d–e**), a mixture of two isomers in 65:35 ratio was obtained, arising from the *Z* and *E* configurations of the tetrasubstituted double bond, as shown by NOESY experiments.

The monosubstituted propargylic carboxylate **1f** led to a mixture of four isomers of type **4** with a 99 % total yield

[Eq. (4)]. The two major isomers, **4f'** with *E* configuration of the acetate-substituted double bond (**4f**₁ (*EZ*)/**4f**₂ (*EE*): 80/20), and the two minor ones, **4f** (**4f**₁ (*ZZ*)/**4f**₂ (*ZE*): 80/20), were obtained in the ratio 85:15. The major products have a silylalkenyl bond with *E* configuration, in contrast to disubstituted derivatives **4a–4e**.


The desilylation of compounds **4** was not easy to perform. Thus, the reaction was carried out with in situ-generated diazomethane, by using trimethylsilyldiazomethane in methanol as solvent with **1a** in the presence of styrene (5 equivalents). After only 15 min of reaction at 60 °C, the corresponding desilylated diene **5** was isolated with a good yield (91 %) [Eq. (5)].

Other diazoalkanes proved less reactive than N₂CHTMS for the formation of dienes. The reaction of **1a** with phenyldiazomethane in the presence of styrene (5 equivalents) at 60 °C after 16 h led to the corresponding conjugated diene **6** in moderate yield (51 %). However,



the phenyl-substituted double bond in **6** has the *E* configuration [Eq. (6)].

At this stage, the reaction mechanism has not been elucidated as the initial adduct of [RuCl(cod)Cp*] with **1a** or N₂CHTMS could not be detected. Two possible approaches can be considered (Scheme 1). The first potential mechanism



Scheme 1. Possible catalytic cycles for the synthesis of dienes from propargylic esters and diazoalkanes. Y = TMS, Ph, or H.

involves the initial formation of the ruthenium vinylcarbene intermediate **A**, after the Rautenstrauch rearrangement of the propargylic carboxylate^[7] in the presence of the $[\text{RuCl}(\text{cod})\text{Cp}^*]$ catalyst. Addition of the diazoalkane then leads to the formation of a ruthenium bis(carbene)–metal intermediate **B**, leading to the formation of the C=C bond and diene **4**, or to the direct reaction of intermediate **A** with diazoalkane to give **4**.^[18] The reaction of intermediate **A** with **1** could lead to formation of **2**.

Another possible mechanism involves the initial interaction of the catalyst with the diazoalkane, leading to carbene **C**, which undergoes [2+2] cycloaddition with the triple bond of **1**, leading to intermediate **D**, as in enyne metathesis,^[12,13] followed by formation of the vinylcarbene–ruthenium intermediate **E**.^[19] The 1,2-shift of the acetate to the carbene carbon of **E** would form **4**.^[3d,5g] The preferred *Z* configuration of the silylalkenyl bond may arise from the interaction of the silyl group with the chloride (upper cycle),^[13] or by ring-opening of intermediate **D**, similar to that of 3-silylcyclobutenes (lower cycle).^[20]

In summary, the $[\text{Ru}(\text{Cl})\text{Cp}^*]$ -catalyzed formation of conjugated dienes proceeded with good yields and selectivities by cross-coupling of two carbene fragments arising from two different sources, a vinylcarbene fragment resulting from a propargylic ester rearrangement and a diazoalkane carbene. This method provides a novel approach to the synthesis of dienes, precursors of α,β -unsaturated silylated ketones, and has potential, owing to the large number of in situ generated metal–vinylcarbenes, to create a variety of functionalized dienes. This C=C bond forming reaction is currently under study.

Experimental Section

Typical procedure for ruthenium-catalyzed cross-coupling of carbenes: $[\text{RuCl}(\text{cod})\text{Cp}^*]$ (5 mol %) was added as a solid to a solution of propargyl acetate **1a–1e** (1.2 mmol), styrene (5 equiv) and trimethylsilyldiazomethane (1.1 equiv, 2 M solution in diethyl ether) in dioxane (1 mL) at 60 °C under an inert atmosphere. The reaction mixture was stirred for 5 min–2 h. The solvent was removed under vacuum, and the products were isolated by column chromatography on silica gel, eluted with mixed solvent (pentane/diethyl ether).

Full experimental details and characterization data are given in the Supporting Information.

Received: October 14, 2008

Revised: November 20, 2008

Published online: January 19, 2009

Keywords: carbenes · catalysis · cross-coupling · dienes · ruthenium

- [1] N. Chatani, K. Kataoka, S. Murai, *J. Am. Chem. Soc.* **1998**, *120*, 9104.
- [2] B. P. Peppers, S. T. Diver, *J. Am. Chem. Soc.* **2004**, *126*, 9524.
- [3] a) K. Miki, K. Ohe, S. Uemura, *J. Org. Chem.* **2003**, *68*, 8505; b) K. Miki, K. Ohe, S. Uemura, *Tetrahedron Lett.* **2003**, *44*, 2019; c) K. Miki, M. Fujita, S. Uemura, K. Ohe, *Org. Lett.* **2006**, *8*, 1741; d) K. Ohe, M. Fujita, H. Matsumoto, Y. Tai, K. Miki, *J. Am. Chem. Soc.* **2006**, *128*, 9270; e) Y. Nakanishi, K. Miki, K. Ohe, *Tetrahedron* **2007**, *63*, 12138.
- [4] a) A. Fürstner, F. Stelzer, H. Szillat, *J. Am. Chem. Soc.* **2001**, *123*, 11863; b) M. Méndez, M. P. Muñoz, C. Nevado, D. J. Cárdenas, A. M. Echavarren, *J. Am. Chem. Soc.* **2001**, *123*, 10511; c) E. Mainetti, V. Mouriès, L. Fensterbank, M. Malacria, J. Marco-Contelles, *Angew. Chem.* **2002**, *114*, 2236; *Angew. Chem. Int. Ed.* **2002**, *41*, 2132; d) V. Mamane, T. Gress, H. Krause, A. Fürstner, *J. Am. Chem. Soc.* **2004**, *126*, 8654; e) E. J. Cho, M. Kim, D. Lee, *Org. Lett.* **2006**, *8*, 5413; f) B. A. Bhanu Prasad, F. K. Yoshimoto, R. Sarpong, *J. Am. Chem. Soc.* **2005**, *127*, 12468.
- [5] a) For a review see: E. Jiménez-Núñez, A. M. Echavarren, *Chem. Rev.* **2008**, *108*, 3326; b) M. R. Luzung, J. P. Markham, F. D. Toste, *J. Am. Chem. Soc.* **2004**, *126*, 10858; c) C. Nieto-Oberhuber, M. P. Muñoz, E. Buñuel, C. Nevado, D. J. Cárdenas, A. M. Echavarren, *Angew. Chem.* **2004**, *116*, 2456; *Angew. Chem. Int. Ed.* **2004**, *43*, 2402; d) D. J. Gorin, N. R. Davis, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 11260; e) C. Nieto-Oberhuber, S. Lopez, M. P. Muñoz, D. J. Cárdenas, E. Buñuel, C. Nevado, A. M. Echavarren, *Angew. Chem.* **2005**, *117*, 6302; *Angew. Chem. Int. Ed.* **2005**, *44*, 6146; f) P. Y. Toullec, E. Genin, L. Leseurre, J. P. Genêt, V. Michelet, *Angew. Chem.* **2006**, *118*, 7587; *Angew. Chem. Int. Ed.* **2006**, *45*, 7427; g) E. J. Cho, M. Kim, D. Lee, *Eur. J. Org. Chem.* **2006**, 3074.
- [6] a) C. Bruneau, *Angew. Chem.* **2005**, *117*, 2380; *Angew. Chem. Int. Ed.* **2005**, *44*, 2328; b) J. Marco-Contelles, E. Soriano, *Chem. Eur. J.* **2007**, *13*, 1350; c) V. Michelet, P. Y. Toullec, J. P. Genêt, *Angew. Chem.* **2008**, *120*, 4338; *Angew. Chem. Int. Ed.* **2008**, *47*, 4268.
- [7] V. Rautenstrauch, *J. Org. Chem.* **1984**, *49*, 950.
- [8] a) X. Shi, D. J. Gorin, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 5802; b) N. Marion, P. de Frémont, G. Lemièrre, E. D. Stevens, L. Fensterbank, M. Malacria, S. P. Nolan, *Chem. Commun.* **2006**, 2048; c) M. J. Johansson, D. J. Gorin, S. T. Staben, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 18002.
- [9] A. Fürstner, P. Hannen, *Chem. Commun.* **2004**, 2546.
- [10] L. Peng, X. Zhang, S. Zhang, J. Wang, *J. Org. Chem.* **2007**, *72*, 1192.

- [11] a) M. A. Sierra, J. C. del Amo, M. J. Mancheño, M. Gómez-Gallego, *J. Am. Chem. Soc.* **2001**, *123*, 851; b) F. Robin-Le Guen, P. Le Poul, B. Caro, N. Faux, N. Le Poul, S. J. Green, *Tetrahedron Lett.* **2002**, *43*, 3967.
- [12] a) A. Kinoshita, M. Mori, *Synlett* **1994**, 1020; b) S.-H. Kim, N. B. Bowden, R. H. Grubbs, *J. Am. Chem. Soc.* **1994**, *116*, 10801; c) S.-H. Kim, W. J. Zuercher, N. B. Bowden, R. H. Grubbs, *J. Org. Chem.* **1996**, *61*, 1073.
- [13] a) F. Monnier, C. Vovard-Le Bray, D. Castillo, V. Aubert, S. Dérien, P. H. Dixneuf, L. Toupet, A. Ienco, C. Mealli, *J. Am. Chem. Soc.* **2007**, *129*, 6037; b) F. Monnier, D. Castillo, S. Dérien, L. Toupet, P. H. Dixneuf, *Angew. Chem.* **2003**, *115*, 5632; *Angew. Chem. Int. Ed.* **2003**, *42*, 5474.
- [14] a) J. Le Paih, F. Monnier, S. Dérien, P. H. Dixneuf, E. Clot, O. Eisenstein, *J. Am. Chem. Soc.* **2003**, *125*, 11964; b) C. Gemel, A. La Pensée K. Mauthner, K. Mereiter, R. Schmid, K. Kirchner, *Monatsh. Chem.* **1997**, *128*, 1189.
- [15] Higher temperature (60°C) favored the classical dimerization of terminal alkyne **1a** before its rearrangement by acetate transfer, into enyne **3a**. This process is known to proceed by insertion of the vinylidene intermediate into a Ru–alkynyl bond.^[16]
- [16] a) C. Bianchini, P. Frediani, D. Masi, M. Peruzzini, F. Zanobini, *Organometallics* **1994**, *13*, 4616; b) C. S. Yi, N. Liu, *Organometallics* **1996**, *15*, 3968.
- [17] a) A. W. Stumpf, E. Saive, A. Demonceau, A. F. Noels, *J. Chem. Soc. Chem. Commun.* **1995**, 1127; b) A. Demonceau, A. W. Stumpf, E. Saive, A. F. Noels, *Macromolecules* **1997**, *30*, 3127.
- [18] This process is proposed for the ruthenium-catalyzed dimerization of diazoalkanes into symmetrical alkenes by addition of an electrophilic metal carbene carbon to the nucleophilic diazoalkane carbon. When Grubbs catalyst is used it does not lead to the cross coupling of carbenes. See: a) D. M. Hodgson, D. Angrish, *Chem. Commun.* **2005**, 4902; b) H. M. Lee, C. Bianchini, G. Jia, P. Barbaro, *Organometallics* **1999**, *18*, 1961.
- [19] These initial steps have been proposed for the addition of diazoalkane to enynes^[13] or to alkynes: J. Le Paih, S. Dérien, I. Özdemir, P. H. Dixneuf, *J. Am. Chem. Soc.* **2000**, *122*, 7400.
- [20] M. Hasegawa, M. Murakami, *J. Org. Chem.* **2007**, *72*, 3764.