

A Journal of the Gesellschaft Deutscher Chemiker


Angewandte Chemie

International Edition



www.angewandte.org

2009–48/23



Mass spectrometry of biomolecules ...

... has developed into a key technology in bioanalytics. As described by C. R. Gebhardt, M. Dürr et al. in their Communication on page 4162 ff., surface collisions of neutral molecular clusters can be used for soft formation of the required gas-phase biomolecular ions. Upon impact on a surface, the clusters provide the energy for desorption of the adsorbed biomolecules and serve as a transient matrix. Subsequent evaporative cooling prevents fragmentation and finally leads to an isolated biomolecular ion in the gas phase.

WILEY-VCH

Inside Cover

**Christoph. R. Gebhardt,* Anna Tomsic, Hartmut Schröder,
Michael Dürr,* and Karl L. Kompa**

Mass spectrometry of biomolecules has developed into a key technology in bioanalytics. As described by C. R. Gebhardt, M. Dürr et al. in their Communication on page 4162 ff., surface collisions of neutral molecular clusters can be used for soft formation of the required gas-phase biomolecular ions. Upon impact on a surface, the clusters provide the energy for desorption of the adsorbed biomolecules and serve as a transient matrix. Subsequent evaporative cooling prevents fragmentation and finally leads to an isolated biomolecular ion in the gas phase.

