

Activation of Hydrogen by Palladium(0): Formation of the Mononuclear Dihydride Complex *trans*-[Pd(H)₂(IPr)(PCy₃)]**

Serena Fantasia, Jonathan D. Egbert, Václav Jurčík, Catherine S. J. Cazin, Heiko Jacobsen, Luigi Cavallo, D. Michael Heinekey,* and Steven P. Nolan*

Metal-catalyzed hydrogenation reactions have been the subject of intensive investigation in organometallic chemistry and catalysis.^[1] Late transition metals employed to catalyze this transformation have exhibited impressive performances. In its homogeneous incarnation, transition-metal catalyzed hydrogenation is best mediated by rhodium (notably Wilkinson's catalyst)^[2] or iridium^[3] complexes. Other metals have fared better in heterogeneous systems, as is the case for palladium whose use in its supported form, Pd/C, is ubiquitous.^[1b]

Palladium-catalyzed homogeneous hydrogenation mainly makes use of Pd^{II} complexes, wherein a heterolytic activation of the dihydrogen molecule is believed to occur.^[4] Less attention has focused on Pd⁰ systems,^[5] in view of the reported and rationalized low reactivity of Pd⁰ complexes towards H₂.^[6] However, in the past decade, efficient hydrogenation systems based on Pd⁰ have been reported.^[7] Mechanistic investiga-

tions suggest that the catalytic manifold involves coordination of the substrate, followed by activation of H₂ to give Pd^{II} species.^[8] In this context, the possible involvement of monomeric Pd^{II} dihydride intermediates, formed by homolytic activation of H₂, is generally ruled out since such species have not, to date, been detected.^[9] Remarkably, oxidative addition of H₂ to afford dihydride complexes is a well-documented reaction for late transition metals.^[10] One of the few exceptions to this generally acknowledged fact is palladium. Indeed, to our knowledge, only polynuclear complexes are known for Pd compounds bearing more than one hydride ligand.^[11]

As part of our investigations on the use of *N*-heterocyclic carbenes (NHCs) as stabilizing ligands,^[12] a synthetic strategy leading to 14-electron Pd⁰ complexes was developed.^[13] One of these complexes, [Pd(SIPr)(PCy₃)] (SIPr = *N,N'*-bis(2,6-diisopropylphenyl)imidazolin-2-ylidene), was successfully used to promote hydrogenation reactions under very mild reaction conditions.^[14] During these catalytic studies, the stability of [Pd(SIPr)(PCy₃)] under hydrogenation conditions was investigated, and NMR spectroscopic studies indicated the presence of a hydride complex which may have resulted from an oxidative addition of H₂ on [Pd(SIPr)(PCy₃)]. However, only partial conversion was detected and the isolation and full characterization of this species was not achieved. Herein, we report, through judicious ancillary ligand selection, the isolation and full characterization of the first monomeric palladium dihydride complex.

As mentioned above, [Pd(SIPr)(PCy₃)] only reacts sparingly with H₂. A better candidate for such experiment might be one with a smaller steric bulk. Preliminary experiments were performed with [Pd(IPr)(PPh₃)] (IPr = *N,N'*-bis-(2,6-diisopropylphenyl)imidazol-2-ylidene) which showed that, after 24 h, under 1 atm of H₂, a hydride species was formed (¹H NMR: weak doublet at $\delta = -4.21$ ppm with $J = 17$ Hz) but product formation was incomplete and subsequent decomposition to palladium black took place. We thus opted for [Pd(IPr)(PCy₃)], as it contains a more electron-rich phosphane, which might favor product formation, and because IPr is slightly less hindered than its SIPr congener. To our delight, exposure of [Pd(IPr)(PCy₃)] (**1**) to H₂ (1 atm) led to almost quantitative conversion of **1** into a novel hydride species **2**.

The ³¹P{¹H} NMR spectrum of **2** exhibited a single resonance at $\delta_P = 68.2$ ppm ($\delta_P(\mathbf{1}) = 49.3$ ppm). In the ¹H NMR spectrum, the imidazole and the isopropyl resonances were shifted downfield by approximately 0.24 ppm with respect to **1**.^[15] A doublet appeared at $\delta_H = -4.81$ ppm ($^2J_{H,P} = 16$ Hz), integrating for two protons, which suggested the

[*] J. D. Egbert, Prof. Dr. D. M. Heinekey
Department of Chemistry, University of Washington
Box 351700, Seattle, WA 98195-1700 (USA)
Fax: (+1) 206-616-4209
E-mail: heinekey@chem.washington.edu

Dr. S. Fantasia,^[4] Dr. V. Jurčík, Dr. C. S. J. Cazin, Prof. Dr. S. P. Nolan
Institute of Chemical Research of Catalonia (ICIQ)
Av. Països Catalans 16, 43007 Tarragona (Spain)

Dr. V. Jurčík, Dr. C. S. J. Cazin, Prof. Dr. S. P. Nolan
Present address: University of St Andrews, School of Chemistry
St Andrews, KY16 9ST (UK)
E-mail: sn17@st-andrews.ac.uk

Dr. H. Jacobsen
KemKom, 1215 Ursulines Ave, New Orleans, LA 70116 (USA)

Prof. Dr. L. Cavallo
Dipartimento di Chimica, Università di Salerno
Via ponte don Melillo, 84084 Fisciano (SA) (Italy)

[†] Present address: Laboratorium für Anorganische Chemie
ETH Zürich, Wolfgang-Pauli-Strasse 110, 8093 Zürich (Switzerland)

[**] S.F. and S.P.N. gratefully thank the MEC (Spain) and the ICIQ Foundation for financial support. S.P.N. acknowledges the ERC for support in the form of an Advanced Researcher Grant (FUNCAT). J.D.E. and D.M.H. acknowledge the US NSF. We thank Dr. W. Kaminsky, Dr. J. Benet-Buchholz, and E. Escudero Adán for X-ray structures. L.C. thanks the ENEA (www.enea.it) and the HPC team for support and for using ENEA-GRID and the HPC facilities CRESCO (www.cresco.enea.it) Portici (Naples), Italy.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200900463>. CCDC 707593 (**2**) and 707594 (**4**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif

presence of two hydride ligands *cis* to the phosphorus atom.^[16] However, to confirm this hypothesis, and to rule out other possibilities, such as a dynamic *cis* dihydride or a dihydrogen complex, further NMR spectroscopic experiments were conducted.

The relaxation time T_1 of the hydride resonance in complex **2** was measured as a function of temperature using a standard inversion recovering sequence. The minimum T_1 of 420 ms occurred at 270 K (500 MHz). Dihydrides and dihydrogen complexes can be distinguished by their minimum T_1 value, by following the method of Halpern and co-workers.^[17] This method is not straightforward when multiple nuclei contribute to the overall relaxation rate, defined as R ($R=1/T_1$). For complex **2**, the contributions of the phosphorus atom and the other H atoms present in the molecule have to be considered. To determine these contributions, we prepared the monohydride complex $[\text{Pd}(\text{H})(\text{Cl})(\text{IPr})(\text{PCy}_3)]$ (**3**). Complex **3** was formed quantitatively as a white solid by addition of an equimolar amount of HCl (4 M in dioxane) to a THF solution of **1**.

Halogeno-hydride complexes of Pd^{II} have been extensively studied and well characterized.^[18] However, palladium hydride complexes bearing NHC ligands remain rare,^[19] possibly as a result of the instability of such complexes towards reductive elimination of the imidazolium salt.^[20] Complex **3** is indefinitely stable both in the solid state and in solution (even upon heating at 80 °C) under Ar atmosphere, giving no sign of reductive elimination of the IPr hydrochloride salt. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **3** displayed a singlet at $\delta_{\text{P}} = 48.2$ ppm, while in the ^1H NMR spectrum the hydride resonance appears as a doublet at $\delta_{\text{H}} = -14.89$ ppm ($^2J_{\text{H,P}} = 2$ Hz). In the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum, the carbenic carbon signal appeared as a doublet centered at $\delta_{\text{C}} = 190.5$ ppm ($^2J_{\text{C,P}} = 137$ Hz).

The minimum relaxation time constant for the hydride resonance in **3**, $T_1 = 590$ ms, occurred at 270 K (500 MHz). Comparison of the relaxation rate in **2** and **3** ($R_2 - R_3$) allowed for determination of the HH relaxation-rate contribution, R_{HH} , in **2**, which is 0.69 s^{-1} .^[21] This value excluded the possibility of a $\eta^2\text{-H}_2$ complex,^[22] and suggested formulation of **2** as the *trans* dihydride complex. Its formation would then proceed according to Equation (1).



Reaction of **1** with HD gas [Eq. (2)], afforded complex $[\text{D}]\text{-2}$, for which the ^1H NMR hydride signal appeared as a multiplet centered at $\delta = -4.72$ ppm ($^2J_{\text{H,D}} = 7$ Hz, $^2J_{\text{H,P}} = 16$ Hz; Figure 1). The value for the H–D coupling constant is consistent with the formation of the *trans* isomer.^[23] Interestingly, in addition to the resonance of complex $[\text{D}]\text{-2}$, the hydride resonance of complex **2** was also present in the ^1H NMR spectrum. Comparison of the integration value of the two hydride signals with the integration of the isopropyl CH resonance of the IPr ligand (3 vs. 12), strongly suggested the presence of the bis-deuterated complex $[\text{D}_2]\text{-2}$. The presence of all three isotopomers was confirmed by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy (Figure 1), in which three resonances, integrated for their statistical distribution, appeared: a singlet at $\delta_{\text{P}} = 68.22$ ppm (**2**), a 1:1:1 triplet at $\delta_{\text{P}} = 68.28$ ppm ($[\text{D}]\text{-2}$) and a five-line pattern at $\delta_{\text{P}} = 68.33$ ppm ($[\text{D}_2]\text{-2}$). The mechanism of this isotope exchange reaction is under investigation.




Figure 1. a) ^1H and b) $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of the three isotopomers **2**, $[\text{D}]\text{-2}$, and $[\text{D}_2]\text{-2}$, formed from the reaction of **1** with HD gas.

Density functional (DFT) calculations were carried out to assess the relative stabilities of the *cis* and the *trans* isomers. We found the *trans* isomer to be lower in energy by $6.5 \text{ kcal mol}^{-1}$ (in THF) compared to the *cis* isomer. We then examined the thermodynamics of the hydrogenation reaction (1), and found it to be exothermic by $3.1 \text{ kcal mol}^{-1}$.^[15]

Furthermore, the structure of the *trans*-dihydride complex **2** was unambiguously confirmed by single-crystal X-ray diffraction (Figure 2).







Figure 2. ORTEP representation of **2** $[\text{Pd}(\text{H})_2(\text{IPr})(\text{PCy}_3)]$. Thermal ellipsoids are set at 50% probability. H atoms, with the exception of H1a and H1b, are omitted for clarity.

The metal center in **2** has an approximately square planar coordination geometry, slightly distorted by steric interactions between the IPr and PCy₃ ligands. The two hydrogen atoms on Pd were located and refined with isotropic thermal parameters. The location of the hydrogen atoms coincides with free voids surrounding the Pd center.

The formation of **2** proved to be extremely sensitive to oxygen and water. When extreme precautions were not taken to avoid the presence of air and H₂O, the amount of dihydride complex **2** present in solution was substantially lowered. Indeed, in the presence of O₂, the peroxo complex [Pd(η²-O₂)(IPr)(PCy₃)] **4** was formed.^[15] As expected, the ¹³C{¹H} resonance for the Pd carbene carbon atom in complex **4** has a coupling constant, ²J_{C,P} = 16 Hz, that was considerably lower than the corresponding coupling constants for complexes **1**, **2**, and **3** (²J_{C,P} = 90, 138, and 137 Hz, respectively). This value reflects the *cis* configuration of the IPr and the PCy₃ ligands in **4**. In this reaction, it remains unclear whether coordination of O₂ takes place on the starting material **1** or on the dihydride product **2**, as preliminary studies suggested that the dihydride species is more reactive towards dioxygen than **1**.

The Pd–C_{carbene} and Pd–P bond lengths in **4** are substantially longer than those in **1** and **2** (Table 1). As expected, the

Table 1: Selected bond lengths [Å] and angles [°] for complexes **1**, **2**, and **4**.

			
	1 [13]	2	4
Pd–C	2.0292(9)	2.009(2)	2.056(3)
Pd–P	2.2212(3)	2.2385(6)	2.3303(7)
C–Pd–P	170.88(2)	173.08(7)	105.24(7)

C–Pd–P angle found for **4** was much smaller than the those for **1** and **2**, confirming in the first case a *cis* arrangement of the NHC and PCy₃, and a *trans* arrangement for the dihydride complex.

To examine whether there was truly something unique with the system bearing mixed ligands, it seemed of interest to investigate the reactivity of homoleptic Pd⁰ complexes towards H₂. Reactions of [Pd(PCy₃)₂] and [Pd(IPr)₂] with H₂ in [D₆]benzene were monitored by ¹H NMR spectroscopy. In the case of [Pd(PCy₃)₂], no hydride signal was detected even after several weeks or when experiments were carried out at low temperature. For [Pd(IPr)₂], a very weak signal at δ_H = –4.80 ppm was detected, but the reaction did not proceed to any significant extent. These observations, as well as those for [Pd(SIPr)(PCy₃)] and [Pd(IPr)(PPh₃)], emphasize that ligand tuning is crucial in the success of the H–H bond activation reaction. In the case of **1**, the stereoelectronic properties of both the NHC^[24] and phosphane^[25] ligands lead to an optimal situation for the detection and isolation of palladium species resulting from H₂ activation.

In summary, we have reported the activation of H₂ by a mixed NHC-phosphane Pd⁰ complex, leading to the formation of an unprecedented monomeric dihydride palladium species. The isolation of a dihydride complex demonstrated the feasibility of direct activation of H₂ by a Pd⁰ complex. These results may pave the way for mechanistic studies on catalytic hydrogenation and for the development of new catalytic systems. Further studies aimed at understanding the stability and reactivity of this species are ongoing in our laboratories.

Experimental Section

All manipulations were carried out under an Argon atmosphere, using standard dry box, Schlenk and vacuum line techniques. [Pd(IPr)(L)] (L = PPh₃, PCy₃, IPr),^[13] [Pd(SIPr)(PCy₃)],^[13] and [Pd(PCy₃)₂]^[26] were synthesized according to literature procedures. Anhydrous solvents were purchased from Aldrich and used as received unless otherwise specified. NMR spectra were recorded on Bruker instruments (400 MHz and 500 MHz) in [D₆]benzene. Variable-temperature NMR spectroscopy was carried out in [D₈]toluene. Mass spectra were recorded on a Waters LCT Premier instrument. Elemental analyses were performed by Universidad Complutense de Madrid and Columbia Analytical Services, Inc., Tucson, AZ.

2 (*trans*-[Pd(H)₂(IPr)(PCy₃)]): A J. Young NMR tube was charged with [Pd(IPr)(PCy₃)] (**1**, 5.0 mg, 6.4 μmol) and then filled with C₆D₆ (dried over Na/K alloy) by vacuum transfer. After freezing the solution, the atmosphere over the solution was replaced with H₂ gas (1 atm). After 24 h, the solution was analyzed by NMR spectroscopy. ¹H NMR (500 MHz): δ = 7.23 (t, *J* = 10 Hz, 2H, CH-*p*), 7.18 (d, *J* = 10 Hz, 4H, CH-*m*), 6.73 (s, 2H, CH imid), 3.25 (sept, *J* = 5 Hz, 4H, CH(CH₃)₂), 2.0–1.0 (m, CH₂ + CH Cy), 1.59 (d, *J* = 5 Hz, CH(CH₃)₂), 1.19 (d, *J* = 5 Hz, CH(CH₃)₂), –4.81 ppm (d, *J*_{P,H} = 16 Hz, 2H, Pd–H); ¹³C{¹H} NMR (125.8 MHz): δ = 200.5 (d, *J*_{C,P} = 138 Hz, C carbene), 146.8 (C arom), 136.9 (C arom), 128.6 (CH arom), 122.8 (CH arom), 127.7 ppm (d, *J*_{C,P} = 4 Hz, CH imid); ³¹P{¹H} NMR (202.5 MHz): δ = 68.1 ppm; elemental analysis calcd for C₄₅H₇₁N₃PPd: C 69.52, H 9.20, N 3.60; found: C 70.42, H 9.06, N 3.47.

3 (*trans*-[Pd(H)(Cl)(IPr)(PCy₃)]): Compound **1** (42.0 mg, 54 μmol) was dissolved in THF (1 mL) and treated with HCl (20 μL of a 4 M solution in dioxane). After stirring at room temperature for 4 h, evaporation of the solvent afforded the product (44 mg, quantitative yield). ¹H NMR (400 MHz): δ = 7.28 (t, *J* = 8 Hz, 2H, CH-*p*), 7.21 (d, *J* = 8 Hz, 4H, CH-*m*), 6.70 (s, 2H, CH imid), 3.33 (sept, *J* = 8 Hz, 4H, CH(CH₃)₂), 1.87 (br q, *J* = 12 Hz, 3H, CH Cy), 1.73 (br d, *J* = 12 Hz, 6H, CH₂), 1.65 (br m, 9H, CH₂), 1.57 (d, *J* = 8 Hz, 12H, CH(CH₃)₂), 1.29 (br q, *J* = 12 Hz, 6H, CH₂), 1.16 (br m, 9H, CH₂), 1.12 (d, *J* = 8 Hz, 12H, CH₂), –14.99 ppm (d, *J* = 2 Hz, 1H, Pd–H); ¹³C{¹H} NMR (125.8 MHz): δ = 190.5 (d, *J*_{C,P} = 137 Hz, C carbene), 146.8 (C arom), 136.7 (C arom), 129.5 (CH arom), 123.5 (CH arom), 123.2 (d, *J*_{C,P} = 4 Hz, CH imid), 34.1 (d, *J*_{C,P} = 20 Hz, CH Cy), 30.2 (d, *J*_{C,P} = 2 Hz, CH₂), 28.6 (CH(CH₃)₂), 27.5 (d, *J*_{C,P} = 11 Hz, CH₂), 26.5 (CH₂), 25.9 (CH(CH₃)₂), 22.5 ppm (CH(CH₃)₂); ³¹P{¹H} NMR (162 MHz): δ = 48.2 ppm; ESI MS: *m/z* calcd for C₄₅H₇₀ClN₂PPd: 810.4; found: 810.4; elemental analysis calcd for C₄₉H₇₈ClN₂O₂PPd [*M*+dioxane]: C 65.39, H 8.74, N 3.11; found: C 65.32, H 8.43, N 3.41.

4 [Pd(η²-O₂)(IPr)(PCy₃)]: Compound **1** (100.0 mg, 127 μmol) was dissolved in benzene (10 mL). Air was bubbled through the solution for 15 min after which time evaporation of the solvent afforded **4** (102.0 mg; quantitative yield). ¹H NMR (400 MHz): δ = 7.24 (t, *J* = 3 Hz, 2H, CH-*p*), 7.17 (under C₆D₆ signal, 4H, CH-*m*), 6.65 (s, 2H, CH imid), 3.22 (br m, 4H, CH(CH₃)₂), 1.72–1.59 (m, 27H), 1.25 (br m, 12H, Cy), 1.10 (br m, 6H, Cy), 1.05 ppm (d, *J* = 8 Hz, 12H, CH(CH₃)₂); ¹³C{¹H} NMR (100.6 MHz): δ = 192.0 (d, *J*_{C,P} = 16 Hz,

C carbene), 146.3 (C arom), 136.9 (C arom), 129.9 (CH arom), 125.0 (CH arom), 124.1 (s, CH imid), 33.5 (d, $J_{\text{CP}} = 15$ Hz, CH Cy), 30.0 (d, $J_{\text{CP}} = 2$ Hz, CH₂), 28.8 (CH(CH₃)₂), 27.4 (d, $J_{\text{CP}} = 11$ Hz, CH₂), 26.6 (CH₂), 26.2 (CH(CH₃)₂), 23.3 ppm (CH(CH₃)₂); ³¹P{¹H} NMR (162 MHz): $\delta = 53.7$ ppm; MS (MALDI-TOF): m/z calcd for C₄₅H₆₉N₂PPd [M-(O₂)]: 774.4; found 774.4; elemental analysis calcd for C₄₅H₇₃N₂O₄PPd [M+2H₂O]: C 64.08, H 8.72, N 3.32; found: C 64.26, H 8.40, N 3.20.

Received: January 24, 2009

Revised: March 18, 2009

Published online: June 12, 2009

Keywords: carbenes · homogeneous catalysis · hydrides · hydrogen · palladium

- [1] a) *Handbook of Homogeneous Hydrogenation* (Eds.: J. G. de Vries, C. J. Elsevier), Wiley-VCH, Weinheim, **2007**; b) B. Chen, U. Dingerdissen, J. G. E. Krauter, H. G. J. Lansink Rotgerink, K. Moebus, D. J. Ostgard, P. Panster, T. H. Riermeier, S. Seebald, T. Tacke, H. Trauthwein, *Appl. Catal. A* **2005**, *280*, 17–46; c) H.-U. Blaser, C. Malan, B. Pugin, F. Spindler, H. Steiner, M. Studer, *Adv. Synth. Catal.* **2003**, *345*, 103–151; d) S. Nishimura, *Handbook of Heterogeneous Catalytic Hydrogenation for Organic Synthesis*, Wiley, Chichester, UK, **2001**.
- [2] For a review on Wilkinson's catalyst, see: F. H. Jardine, *Prog. Inorg. Chem.* **1981**, *28*, 63–202.
- [3] a) A. Pfaltz, J. Blankenstein, R. Hilgraf, E. Hörmann, S. McIntyre, F. Menges, M. Schönleber, S. P. Smidt, B. Wüstenberg, N. Zimmermann, *Adv. Synth. Catal.* **2003**, *345*, 33–43; b) R. H. Crabtree, *Acc. Chem. Res.* **1979**, *12*, 331–337.
- [4] a) P. Pelagatti, A. Bacchi, M. Carcelli, M. Costa, A. Fochi, P. Ghidini, E. Leporati, M. Masi, C. Pelizzi, G. Pelizzi, *J. Organomet. Chem.* **1999**, *583*, 94–105; b) P. Pelagatti, A. Venturini, A. Leporati, M. Carcelli, M. Costa, A. Bacchi, G. Pelizzi, C. Pelizzi, *J. Chem. Soc. Dalton Trans.* **1998**, 2715–2722; c) V. V. S. Reddy, *J. Mol. Catal.* **1988**, *45*, 73–84; d) M. M. Taqui Khan, B. Taqui Khan, S. Begum, *J. Mol. Catal.* **1988**, *45*, 305–317; e) D. R. Armstrong, O. Novaro, M. E. Ruiz-Vizcaya, R. Linarte, *J. Catal.* **1977**, *48*, 8–20; f) G. Carturan, G. Strukul, *J. Organomet. Chem.* **1978**, *157*, 475–481; g) G. H. Olivé, S. Olivé, *J. Mol. Catal.* **1975**, *1*, 121–135; h) G. Henrici-Olivé, S. Olivé, *Angew. Chem.* **1974**, *86*, 561–562; *Angew. Chem. Int. Ed. Engl.* **1974**, *13*, 549–550; i) For an early review see: F. R. Hartley, *Chem. Rev.* **1969**, *69*, 799–844.
- [5] a) For a review on Pd⁰ cluster-mediated hydrogenation, see: I. I. Moiseev, M. N. Vargaftik, *New J. Chem.* **1998**, *22*, 1217–1227; b) R. van Asselt, C. J. Elsevier, *J. Mol. Catal.* **1991**, *65*, L13–L19; c) R. U. Kirss, R. Eisenberg, *Inorg. Chem.* **1989**, *28*, 3372–3378; d) E. W. Stern, P. K. Maples, *J. Catal.* **1972**, *27*, 120–133; e) Y. Takahashi, Ts. Ito, S. Sakai, Y. Ishii, *J. Chem. Soc. D* **1970**, 1065–1066.
- [6] a) L. Andrews, X. Wang, M. E. Alikhani, L. Manceron, *J. Phys. Chem. A* **2001**, *105*, 3052–3063; b) L. Andrews, L. Manceron, M. E. Alikhani, X. Wang, *J. Am. Chem. Soc.* **2000**, *122*, 11011–11012; c) G. A. Ozin, J. Garcia-Prieto, *J. Am. Chem. Soc.* **1986**, *108*, 3099–3100; d) C. Jarque, O. Novaro, M. E. Ruiz, J. Garcia-Prieto, *J. Am. Chem. Soc.* **1986**, *108*, 3507–3509; e) J. J. Low, W. A. Goddard III, *J. Am. Chem. Soc.* **1984**, *106*, 8321–8322; f) J. P. Collman, *Acc. Chem. Res.* **1968**, *1*, 136–143; g) R. Ugo, *Coord. Chem. Rev.* **1968**, *3*, 319–344; h) S. Carrá, R. Ugo, *Inorg. Chim. Acta* **1967**, *1*, 49–63; i) J. Halpern, *Adv. Catal.* **1959**, *11*, 301–344; j) J. Halpern, *J. Phys. Chem.* **1959**, *63*, 398–403.
- [7] a) M. W. van Laren, C. J. Elsevier, *Angew. Chem.* **1999**, *111*, 3926–3929; *Angew. Chem. Int. Ed.* **1999**, *38*, 3715–3717; b) C. Borriello, M. L. Ferrara, I. Orabona, A. Panunzi, F. Ruffo, *J. Chem. Soc. Dalton Trans.* **2000**, 2545–2550; c) M. W. van Laren, M. A. Duin, C. Klerk, M. Naglia, D. Rogolino, P. Pelegatti, A. Bacchi, C. Pelizzi, C. J. Elsevier, *Organometallics* **2002**, *21*, 1546–1553; d) D. Evrard, K. Groison, Y. Mugnier, P. D. Harvey, *Inorg. Chem.* **2004**, *43*, 790–796; e) J. W. Sprengers, J. Wassenaar, N. D. Clement, K. J. Cavell, C. J. Elsevier, *Angew. Chem.* **2005**, *117*, 2062–2065; *Angew. Chem. Int. Ed.* **2005**, *44*, 2026–2029; f) H. Guo, Z. Zheng, F. Yu, S. Ma, A. Holuigue, D. S. Tromp, C. J. Elsevier, Y. Yu, *Angew. Chem.* **2006**, *118*, 5119–5122; *Angew. Chem. Int. Ed.* **2006**, *45*, 4997–5000; g) H. Sajiki, S. Mori, T. Ohkubo, T. Ikawa, A. Kume, T. Maegawa, Y. Monguchi, *Chem. Eur. J.* **2008**, *14*, 5109–5111; h) P. Hauwert, G. Maestri, J. W. Sprengers, M. Catellani, C. J. Elsevier, *Angew. Chem.* **2008**, *120*, 3267–3270; *Angew. Chem. Int. Ed.* **2008**, *47*, 3223–3226.
- [8] a) A. M. Kluwer, T. S. Koblenz, T. Jonischkeit, K. Woelk, C. J. Elsevier, *J. Am. Chem. Soc.* **2005**, *127*, 15470–15480; b) J. López-Serrano, S. B. Duckett, S. Aiken, K. Q. Almeida Leñero, E. Drent, J. P. Dunne, D. Konya, A. C. Whitwood, *J. Am. Chem. Soc.* **2007**, *129*, 6513–6527.
- [9] To the best of our knowledge, the only report of an isolated monomeric palladium dihydride species was reported by Kudo et al. wherein the isolation, in 3.5% yield, of [Pd(PCy₃)₂(H)₂] from the reaction of [Pd(acac)₂], PCy₃, and AlEt₃ is disclosed. The complex was characterized solely by IR spectroscopy, see: K. Kudo, M. Hidai, Y. Uchida, *J. Organomet. Chem.* **1973**, *56*, 413–418.
- [10] G. S. McGrady, G. Guilera, *Chem. Soc. Rev.* **2003**, *32*, 383–392, and references therein.
- [11] a) M. D. Fryzuk, B. R. Lloyd, G. K. B. Clentsmith, S. J. Rettig, *J. Am. Chem. Soc.* **1994**, *116*, 3804–3812; b) S. Chan, W.-T. Wong, *J. Chem. Soc. Dalton Trans.* **1995**, 3987–3994; c) J. W.-S. Hiu, W.-T. Wong, *J. Chem. Soc. Dalton Trans.* **1997**, 1515–1520; d) S. Kuwata, S. Kabashima, Y. Ishii, M. Hidai, *J. Am. Chem. Soc.* **2001**, *123*, 3826–3827; e) S. Du, J. C. Jeffery, J. A. Kautz, X. L. Lu, T. D. McGrath, T. A. Miller, T. Riis-Johannessen, F. G. A. Stone, *Inorg. Chem.* **2005**, *44*, 2815–2825.
- [12] N. M. Scott, S. P. Nolan, *Eur. J. Inorg. Chem.* **2005**, 1815–1828.
- [13] S. Fantasia, S. P. Nolan, *Chem. Eur. J.* **2008**, *14*, 6987–6993.
- [14] V. Jurčík, S. P. Nolan, C. S. J. Cazin, *Chem. Eur. J.* **2009**, *15*, 2509–2511.
- [15] For more details, see the Supporting Information.
- [16] *trans* J(P,H) coupling constants are usually ≥ 160 Hz, whereas for *cis* couplings, values are 0–20 Hz; for selected examples, see: a) B. Kellenberger, S. J. Young, J. K. Stille, *J. Am. Chem. Soc.* **1985**, *107*, 6105–6107; b) P. Leoni, M. Sommovigo, M. Pasquali, S. Midollini, D. Braga, P. Sabatino, *Organometallics* **1991**, *10*, 1038–1044; c) K. Schager, K. Seevogel, K.-R. Pörschke, M. Kessler, C. Krüger, *J. Am. Chem. Soc.* **1996**, *118*, 13075–13076; d) P. J. Pérez, J. C. Calabrese, E. B. Bunel, *Organometallics* **2001**, *20*, 337–345; e) O. V. Ozerov, C. Guo, L. Fan, B. M. Foxman, *Organometallics* **2004**, *23*, 5573–5580; f) I. D. Hills, G. C. Fu, *J. Am. Chem. Soc.* **2004**, *126*, 13178–13179.
- [17] a) P. J. Desrosiers, L. Cai, Z. Lin, R. Richards, J. Halpern, *J. Am. Chem. Soc.* **1991**, *113*, 4173–4184; b) C. A. Bayse, R. L. Luck, E. J. Schelter, *Inorg. Chem.* **2001**, *40*, 3463–3467.
- [18] V. V. Grushin, *Chem. Rev.* **1996**, *96*, 2011–2033.
- [19] a) N. D. Clement, K. J. Cavell, C. Jones, C. J. Elsevier, *Angew. Chem.* **2004**, *116*, 1297–1299; *Angew. Chem. Int. Ed.* **2004**, *43*, 1277–1279; b) M. M. Konnick, B. A. Gandhi, I. A. Guzei, S. S. Stahl, *Angew. Chem.* **2006**, *118*, 2970–2973; *Angew. Chem. Int. Ed.* **2006**, *45*, 2904–2907.
- [20] a) D. S. McGuinness, K. J. Cavell, B. W. Skelton, A. H. White, *Organometallics* **1999**, *18*, 1596–1605; b) P. L. Arnold, F. G. N. Cloke, T. Geldbach, P. B. Hitchcock, *Organometallics* **1999**, *18*, 3228–3233.

- [21] Considering the overall relaxation rate of **2** and **3**, given by $R_2 = R_{HH} + R_{HP}$ and $R_3 = R_{HP}$, the HH contribution is calculated as $R_{HH} = R_2 - R_3 = (1/T_{12} - 1/T_{13})$.
- [22] H–H distances in $\eta^2\text{-H}_2$ transition-metal complexes are in the range 0.80–0.90 Å (compared to 0.74 Å for free H_2). See: a) R. H. Morris, J. F. Sawyer, M. Shiralian, J. D. Zubkowski, *J. Am. Chem. Soc.* **1985**, *107*, 5581–5582; b) G. J. Kubas, R. R. Ryan, B. I. Swanson, P. J. Vergamini, H. J. Wasserman, *J. Am. Chem. Soc.* **1984**, *106*, 451–452.
- [23] $J_{\text{H,D}}$ values in $\eta^2\text{-H}_2$ transition-metal complexes are in the range 30–35 Hz, whereas for complexes containing hydride and deuteride ligands, $J(\text{H,D}) < 3$ Hz. The slightly higher value detected in our case might be a result of the *trans* geometry, see for example: a) G. J. Kubas, C. J. Unkefer, B. I. Swanson, E. Fukushima, *J. Am. Chem. Soc.* **1986**, *108*, 7000–7009; b) G. J. Kubas, R. R. Ryan, D. A. Wroblewski, *J. Am. Chem. Soc.* **1986**, *108*, 1339–1341; c) R. Gelabert, M. Moreno, J. M. Lluch, A. Lledos, V. Pons, D. M. Heinekey, *J. Am. Chem. Soc.* **2004**, *126*, 8813–8822, and references therein.
- [24] a) S. Díez-González, S. P. Nolan, *Coord. Chem. Rev.* **2007**, *251*, 874–883; b) H. Jacobsen, A. Correa, A. Poater, C. Costabile, L. Cavallo, *Coord. Chem. Rev.* **2009**, *253*, 687–703; c) F. E. Hahn, M. C. Jahnke, *Angew. Chem.* **2008**, *120*, 3166–3216; *Angew. Chem. Int. Ed.* **2008**, *47*, 3122–3172; d) P. de Frémont, N. Marion, S. P. Nolan, *Coord. Chem. Rev.* **2009**, *253*, 862–892.
- [25] C. A. Tolman, *Chem. Rev.* **1977**, *77*, 313–348.
- [26] V. V. Grushin, C. Bensimon, H. Alper, *Inorg. Chem.* **1994**, *33*, 4804–4806.