


# Chemoselective Asymmetric N-Allylic Alkylation of Indoles with Morita–Baylis–Hillman Carbonates\*\*

Hai-Lei Cui, Xin Feng, Jing Peng, Jie Lei, Kun Jiang, and Ying-Chun Chen\*

The indole framework represents a key structural motif in a large number of biologically active natural products and pharmaceutical compounds.<sup>[1]</sup> Consequently, modifications on the indole structure,<sup>[2]</sup> including the development of enantioselective variants,<sup>[3]</sup> have triggered increasing interests. Because of the inherent nucleophilic characteristics of indole compounds, their reactions preferentially take place at the C3-position of the ring system. As a result, the majority of enantioselective reactions of indoles focus on the C3-selective addition of electron-rich indoles to electrophilic imines, epoxides, carbonyl compounds,  $\alpha,\beta$ -unsaturated carbonyl compounds, and nitroalkenes etc., thus leading to the formation of diversely structured enantioenriched C3-functionalized indole derivatives.<sup>[4]</sup> Recently, the enantioselective synthesis of C2-substituted indoles has also been realized through the asymmetric alkylation of 4,7-dihydroindoles and subsequent oxidation.<sup>[5]</sup>


In sharp contrast to the progress in enantioselective alkylation at the C3- or C2-positions, the asymmetric N-alkylation of indoles has been underdeveloped: probably because of the privileged C3-chemoselectivity of indole compounds. The limited examples include the palladium-catalyzed N-allylic alkylation of 3-substituted indoles developed by Trost et al.<sup>[6a]</sup> and the enantioselective intramolecular aza-Michael addition of tethered indole-2-carboxylates under chiral phase-transfer catalysis developed by Bandini et al.<sup>[6b]</sup> Indeed, N-alkylated indoles have been applied as useful intermediates for the synthesis of polyheterocycles<sup>[7]</sup> and occur widely among natural products and biologically active pharmaceuticals (Scheme 1).<sup>[8]</sup> For example, mitomycin C exhibits potent antitumour activity and is used clinically in the treatment of certain cancers.<sup>[8b]</sup> Yuremamine, which was recently isolated from the stem bark of *Mimosa hostilis*, shows hallucinogenic and psychoactive effects.<sup>[9]</sup> Pyrrolo[3,2,1-ij]-quinoline derivatives are active antihistamines and inhibitors of leukotriene, and they offer the possibility of a novel multimediator approach to the treatment of allergic disea-



Scheme 1. Biologically active N-alkylated indole derivatives.

ses.<sup>[8c]</sup> Therefore, it would be extremely desirable to develop effective protocols to access optically pure N-alkylated indoles.

Recently, the allylic alkylation of Morita–Baylis–Hillman (MBH) adducts catalyzed by a metal-free organic Lewis base has emerged as an attractive strategy to prepare multifunctional compounds.<sup>[10]</sup> Our research group has also reported that modified cinchona alkaloids are outstanding catalysts for the asymmetric allylic alkylation of  $\alpha,\alpha$ -dicyanoolefins with MBH carbonates.<sup>[11]</sup> We anticipated that, as outlined in Scheme 2, the deprotonation of the acidic NH group of the



Scheme 2. Proposed N- or C3-selective allylic alkylation of indoles with MBH carbonates catalyzed by a tertiary amine. Boc = *tert*-butoxycarbonyl.

indole ring by the *tert*-butoxy anion generated in situ would occur, and N- or C3-selective asymmetric allylic alkylation would followed to deliver valuable chiral indole derivatives.<sup>[12,13]</sup>

To our delight, we found that DABCO smoothly catalyzed the allylic alkylation of indole **1a** and MBH carbonate **2a** at 50°C. The reaction showed excellent chemoselectivity, and the N-alkylation compound **3a** was isolated as the sole product (Table 1, entry 1).<sup>[14]</sup> Encouraged by this preliminary

[\*] H.-L. Cui, X. Feng, J. Peng, J. Lei, K. Jiang, Prof. Dr. Y.-C. Chen  
Key Laboratory of Drug-Targeting and Drug Deliver System of the  
Education Ministry, Department of Medicinal Chemistry  
West China School of Pharmacy, Sichuan University  
Chengdu, 610041 (PR China)  
Fax: (+86) 28-8550-2609  
E-mail: ycchenhuaxi@yahoo.com.cn


Prof. Dr. Y.-C. Chen

State Key Laboratory of Biotherapy, West China Hospital  
Sichuan University, Chengdu, 610041 (PR China)

[\*\*] We are grateful for the financial support from NSFC (20772084).

Supporting information for this article is available on the WWW  
under <http://dx.doi.org/10.1002/anie.200902093>.

**Table 1:** Screening studies of organocatalytic N-allylic alkylation of indole **1a** with MBH carbonate **2a**.<sup>[a]</sup>




| Entry | Catalyst                 | Solvent           | <i>t</i> [h] | Yield [%] <sup>[b]</sup> | <i>ee</i> [%] <sup>[c]</sup> |
|-------|--------------------------|-------------------|--------------|--------------------------|------------------------------|
| 1     | DABCO                    | DCE               | 20           | 93                       | –                            |
| 2     | (DHQD) <sub>2</sub> AQN  | DCE               | 24           | 99                       | 24                           |
| 3     | (DHQD) <sub>2</sub> PHAL | DCE               | 42           | 96                       | 67                           |
| 4     | (DHQD) <sub>2</sub> PYR  | DCE               | 42           | 96                       | 48                           |
| 5     | (DHQ) <sub>2</sub> AQN   | DCE               | 42           | 62                       | –23                          |
| 6     | TMSQD                    | DCE               | 23           | 92                       | 12                           |
| 7     | BzQD                     | DCE               | 59           | 43                       | 0                            |
| 8     | (DHQD) <sub>2</sub> PHAL | toluene           | 59           | 79                       | 91                           |
| 9     | (DHQD) <sub>2</sub> PHAL | <i>m</i> -xylene  | 59           | 80                       | 91                           |
| 10    | (DHQD) <sub>2</sub> PHAL | mesitylene        | 59           | 82                       | 91                           |
| 11    | (DHQD) <sub>2</sub> PHAL | PhCF <sub>3</sub> | 36           | 92                       | 83                           |
| 12    | (DHQ) <sub>2</sub> PHAL  | mesitylene        | 60           | 76                       | –52                          |

[a] Unless otherwise noted, reactions were performed with 0.1 mmol of **1a**, 0.2 mmol of **2a**, and 0.01 mmol of catalyst in 1 mL solvent. [b] Yield of isolated product. [c] Determined by HPLC on a chiral stationary phase. DABCO = 1,4-diazabicyclo[2.2.2]octane, (DHQD)<sub>2</sub>AQN = hydroquinidine (anthraquinone-1,4-diyl) diether, (DHQD)<sub>2</sub>PHAL = hydroquinidine 1,4-phthalazinediyl diether, (DHQD)<sub>2</sub>PYR = hydroquinidine-2,5-diphenyl-4,6-pyrimidinediyl diether, (DHQ)<sub>2</sub>AQN = hydroquinine (anthraquinone-1,4-diyl) diether, TMSQD = 9-trimethylsilylquinidine, BzQD = 9-quinidyl benzoate, (DHQ)<sub>2</sub>PHAL = hydroquinine 1,4-phthalazinediyl diether, DCE = 1,2-dichloroethane.

result, we examined the enantioselective variant in the presence of chiral tertiary amine catalysts. A modified cinchona alkaloid<sup>[15]</sup> (DHQD)<sub>2</sub>AQN (10 mol %) showed high catalytic activity, and **3a** was obtained in quantitative yield after 24 hours, albeit with poor enantioselectivity (24% *ee*; Table 1, entry 2). Then we tested other tertiary amines derived from cinchona alkaloids (Table 1, entries 3–7) and found that (DHQD)<sub>2</sub>PHAL served as the best catalyst, which gave the product with 67% *ee* (Table 1, entry 3). Subsequently, we screened more parameters to optimize the reaction. Pleasingly, the choice of solvent had a significant effect on enantiocontrol. An excellent *ee* value (91%) was attained in toluene even though the reaction rate was slightly slower (Table 1, entry 8). Similar results were observed in *m*-xylene and mesitylene (Table 1, entries 9 and 10). A higher reactivity was observed in PhCF<sub>3</sub>, but the *ee* value was lower (Table 1, entry 11). In addition, (DHQ)<sub>2</sub>PHAL gave **3a** with the opposite configuration, although the enantioselectivity was modest under the optimal reaction conditions (Table 1, entry 12).

Next, the scope of the new N-allylic alkylation reaction was explored with a variety of indoles **1** and MBH carbonates **2**. The reaction was generally conducted with 10 mol % of (DHQD)<sub>2</sub>PHAL in mesitylene at 50 °C. The results are summarized in Table 2. In general, for MBH carbonate **2a**, variation in the substitution pattern on the indole unit was well tolerated. 3-Methylindole (**1b**) gave 83% *ee* in 93% yield (Table 2, entry 2). Notably, both 3-indolylpropanoate (**1c**) and the protected tryptamine **1d** gave high enantioselectivities with good yields (Table 2, entries 3 and 4). The effects of having a substituent at the 5-position of the indole unit was also explored (Table 2, entries 5–7). Indole **1f** bearing a strong electron-withdrawing group (NO<sub>2</sub>) gave decreased enantioselectivity, even though higher reactivity was observed (Table 2, entry 6).<sup>[13]</sup> 7-Bromoindole (**1h**) exhibited lower reactivity as a result of steric hindrance. Four equivalent of MBH carbonate **2a** was required to ensure a better conversion, and a moderate *ee* value was obtained (Table 2, entry 8). Importantly, indole-2-carboxylates **1i** and **1j** could be successfully alkylated. In these cases (DHQD)<sub>2</sub>AQN proved to be a superior catalyst in PhCF<sub>3</sub>. High enantioselectivities were afforded when the diphenylmethyl esters were used (Table 2, entries 9 and 10).<sup>[16]</sup> Moreover, a number of MBH carbonates have also been explored in the alkylation reaction with indoles. High enantioselectivities (81–93% *ee*) and moderate to good yields (71–99%) were attained for MBH carbonates bearing electron-deficient or electron-rich aryl or heteroaryl groups (Table 2, entries 11–

**Table 2:** Asymmetric N-allylic alkylation of indoles **1** with MBH carbonates **2**.<sup>[a]</sup>



**1a** R<sup>1</sup> = R<sup>2</sup> = R<sup>3</sup> = R<sup>4</sup> = H

**1b** R<sup>1</sup> = R<sup>3</sup> = R<sup>4</sup> = H, R<sup>2</sup> = Me

**1c** R<sup>1</sup> = R<sup>3</sup> = R<sup>4</sup> = H

R<sup>2</sup> = CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>Me

**1d** R<sup>1</sup> = R<sup>3</sup> = R<sup>4</sup> = H

R<sup>2</sup> = CH<sub>2</sub>CH<sub>2</sub>NPhth

**1e** R<sup>1</sup> = R<sup>2</sup> = R<sup>4</sup> = H, R<sup>3</sup> = Br

**1f** R<sup>1</sup> = R<sup>2</sup> = R<sup>4</sup> = H, R<sup>3</sup> = NO<sub>2</sub>

**1g** R<sup>1</sup> = R<sup>2</sup> = R<sup>4</sup> = H, R<sup>3</sup> = MeO

**1h** R<sup>1</sup> = R<sup>2</sup> = R<sup>3</sup> = H, R<sup>4</sup> = Br

**1i** R<sup>1</sup> = COOCHPh<sub>2</sub>

R<sup>2</sup> = R<sup>3</sup> = R<sup>4</sup> = H

**1j** R<sup>1</sup> = COOCHPh<sub>2</sub>

R<sup>2</sup> = R<sup>4</sup> = H, R<sup>3</sup> = Me


| Entry             | <b>1</b>  | Ar              | <i>t</i> [h] | Product   | Yield [%] <sup>[b]</sup> | <i>ee</i> [%] <sup>[c]</sup> |
|-------------------|-----------|-----------------|--------------|-----------|--------------------------|------------------------------|
| 1                 | <b>1a</b> | Ph              | 59           | <b>3a</b> | 82                       | 91                           |
| 2                 | <b>1b</b> | Ph              | 94           | <b>3b</b> | 93                       | 83                           |
| 3                 | <b>1c</b> | Ph              | 116          | <b>3c</b> | 68                       | 90                           |
| 4                 | <b>1d</b> | Ph              | 54           | <b>3d</b> | 69                       | 85                           |
| 5                 | <b>1e</b> | Ph              | 54           | <b>3e</b> | 97                       | 81                           |
| 6 <sup>[d]</sup>  | <b>1f</b> | Ph              | 66           | <b>3f</b> | 91                       | 62                           |
| 7                 | <b>1g</b> | Ph              | 46           | <b>3g</b> | 99                       | 87                           |
| 8 <sup>[e]</sup>  | <b>1h</b> | Ph              | 90           | <b>3h</b> | 79                       | 71                           |
| 9 <sup>[f]</sup>  | <b>1i</b> | Ph              | 31           | <b>3i</b> | 91                       | 90                           |
| 10 <sup>[f]</sup> | <b>1j</b> | Ph              | 24           | <b>3j</b> | 91                       | 86                           |
| 11                | <b>1a</b> | <i>p</i> -MeOPh | 38           | <b>3k</b> | 72                       | 89                           |
| 12                | <b>1a</b> | <i>p</i> -ClPh  | 47           | <b>3l</b> | 89                       | 80                           |
| 13                | <b>1a</b> | 2-furyl         | 16           | <b>3m</b> | 77                       | 90                           |
| 14                | <b>1a</b> | <i>m</i> -MePh  | 67           | <b>3n</b> | 71                       | 87                           |
| 15                | <b>1d</b> | <i>p</i> -ClPh  | 96           | <b>3o</b> | 79                       | 85 <sup>[g]</sup>            |
| 16 <sup>[f]</sup> | <b>1i</b> | <i>p</i> -MeOPh | 40           | <b>3p</b> | 99                       | 93                           |

[a] Unless otherwise noted, reactions were performed with 0.1 mmol of **1**, 0.2 mmol of **2**, and 10 mol % of (DHQD)<sub>2</sub>PHAL in 1.0 mL mesitylene at 50 °C. [b] Yield of isolated product. [c] Determined by HPLC on a chiral stationary phase. [d] At 25 °C. [e] 0.4 mmol of **2a** was used. [f] With 10 mol % of (DHQD)<sub>2</sub>AQN in PhCF<sub>3</sub>. [g] The absolute configuration of **3o**<sup>[17]</sup> was determined by X-ray analysis (see the Supporting Information). The absolute configurations of the other products were assigned by analogy. Phth = phthaloyl.

lectivities with good yields (Table 2, entries 3 and 4). The effects of having a substituent at the 5-position of the indole unit was also explored (Table 2, entries 5–7). Indole **1f** bearing a strong electron-withdrawing group (NO<sub>2</sub>) gave decreased enantioselectivity, even though higher reactivity was observed (Table 2, entry 6).<sup>[13]</sup> 7-Bromoindole (**1h**) exhibited lower reactivity as a result of steric hindrance. Four equivalent of MBH carbonate **2a** was required to ensure a better conversion, and a moderate *ee* value was obtained (Table 2, entry 8). Importantly, indole-2-carboxylates **1i** and **1j** could be successfully alkylated. In these cases (DHQD)<sub>2</sub>AQN proved to be a superior catalyst in PhCF<sub>3</sub>. High enantioselectivities were afforded when the diphenylmethyl esters were used (Table 2, entries 9 and 10).<sup>[16]</sup> Moreover, a number of MBH carbonates have also been explored in the alkylation reaction with indoles. High enantioselectivities (81–93% *ee*) and moderate to good yields (71–99%) were attained for MBH carbonates bearing electron-deficient or electron-rich aryl or heteroaryl groups (Table 2, entries 11–


16). MBH carbonates with  $\beta$ -alkyl substitution were tested but showed much lower reactivity.

We have expanded the substrate scope by replacing the indole unit with methyl pyrrole-2-carboxylate **4**. The desired N-allylic alkylation product **5** was obtained in 80 % yield with 73 % *ee* (Scheme 3).<sup>[18]</sup>



**Scheme 3.** N-Allylic alkylation of pyrrole-2-carboxylate.

Apart from the traditional derivation of the C=C bond of the allylic products,<sup>[19]</sup> some fused-ring systems could be efficiently synthesized from the newly prepared N-allylic indole compounds. Although the direct oxidative Heck reaction of **3b** resulted in limited success,<sup>[2d]</sup> a regioselective bromination was easily realized by simply treating **3b** with NBS (Scheme 4). The subsequent palladium-catalyzed intra-



**Scheme 4.** Synthesis of the fused, cyclic indole systems. AIBN = 2,2'-azobisisobutyronitrile, NBS = *N*-bromosuccinimide.

molecular Heck reaction of **6** afforded the desired pyrrolo-[1,2-*a*]indole framework **7**, which possesses the core structure of the natural product Yuremamine (61 % yield, 80 % *ee*; see Scheme 1).<sup>[20]</sup> On the other hand, an intramolecular radical cyclization or Heck reaction of **3h** could be smoothly conducted to deliver pyrrolo[3,2,1-*ij*]quinoline derivatives **8** and **9**, respectively. These derivatives might find valuable applications in medicinal chemistry.<sup>[8c]</sup>

In conclusion, we have developed the first asymmetric and chemoselective N-allylic alkylation of indoles with Morita-Baylis-Hillman carbonates. This method was achieved by employing metal-free catalysis of modified cinchona alkaloids. Either electron-rich or electron-deficient indoles could be employed and moderate to excellent enantioselectivity was achieved (62–93 % *ee*). This reaction also provides a convenient method to prepare multiply functionalized pyrrolo[1,2-*a*]indole and pyrrolo[3,2,1-*ij*]quinoline deriva-

tives, which may have potential use in the synthesis of biologically important materials. This work is currently underway in our laboratory.

Received: April 19, 2009

Revised: May 18, 2009

Published online: June 24, 2009

**Keywords:** chemoselectivity · indoles · Morita-Baylis-Hillman carbonates · N-allylic alkylation · organocatalysis

- [1] a) D. J. Faulkner, *Nat. Prod. Rep.* **2002**, *19*, 1; b) M. Amat, N. Llor, J. Bosch, X. Solans, *Tetrahedron* **1997**, *53*, 719; c) A. Kleeman, J. Engel, B. Kutscher, D. Reichert, *Pharmaceutical Substances*, 4th ed., Thieme, New York, **2001**.
- [2] For recent reviews, see: a) S. Cacchi, G. Fabrizi, *Chem. Rev.* **2005**, *105*, 2873; b) G. R. Humphrey, J. T. Kueth, *Chem. Rev.* **2006**, *106*, 2875; c) G. Zeni, R. C. Larock, *Chem. Rev.* **2006**, *106*, 4644; for selected examples, see: d) N. P. Grimster, C. Gauntlett, C. R. A. Godfrey, M. J. Gaunt, *Angew. Chem.* **2005**, *117*, 3185; *Angew. Chem. Int. Ed.* **2005**, *44*, 3125; e) C. Liu, X. Han, X. Wang, R. A. Widenhoefer, *J. Am. Chem. Soc.* **2004**, *126*, 3700; f) H. Huang, R. Peters, *Angew. Chem.* **2009**, *121*, 612; *Angew. Chem. Int. Ed.* **2009**, *48*, 604.
- [3] For recent reviews, see: a) M. Bandini, A. Melloni, A. Umani-Ronchi, *Angew. Chem.* **2004**, *116*, 560; *Angew. Chem. Int. Ed.* **2004**, *43*, 550; b) M. Bandini, S. Tommasi, A. Umani-Ronchi, *Synlett* **2005**, 1199; c) T. B. Poulsen, K. A. Jørgensen, *Chem. Rev.* **2008**, *108*, 2903.
- [4] For selected examples, see: a) J. F. Austin, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2002**, *124*, 1172; b) R. P. Herrera, V. Sgarzani, L. Bernardi, A. Ricci, *Angew. Chem.* **2005**, *117*, 6734; *Angew. Chem. Int. Ed.* **2005**, *44*, 6576; c) K. B. Jensen, J. Thorhauge, R. G. Hazell, K. A. Jørgensen, *Angew. Chem.* **2001**, *113*, 164; *Angew. Chem. Int. Ed.* **2001**, *40*, 160; d) H. Li, Y.-Q. Wang, L. Deng, *Org. Lett.* **2006**, *8*, 4063; e) D. A. Evans, K. R. Fandrick, H.-J. Song, *J. Am. Chem. Soc.* **2005**, *127*, 8942; f) Q. Kang, Z.-A. Zhao, S.-L. You, *J. Am. Chem. Soc.* **2007**, *129*, 1484; g) Y.-X. Jia, J. Zhong, S.-F. Zhu, C.-M. Zhang, Q.-L. Zhou, *Angew. Chem.* **2007**, *119*, 5661; *Angew. Chem. Int. Ed.* **2007**, *46*, 5565; h) A. J. Boersma, B. L. Feringa, G. Roelfes, *Angew. Chem.* **2009**, *121*, 3396; *Angew. Chem. Int. Ed.* **2009**, *48*, 3346.
- [5] a) D. A. Evans, K. R. Fandrick, *Org. Lett.* **2006**, *8*, 2249; b) D. A. Evans, K. R. Fandrick, H.-J. Song, K. A. Scheidt, R. Xu, *J. Am. Chem. Soc.* **2007**, *129*, 10029; c) Q. Kang, X.-J. Zheng, S.-L. You, *Chem. Eur. J.* **2008**, *14*, 3539; d) Y.-F. Sheng, G.-Q. Li, Q. Kang, A.-J. Zhang, S.-L. You, *Chem. Eur. J.* **2009**, *15*, 3351.
- [6] a) B. M. Trost, M. J. Krische, V. Berl, E. M. Grenzer, *Org. Lett.* **2002**, *4*, 2005; b) M. Bandini, A. Eichholzer, M. Tragni, A. Umani-Ronchi, *Angew. Chem.* **2008**, *120*, 3282; *Angew. Chem. Int. Ed.* **2008**, *47*, 3238.
- [7] a) G. A. Molander, M. H. Schmitt, *J. Org. Chem.* **2000**, *65*, 3767; b) M. Tanaka, M. Ubukata, T. Matsuo, K. Yasue, K. Matsumoto, Y. Kajimoto, T. Ogo, T. Inaba, *Org. Lett.* **2007**, *9*, 3331.
- [8] a) E. O. M. Orlemans, W. Verboom, M. W. Scheltinga, D. N. Reinhoudt, P. Lelieveld, H. H. Fiebig, B. R. Winterhalter, J. A. Double, M. C. Bibby, *J. Med. Chem.* **1989**, *32*, 1612; b) U. Galm, M. H. Hager, S. G. V. Lanen, J. Ju, J. S. Thorson, B. Shen, *Chem. Rev.* **2005**, *105*, 739; c) D. Paris, M. Cottin, P. Demonchaux, G. Augert, P. Dupassieux, P. Lenoir, M. J. Peck, D. Jasserand, *J. Med. Chem.* **1995**, *38*, 669; d) R. M. Wilson, R. K. Thalji, R. G. Bergman, J. A. Ellman, *Org. Lett.* **2006**, *8*, 1745; e) L. S.

- Fernandez, M. S. Buchanan, A. R. Carroll, Y. J. Feng, R. J. Quinn, V. M. Avery, *Org. Lett.* **2009**, *11*, 329.
- [9] J. J. Vepsäläinen, S. Auriola, M. Tukiainen, N. Ropponen, J. Callaway, *J. Planta Med.* **2005**, *71*, 1049.
- [10] a) C.-W. Cho, M. J. Krische, *Angew. Chem.* **2004**, *116*, 6857; *Angew. Chem. Int. Ed.* **2004**, *43*, 6689; b) D. J. V. C. van Steenis, T. Marcelli, M. Lutz, A. L. Spek, J. H. van Maarseveen, H. Hiemstra, *Adv. Synth. Catal.* **2007**, *349*, 281; c) Y.-Q. Jiang, Y.-L. Shi, M. Shi, *J. Am. Chem. Soc.* **2008**, *130*, 7202.
- [11] H.-L. Cui, J. Peng, X. Feng, W. Du, K. Jiang, Y.-C. Chen, *Chem. Eur. J.* **2009**, *15*, 1574.
- [12] For metal-catalyzed C3-selective allylic alkylation of indoles, see: a) W.-B. Liu, H. He, L.-X. Dai, S.-L. You, *Org. Lett.* **2008**, *10*, 1815; b) A. B. Zaitsev, S. Gruber, P. A. Plüss, P. S. Pregosin, L. F. Veiros, M. Wörle, *J. Am. Chem. Soc.* **2008**, *130*, 11604; c) M. Bandini, A. Melloni, F. Piccinelli, R. Sinisi, S. Tommasi, A. Umani-Ronchi, *J. Am. Chem. Soc.* **2006**, *128*, 1424; d) H. Y. Cheung, W.-Y. Yu, F. L. Lam, T. T.-L. Au-Yeung, Z. Zhou, T. H. Chan, A. S. C. Chan, *Org. Lett.* **2007**, *9*, 4295; either C3- or N-selective allylic alkylation products could be obtained in a racemic form, see: e) M. Bandini, A. Melloni, A. Umani-Ronchi, *Org. Lett.* **2004**, *6*, 3199.
- [13] For a base-catalyzed C3-selective alkylation of indoles, see: M. Bandini, R. Sinisi, *Org. Lett.* **2009**, *11*, 2093.
- [14] The desired N-allylic alkylated product **3a** could not be obtained when the reaction was catalyzed by PPh<sub>3</sub>.
- [15] For a review, see: S.-K. Tian, Y. Chen, J. Hang, L. Tang, P. McDaid, L. Deng, *Acc. Chem. Res.* **2004**, *37*, 621.
- [16] For more screening studies, see the Supporting Information.
- [17] CCDC 732289 (**3o**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- [18] B. M. Trost, G. Dong, *J. Am. Chem. Soc.* **2006**, *128*, 6054.
- [19] a) I. T. Raheem, E. N. Jacobsen, *Adv. Synth. Catal.* **2005**, *347*, 1701; b) see the Supporting Information.
- [20] For the synthesis of the core structure of *Yuremamine* in five steps, see: a) M. B. Johansen, M. A. Kerr, *Org. Lett.* **2008**, *10*, 3497; b) H. Kusama, J. Takaya, N. Iwasawa, *J. Am. Chem. Soc.* **2002**, *124*, 11592.