

A Journal of the Gesellschaft Deutscher Chemiker


# Angewandte Chemie

International Edition

GDCh

www.angewandte.org

2009–48/42



## Although +VI ...

... has long been the highest experimentally known oxidation state of Ir, the +VIII oxidation state has been produced and identified in low-temperature noble gas matrices of iridium tetroxide. M. F. Zhou, S. Riedel, and co-workers describe in their Communication on page 7879 ff. how the compound is formed by co-depositing Ir atoms with mixtures of O<sub>2</sub> and a noble gas onto a CsI surface at 4–6 K. Theoretical analysis indicates that it is a d<sup>1</sup> species with the formal oxidation state Ir<sup>VIII</sup>.

 WILEY-VCH

## Inside Cover

**Yu Gong, Mingfei Zhou,\* Martin Kaupp, and Sebastian Riedel\***

**Although +VI** has long been the highest experimentally known oxidation state of Ir, the +VIII oxidation state has been produced and identified in low-temperature noble gas matrices of iridium tetroxide. M. F. Zhou, S. Riedel, and co-workers describe in their Communication on page 7879 ff. how the compound is formed by co-depositing Ir atoms with mixtures of O<sub>2</sub> and a noble gas onto a CsI surface at 4–6 K. Theoretical analysis indicates that it is a d<sup>1</sup> species with the formal oxidation state Ir<sup>VIII</sup>.

