

A Journal of the Gesellschaft Deutscher Chemiker


Angewandte Chemie

International Edition



www.angewandte.org

2009–48/47



The catalytic conversion ...

... of thiophene derivatives on individual zeolite ZSM-5 crystals was studied by using a combination of multiplex coherent anti-Stokes Raman scattering (CARS) and synchrotron-based IR microscopy. In their Communication on page 8990 ff., B. M. Weckhuysen et al. show how label-free chemical imaging of reactant/reaction products can be achieved with high spatial resolution. Vibrational bands in the CARS and IR spectra evidenced interaction of thiophenes with the zeolite framework, followed by ring opening.

WILEY-VCH

Inside Cover

Marianne H. F. Kox, Katrin F. Domke, James P. R. Day, Gianluca Rago, Eli Stavitski, Mischa Bonn, and Bert M. Weckhuysen*

The catalytic conversion of thiophene derivatives on individual zeolite ZSM-5 crystals was studied by using a combination of multiplex coherent anti-Stokes Raman scattering (CARS) and synchrotron-based IR microscopy. In their Communication on page 8990 ff., B. M. Weckhuysen et al. show how label-free chemical imaging of reactant/reaction products can be achieved with high spatial resolution. Vibrational bands in the CARS and IR spectra evidenced interaction of thiophenes with the zeolite framework, followed by ring opening.

