

A Journal of the Gesellschaft Deutscher Chemiker


# Angewandte Chemie

International Edition



[www.angewandte.org](http://www.angewandte.org)

2009–48/48



## Morphological control ...

... of high-symmetry small-molecular organic microcrystals was achieved by changing the solvent composition and precursor solubility, which substantially alter the crystal growth rate in the  $\langle 100 \rangle$  direction relative to that in the  $\langle 110 \rangle$  direction. In their Communication on page 9121 ff., X. H. Zhang, S. T. Lee, and co-workers describe the surfactant-assisted synthesis of microcrystals with selected shapes that range from cubes, through truncated cubes, to rhombic dodecahedra.

 WILEY-VCH

## Inside Cover

**Xiujuan Zhang,\* Cheng Dong, Juan Antonio Zapien, Shafiq Ismathullakhan, Zhenhui Kang, Jiansheng Jie, Xiaohong Zhang,\* Jack C. Chang, Chun-Sing Lee, and Shuit-Tong Lee\***

**Morphological control** of high-symmetry small-molecular organic microcrystals was achieved by changing the solvent composition and precursor solubility, which substantially alter the crystal growth rate in the  $\langle 100 \rangle$  direction relative to that in the  $\langle 110 \rangle$  direction. In their Communication on page 9121 ff., X. H. Zhang, S. T. Lee, and co-workers describe the surfactant-assisted synthesis of microcrystals with selected shapes that range from cubes, through truncated cubes, to rhombic dodecahedra.

