


Chiral Phosphoric Acid Catalyzed Desymmetrization of *meso*-1,3-Diones: Asymmetric Synthesis of Chiral Cyclohexenones**

Keiji Mori, Takuya Katoh, Tohru Suzuki, Takuya Noji, Masahiro Yamanaka, and Takahiko Akiyama*

Cyclohexenones are important building blocks in synthetic organic chemistry. In particular, Hajos–Parrish^[1] and Wieland–Miescher^[2] ketones are useful synthetic intermediates not only for the preparation of steroids^[3] but also for a range of natural products.^[4] The most facile and conventional method used to obtain these ketones in enantiomerically pure form is the desymmetrization of *meso*-1,3-dicarbonyl compounds, in which (*S*)-proline is commonly used as a highly reliable chiral catalyst (Scheme 1).^[1,2] This protocol consists of two consecutive transformations: 1) desymmetrization of



Scheme 1. Synthetic strategy for the preparation of chiral cyclohexenones.

[*] Dr. K. Mori, T. Katoh, T. Suzuki, T. Noji, Prof. Dr. T. Akiyama
Department of Chemistry, Faculty of Science
Gakushuin University
1-5-1, Mejiro, Toshima-ku, Tokyo, 171-8588 (Japan)
Fax: (+81) 3-5992-1029
E-mail: takahiko.akiyama@gakushuin.ac.jp
Homepage: http://www-cc.gakushuin.ac.jp/~940020/akiyama-site/index_e.html

Prof. Dr. M. Yamanaka
Department of Chemistry, Rikkyo University
3-34-1 Nishi-Ikebukuro, Toshima-ku, Tokyo 171-8501 (Japan)

[**] This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (Japan).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200905271>.

meso-1,3-dione by (*S*)-proline-catalyzed aldol reaction, and 2) dehydration.

As a result of our initial findings,^[5] chiral phosphoric acids **1** derived from (*R*)-BINOL have been extensively studied as versatile chiral Brønsted acid catalysts. They exhibited remarkable asymmetric inducing ability in the nucleophilic addition to imine, the 1,4-addition to α,β -unsaturated compounds, and the transfer hydrogenation with Hantzsch ester.^[6] Although the asymmetric ring-opening of *meso*-aziridines by means of chiral phosphoric acid had already been reported by Antilla and co-workers,^[7a] the chiral phosphoric acid induced desymmetrization of *meso*-1,3-diones still remains a challenge.^[7b,c] The control of stereoselectivity by weak, noncovalent bond interaction (hydrogen bond) is not a trivial issue in comparison with the control by covalent bonds ((*S*)-proline catalysis).^[8] Herein, we report the asymmetric synthesis of synthetically useful chiral cyclohexenones through the desymmetrization of *meso*-1,3-dicarbonyl compounds induced by a chiral phosphoric acid.^[9] By this method, both desymmetrization of the 1,3-dione compound and dehydration could be accomplished in a single-step, one-pot operation, to afford chiral cyclohexenones with excellent enantioselectivity.

An initial study was conducted by treatment of triketone **2** with 10 mol % of **1a** in toluene. Gratifyingly, enone **3a** was obtained in the enantioenriched form (46% *ee*; Table 1,

Table 1: Screening of catalyst (*R*)-**1**.^[a]

Entry	Ar	Yield [%] ^[b]	<i>ee</i> [%] ^[c]
1	Ph (1a)	56 (43)	46
2	SiPh ₃ (1b)	85 (5)	84
3	9-anthryl (1c)	91 (4)	70
4	2,4,6-(<i>i</i> Pr) ₃ C ₆ H ₂ (1d)	92 (5)	90
5 ^[d]	2,4,6-(<i>i</i> Pr) ₃ C ₆ H ₂ (1d)	> 99	90
6 ^[d,e]	2,4,6-(<i>i</i> Pr) ₃ C ₆ H ₂ (1d)	95	90
7 ^[f]	(<i>S</i>)-proline	57 ^[g]	–60

[a] Unless otherwise noted, all reactions were conducted with 0.2 mmol of **2a** in toluene (2.0 mL). [b] Determined by ¹H NMR spectroscopy. Amount of recovered starting material in parenthesis. [c] Determined by HPLC on a chiral stationary phase using a Daicel Chiralcel OD-H column; flow rate = 0.5 mL min^{–1}; eluent = *n*-hexane/*i*PrOH = 5:1. [d] *n*-Hexane was used as the reaction solvent. [e] 5 mol % catalyst loading. [f] Triketone **2** was treated with 10 mol % of (*S*)-proline in DMF at 25 °C for 48 h, and the resulting product was treated with TsOH·H₂O (10 mol %) in benzene at reflux for 20 min. [g] Yield of isolated product. DMF = *N,N*-dimethylformamide, Ts = 4-toluenesulfonyl.

entry 1), even when the reaction was performed at 70°C. Screening of catalysts and reaction conditions (Table 1, entries 2–5) revealed that the use of 5 mol % of 2,4,6-triisopropylphenyl-substituted catalyst **1d** in *n*-hexane at gentle reflux (70°C) was the optimal set of the conditions, and gave **3a** in excellent yield and enantioselectivity (95 %, 90 % *ee*; Table 1, entry 6).^[10] Significantly, the selectivity of the chiral phosphoric acid was overwhelmingly higher than that of (*S*)-proline (57 %, 60 % *ee*; Table 1, entry 7).

The generality of this method was examined under the optimal reaction conditions (Table 2). Ethyl-substituted **3b** was obtained in excellent yield and enantioselectivity (89 %, 92 % *ee*;

Table 2: Substrate scope of the desymmetrization reaction.^[a]

Entry	Product	<i>t</i> [h]	Yield [%]	<i>ee</i> [%] ^[b]
1	3b	48	89	92
2	3c	72	82	94
3 ^[c]	3d	96	94	87
4 ^[c,d]	3e	48	72	90
5 ^[c]	3f	24	90	84
6 ^[c]	3g	24	86	70
7 ^[c]	3h	24	64	82

[a] Unless otherwise noted, all reactions were conducted with 0.2 mmol of **2** and 5 mol % **1d** in *n*-hexane (2.0 mL) at 70°C. [b] Determined by HPLC on a chiral stationary phase. [c] 10 mol % of **1d** was employed. [d] In toluene at 90°C.

92 % *ee*; Table 2, entry 1). In the case of propargyl substrate **3c**, a key synthetic intermediate in the synthesis of the gibbane framework,^[11] the enantioselectivity was also excellent (94 % *ee*; Table 2, entry 2). Benzyl- and phenyl-substituted substrates (**3d** and **3e**) were obtained in good enantioselectivity (87 and 90 % *ee*; Table 2, entries 3 and 4). In these two reactions, higher catalyst loading (10 mol %) and prolonged reaction times (96 and 48 h, respectively) were required. In particular, in the case of the phenyl-substituted substrate **3e** the use of toluene as the reaction solvent was

essential owing to the poor solubility of the substrate. Excellent yield and enantioselectivity were also observed in the case of naphthalene derivative **3f** (90 %, 84 % *ee*; Table 2, entry 5). Chiral phosphoric acid was applied to the asymmetric synthesis of Hajos–Parrish and Wieland–Miescher ketones (**3g** and **3h**), and gave moderate to high yields with good enantioselectivity (70 and 82 % *ee*; Table 2, entries 6 and 7).^[12] The absolute configurations of these cyclohexenone derivatives were surmised by analogy with **3a**.

To clarify the origin of the enantioselectivity, ONIOM (B3LYP/6-31G*:HF/3-21G) calculations^[13] were carried out based on the transition state (TS) controlled by hydrogen bonding, as shown in Scheme 1. In the TS, chiral phosphoric acid could simultaneously activate carbonyl and enol moieties with Brønsted acidic and Lewis basic sites, respectively.^[14] The attack of the TS from the *si* face (**TS_{si}**) was 1.3 kcal mol^{−1} more stable than attack from the *re* face (**TS_{re}**), which is in agreement with the experimental results (Figure 1). The




Figure 1. 3D structures and schematic representation of models of **TS_{re}** and **TS_{si}** calculated by ONIOM (B3LYP/−31G*:HF/3-21G).

energy difference would be mainly caused by the steric repulsion between the aryl moiety of the substrate and the 3,3'-triisopropylphenyl group.

In summary, we have developed the first example of chiral phosphoric acid catalyzed desymmetrization of *meso*-1,3-dicarbonyl compounds. This method could be applied to a wide variety of substrates to give chiral cyclohexenones in high yields and with excellent enantioselectivity. Further investigation of its application to the synthesis of natural products is currently under way in our laboratory.

Experimental Section

General procedure for Table 1, entry 6: Chiral phosphoric acid **1d** (7.5 mg, 0.010 mmol) was added to a solution of triketone **2a** (46 mg, 0.20 mmol) in *n*-hexane (2.0 mL) at room temperature, and the reaction mixture was heated at 70°C. After heating for 24 h, the reaction mixture was directly purified by column chromatography on

silica gel (eluent: CH₂Cl₂/AcOEt 10:1) to give **3a** (40 mg, 0.19 mmol) in 95% yield.

Received: September 21, 2009

Published online: November 18, 2009

Keywords: aldol reaction · asymmetric synthesis · chiral phosphoric acids · computational chemistry · desymmetrization

- [1] a) U. Eder, G. Sauer, R. Wiechert, *Angew. Chem.* **1971**, 83, 492–493; *Angew. Chem. Int. Ed. Engl.* **1971**, 10, 496–497; b) Z. G. Hajos, D. R. Parrish, *J. Org. Chem.* **1974**, 39, 1615–1621; c) R. A. Micheli, Z. G. Hajos, N. Cohen, D. R. Parrish, L. A. Poland, W. Sciamanna, M. A. Scott, P. A. Wehrli, *J. Org. Chem.* **1975**, 40, 675–681; d) Z. G. Hajos, D. R. Parrish, *Org. Synth., Collect. Vol. VII*, Wiley, New York, **1990**, pp. 363–369.
- [2] a) P. Wieland, K. Miescher, *Helv. Chim. Acta* **1950**, 33, 2215–2228; b) J. Gutzwiller, A. Buchschacher, A. Furst, *Synthesis* **1977**, 167–168; c) S. Ramachandran, M. S. Newman, *Org. Synth.* **1961**, 41, 38.
- [3] For selected references, see: a) P. Wieland, H. Unberwasser, G. Annre, K. Miescher, *Helv. Chim. Acta* **1953**, 36, 1231–1241; b) P. Wieland, G. Anner, K. Miescher, *Helv. Chim. Acta* **1953**, 36, 1803–1809; c) T. A. Spencer, T. D. Weaver, R. M. Villarica, R. J. Friary, J. Posler, M. A. Schwartz, *J. Org. Chem.* **1968**, 33, 712–719; d) S. J. Danishefsky, P. Cain, *J. Am. Chem. Soc.* **1975**, 97, 5282–5284.
- [4] For selected references, see: a) E. J. Corey, M. Ohno, R. B. Mitra, P. A. Vatakencherry, *J. Am. Chem. Soc.* **1964**, 86, 478–485; b) E. J. Corey, M. Ohno, R. B. Mitra, R. A. Vatakencherry, *J. Am. Chem. Soc.* **1964**, 86, 478–485; c) J. E. McMurry, S. J. Isser, *J. Am. Chem. Soc.* **1972**, 94, 7132–7137; d) R. C. A. Isaacs, R. M. J. Di Grandi, S. J. Danishefsky, *J. Org. Chem.* **1993**, 58, 3938–3941; e) L. A. Paquette, T.-Z. Wang, M. R. Sivik, *J. Am. Chem. Soc.* **1994**, 116, 11323–11334; f) T. J. Reddy, G. Bordeaux, L. Trimble, *Org. Lett.* **2006**, 8, 5585–5588; g) S. Yamashita, K. Iso, M. Hiram, *Org. Lett.* **2008**, 10, 3413–3415.
- [5] a) T. Akiyama, J. Itoh, K. Yokota, K. Fuchibe, *Angew. Chem.* **2004**, 116, 1592–1594; *Angew. Chem. Int. Ed.* **2004**, 43, 1566–1568; b) D. Uraguchi, M. Terada, *J. Am. Chem. Soc.* **2004**, 126, 5356–5357.
- [6] For reviews on Brønsted acid catalysis, see: a) T. Akiyama, J. Itoh, K. Fuchibe, *Adv. Synth. Catal.* **2006**, 348, 999–1010; b) T. Akiyama, *Chem. Rev.* **2007**, 107, 5744–5758; c) S. J. Connon, *Angew. Chem.* **2006**, 118, 4013–4016; *Angew. Chem. Int. Ed.* **2006**, 45, 3909–3912; d) T. Akiyama in *Acid Catalysis in Modern Organic Synthesis* (Eds.: H. Yamamoto, K. Ishihara), Wiley-VCH, Weinheim, **2008**, pp. 62–107; e) M. Terada, *Chem. Commun.* **2008**, 4097–4112, and references therein.
- [7] a) E. B. Rowland, G. B. Rowland, E. Rivera-Otero, J. C. Antilla, *J. Am. Chem. Soc.* **2007**, 129, 12084–12085; b) D. K. Sala, A. Lattanzi, *Org. Lett.* **2009**, 11, 3330–3333; c) G. L. Hamilton, T. Kanai, F. D. Toste, *J. Am. Chem. Soc.* **2008**, 130, 14984–14986.
- [8] a) S. Bahmanyar, K. N. Houk, *J. Am. Chem. Soc.* **2001**, 123, 12911–12912; b) L. Hoang, S. Bahmanyar, K. N. Houk, B. List, *J. Am. Chem. Soc.* **2003**, 125, 16–17; c) C. Allemann, R. Gordillo, F. R. Clemente, H.-Y. Cheong, K. N. Houk, *Acc. Chem. Res.* **2004**, 37, 558–569.
- [9] For our previous approach to chiral cyclohexenones, see: T. Akiyama, T. Katoh, K. Mori, *Angew. Chem.* **2009**, 121, 4290–4292; *Angew. Chem. Int. Ed.* **2009**, 48, 4226–4228.
- [10] The absolute configuration of **3a** was unambiguously established by single-crystal X-ray analysis of the corresponding bromide. CCDC 743645 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif. See the Supporting Information.
- [11] S. Takano, C. Kasahara, K. Ogasawara, *J. Chem. Soc. Chem. Commun.* **1981**, 635–637.
- [12] The absolute configuration of Hajos–Parrish and Wieland–Miescher ketones were determined by comparing optical rotation values with those reported. E. Lacoste, E. Vaique, M. Berlande, I. Pianet, J. M. Vincent, Y. Landais, *Eur. J. Org. Chem.* **2007**, 167–177.
- [13] Details of chemical models of ONIOM calculations are shown in the Supporting Information. All calculations were performed with the Gaussian 03 package, see: Gaussian03, Revision E.01., M. J. Frisch, et al. Gaussian, Inc., Wallingford CT, **2004**.
- [14] For theoretical studies on chiral phosphoric acid catalysis, see: a) M. Yamanaka, J. Itoh, K. Fuchibe, T. Akiyama, *J. Am. Chem. Soc.* **2007**, 129, 6756–6764; b) L. Simón, J. M. Goodman, *J. Am. Chem. Soc.* **2008**, 130, 8741–8747; c) T. Marcelli, P. Hammar, F. Himo, *Chem. Eur. J.* **2008**, 14, 8562–8571; d) L. Simón, J. M. Goodman, *J. Am. Chem. Soc.* **2009**, 131, 4070–4077; e) M. Yamanaka, T. Hirata, *J. Org. Chem.* **2009**, 74, 3266–3271; f) T. Akiyama, H. Morita, P. Bachu, K. Mori, M. Yamanaka, T. Hirata, *Tetrahedron* **2009**, 65, 4950–4956.