



On these pages, we feature a selection of the excellent work that has recently been published in our sister journals. If you are reading these pages on a

computer, click on any of the items to read the full article. Otherwise please see the DOIs for easy online access through Wiley InterScience.




Lithium Carbenoids

V. Capriati,* S. Florio*

Anatomy of Long-lasting Love Affairs with Lithium Carbenoids: Past and Present Status and Future Prospects

Face-to-face, rising to the challenge! The two Janus-like souls of lithium carbenoids are not a weak but a strength point of their reactivity, instead. A proper knowledge of their structural features, aggregation and solvation can indeed shed light not only on the reasons of their thermal and configurational lability/stability, but can also allow a fine tuning of their reactivity toward more stereoselective and targeted transformations.



Chem. Eur. J.

DOI: 10.1002/chem.200902870




Gold Nanostructures

C.-C. Huang, * Y.-L. Hung, Y.-C. Shiang, T.-Y. Lin, Y.-S. Lin, C.-T. Chen,* H.-T. Chang*

Photoassisted Synthesis of Luminescent Mannose-Au Nanodots for the Detection of Thyroglobulin in Serum

Thyroglobulin assays: Photoassisted synthesis of water-soluble and highly luminescent mannose-modified gold nanodots (Man-Au NDs) by irradiation with an LED have been employed for the detection of cancer marker thyroglobulin (Tg) through an assay based on competition between Tg and Man-Au NDs for interaction with the concanavalin A (Con A).



Chem. Asian J.

DOI: 10.1002/asia.200900346




PDZ Domains

R. Gallardo, Y. Ivarsson, J. Schymkowitz, F. Rousseau, P. Zimmermann*

Structural Diversity of PDZ-Lipid Interactions

Partners in crime: PDZ domains are among the most abundant proteininteraction domains in multicellular organisms. This review provides an overview of the recent progress in research on the role of lipid interactions in the biology of PDZ scaffolds. We summarize the current understanding of PDZ-lipid interactions and the consequences for peptide interactions by emphasizing the structural and biochemical aspects.



ChemBioChem

DOI: 10.1002/cbic.200900616

... on our Sister Journals

Biomembranes


X. Han, A. S. Achalkumar, M. R. Cheetham, S. D. A. Connell, B. R. G. Johnson, R. J. Bushby, S. D. Evans*

A Self-assembly Route for Double Bilayer Lipid Membrane Formation

Biomembranes: A new method of forming double bilayer lipid membranes uses NHS/EDC chemistry to link adjacent bilayers, both of which remain fluid. This approach provides a novel platform for the study of biomembranes, in which the components of the upper membrane are shielded from the solid substrate by a second membrane (see figure); and for studying more complex membrane protein systems which span double lipid bilayers.







Molecular Modeling

A. Stary,* S. J. Wacker, L. Boukharta, U. Zachariae, Y. Karimi-Nejad, J. Åqvist, G. Vriend, B. L. de Groot


Toward a Consensus Model of the hERG Potassium Channel

To achieve a consensus model of hERG we tested seven models based on different S5 alignments by using a combination of static assessment, MD simulations, and docking. Model 6 fulfills all quality criteria and is confirmed by recent mutagenesis experiments.





ChemMedChem




Solar Cells

P. A. Troshin,* E. A. Khakina, M. Egginger, A. E. Goryachev, S. I. Troyanov, A. Fuchsbauer, A. S. Peregudov, R. N. Lyubovskaya, V. F. Razumov, N. S. Sariciftci

Self-Assembly of Thiophene- and Furan-Appended Methanofullerenes with Poly(3-Hexylthiophene) in Organic Solar Cells


Fullerene derivatives with pendant thiophene or furan units selfassemble with poly(3-hexylthiophene) via π - π stacking and form ordered nanostructures, enabling their efficient operation in photovoltaic devices. Such intermolecular stabilization might be considered as a promising way to improve the lifetime of organic photovoltaics in the future.



ChemSusChem

DOI: 10.1002/cssc.200900196

Methanol Synthesis



S. Schimpf, A. Rittermeier, X. Zhang, Z.-A. Li, M. Spasova, M. W. E. van den Berg, M. Farle, Y. Wang, R. A. Fischer, M. Muhler*

Stearate-Based Cu Colloids in Methanol Synthesis: Structural Changes Driven by Strong Metal-Support Interactions

Grand prisms: Zinc stearate-stabilized copper nanoparticles, synthesized by an efficient one-step process, are applied in the continuous liquid-phase synthesis of methanol. The active state is identified as triangular ZnO prisms with truncated edges attached to the spherical Cu particles, which act as a reservoir for ZnO_x species diffusing onto the Cu particles and promoting the catalytic activity.



ChemCatChem DOI: 10.1002/cctc.200900252






Ru Oxaphosphacyclic Carbenes

S. Bolaño, * J. Bravo, * J. Castro, M. Rodríguez-Rocha

Formation of Oxaphosphacarbenes by Reaction of [RuTpCl(PPh₃) (thp)] with Terminal Alkynes and a Propargylic Alcohol

Cationic and neutral five-membered cyclic ruthenium oxaphosphacarbenes were obtained by intramolecular nucleophilic addition of one hydroxymethyl group of the tris (hydroxymethyl) phosphane (thp) ligand to the α -carbon atom of vinylidene or allenylidene intermediates, which were generated by reaction of [TpRuCl(PPh₃) (PTA)] with terminal alkynes and with 1,1-diphenyl-2-propyn-1-ol.



Eur. J. Inorg. Chem.

DOI: 10.1002/ejic.200900988




Dynamic Kinetic Resolution

J. H. Lee, K. Han, M.-J. Kim,* J. Park*

Chemoenzymatic Dynamic Kinetic Resolution of Alcohols and Amines

Dynamic kinetic resolution (DKR) is an attractive process for the production of optically active compounds from racemic mixtures. Combinations of the processes of metal-catalyzed racemization and enzymatic acylation in one-pot fashion have shown the potential of DKR of alcohols and amines. This Microreview summarizes developments in and the scope of chemoenzymatic DKR.



Eur. J. Org. Chem.

DOI: 10.1002/ejoc.200900935

