

A Journal of the Gesellschaft Deutscher Chemiker


Angewandte Chemie

International Edition

GDCh

www.angewandte.org

2010–49/14



C–H activation and reduction ...

... are both apparent in the reaction of $[\text{Sm}^{\text{III}}(\text{AlMe}_4)_3]$ with the heterocycle $(\text{CyNCH}_2)_3$ (Cy = cyclohexyl). As shown by N. W. Mitzel and co-workers in their Communication on page 2611 ff., C–H bonds of methyl groups are activated to form a carbyne complex with an $\{\text{(HC)Al}_3\text{Sm}\}$ unit, and spontaneous reduction to samarium(II) by methyl groups can also be observed.

 WILEY-VCH

Inside Cover

Daniel Bojer, Ajay Venugopal, Beate Neumann, Hans-Georg Stammler, and Norbert W. Mitzel*

C–H activation and reduction are both apparent in the reaction of $[\text{Sm}^{\text{III}}(\text{AlMe}_4)_3]$ with the heterocycle $(\text{CyNCH}_2)_3$ (Cy = cyclohexyl). As shown by N. W. Mitzel and co-workers in their Communication on page 2611 ff., C–H bonds of methyl groups are activated to form a carbyne complex with an $\{(\text{HC})\text{Al}_3\text{Sm}\}$ unit, and spontaneous reduction to samarium(II) by methyl groups can also be observed.

