



Organoiridium complexes ...

... are noteworthy scaffolds for the design of enzyme inhibitors. As described by E. Meggers and co-workers in their Communication on page 3839 ff., a nanomolar and selective iridium-based inhibitor for the receptor protein kinase VEGFR3 (Flt4) has been discovered. This nontoxic octahedral iridium(III) complex, which is synthesized through stereoselective oxidative addition starting from a suitable square-planar iridium(I) precursor, is revealed to interfere with the development of blood vessels *in vivo* in two different zebrafish angiogenesis models.

Inside Cover

**Alexander Wilbuer, Danielle H. Vlecken, Daan J. Schmitz,
Katja Kräling, Klaus Harms, Christoph P. Bagowski, and Eric Meggers***

Organoiridium complexes are noteworthy scaffolds for the design of enzyme inhibitors. As described by E. Meggers and co-workers in their Communication on page 3839 ff., a nanomolar and selective iridium-based inhibitor for the receptor protein kinase VEGFR3 (Flt4) has been discovered. This nontoxic octahedral iridium(III) complex, which is synthesized through stereoselective oxidative addition starting from a suitable square-planar iridium(I) precursor, is revealed to interfere with the development of blood vessels *in vivo* in two different zebrafish angiogenesis models.

