

A Journal of the Gesellschaft Deutscher Chemiker


Angewandte Chemie

International Edition



www.angewandte.org

2010–49/25



A geometrically unprecedented ...

... dodecanuclear nickel(II) elliptical cluster and two decanuclear nickel(II) wheel clusters have been obtained by stepwise introduction of different thiolates at a nickel(II) center, as described by C. Zhang, M. G. Humphrey, K. Tatsumi et al. in their Communication on page 4209 ff. The novel toroidal molecular configurations and strong wavelength-dependent nonlinear optical behavior in these “tiara-like” chromophores have been shown by theoretical studies to be driven by Ni–S π -bonding interactions.

WILEY-VCH

Inside Cover

Chi Zhang,* Tsuyoshi Matsumoto, Marek Samoc, Simon Petrie, Suci Meng, T. Christopher Corkery, Robert Stranger, Jinfang Zhang, Mark G. Humphrey,* and Kazuyuki Tatsumi*

A geometrically unprecedented dodecanuclear nickel(II) elliptical cluster and two decanuclear nickel(II) wheel clusters have been obtained by stepwise introduction of different thiolates at a nickel(II) center, as described by C. Zhang, M. G. Humphrey, K. Tatsumi et al. in their Communication on page 4209 ff. The novel toroidal molecular configurations and strong wavelength-dependent nonlinear optical behavior in these “tiara-like” chromophores have been shown by theoretical studies to be driven by Ni–S π -bonding interactions.

