

A Journal of the Gesellschaft Deutscher Chemiker


# Angewandte Chemie

International Edition



www.angewandte.org

2010–49/36



## Photoactive inorganic–organic nanohybrids ...

... are constructed from CdTe quantum dots (QDs) and single-walled carbon nanotubes (SWNTs). In their Communication on page 6425 ff. D. M. Guldi and co-workers report that the introduction of surface stabilizers to QDs facilitates both the covalent attachment of a pyrene group and the anchoring onto the SWNTs. Complementary luminescence and ultrafast transient absorption analyses indicate the formation of a charge-separated state; this paves the way for the successful integration into photovoltaic devices.

 WILEY-VCH

## Inside Cover

**Christian Schulz-Drost, Vito Sgobba, Christina Gerhards,  
Susanne Leubner, Rafael M. Krick Calderon, Andrés Ruland, and  
Dirk M. Guldi\***

**Photoactive inorganic–organic nanohybrids** are constructed from CdTe quantum dots (QDs) and single-walled carbon nanotubes (SWNTs). In their Communication on page 6425 ff. D. M. Guldi and co-workers report that the introduction of surface stabilizers to QDs facilitates both the covalent attachment of a pyrene group and the anchoring onto the SWNTs. Complementary luminescence and ultrafast transient absorption analyses indicate the formation of a charge-separated state; this paves the way for the successful integration into photovoltaic devices.

