

A Journal of the Gesellschaft Deutscher Chemiker


# Angewandte Chemie

International Edition



www.angewandte.org

2010–49/40



## The role of quantum tunneling ...

... of hydrogen atoms in the formation of dihydrogen in space has been investigated. In their Communication on page 7350 ff., T.P.M. Goumans and J. Kästner use a novel implementation of harmonic quantum transition-state theory to study the reaction rate of hydrogen atoms with benzene. This model of the rate-limiting step of H<sub>2</sub> formation on carbonaceous dust grains in space shows that hydrogen reacts about two orders of magnitude faster than the heavier deuterium. Background image: NASA/JPL-Caltech/STScI/CXC/UofA/ESA/AURA/JHU.

 WILEY-VCH

## Inside Cover

**Theodorus P. M. Goumans\* and Johannes Kästner**

**The role of quantum tunneling** of hydrogen atoms in the formation of dihydrogen in space has been investigated. In their Communication on page 7350 ff., T.P.M. Goumans and J. Kästner use a novel implementation of harmonic quantum transition-state theory to study the reaction rate of hydrogen atoms with benzene. This model of the rate-limiting step of  $\text{H}_2$  formation on carbonaceous dust grains in space shows that hydrogen reacts about two orders of magnitude faster than the heavier deuterium. Background image: NASA/JPL-Caltech/STScI/CXC/UofA/ESA/AURA/JHU.

