

A Journal of the Gesellschaft Deutscher Chemiker


Angewandte Chemie

International Edition



www.angewandte.org

2010–49/42



A one-dimensional n-type ...

... semiconducting nanotube is formed by the self-assembly of a naphthalenetetracarboxylic acid diimide-lysine bolaamphiphile. In their Communication on page 7688 ff., J. R. Parquette and co-workers show how the nanotubes aggregate into a monolayer membrane that curves into rings, which then stack into tubes. Solid-state NMR studies show that there is exceptional conformational homogeneity among the constituent molecules that comprise the nanotubes, leading to efficient energy migration within the assemblies.

 WILEY-VCH

Inside Cover

**Hui Shao, James Seifert, Natalie C. Romano, Min Gao,
Jonathan J. Helmus, Christopher P. Jaroniec, David A. Modarelli, and
Jon R. Parquette***

A one-dimensional n-type semiconducting nanotube is formed by the self-assembly of a naphthalenetetracarboxylic acid diimide–lysine bolaamphiphile. In their Communication on page 7688 ff., J. R. Parquette and co-workers show how the nanotubes aggregate into a monolayer membrane that curves into rings, which then stack into tubes. Solid-state NMR studies show that there is exceptional conformational homogeneity among the constituent molecules that comprise the nanotubes, leading to efficient energy migration within the assemblies.

