

A Journal of the Gesellschaft Deutscher Chemiker


# Angewandte Chemie

50  
YEARS

International Edition

www.angewandte.org

2011–50/7



## Two types of $\text{HF}_2^-$ bridges ...

... are found in the rectangular polymeric network formed from  $\text{Cu}^{\text{II}}$ ,  $\text{FHF}^-$ , and pyrazine (pyz). The resulting linear  $\text{Cu}(\text{FHF})_2\text{-Cu}$  chains are linked into 2D sheets by pyz units. The  $\text{FHF}^-$  coordination in the  $\text{Cu}(\text{FHF})_2\text{-Cu}$  chains mimics motifs found in transition metal/azide systems. In their Communication on page 1573 ff. J. L. Manson et al. also show that stronger spin exchange occurs along the  $\text{HF}_2^-$  bridges than the  $\text{Cu-pyz-Cu}$  bridges.

 WILEY-VCH

## Back Cover

**Jamie L. Manson,\* Michelle L. Warter, John A. Schlueter, Tom Lancaster, Andrew J. Steele, Stephen J. Blundell, Francis L. Pratt, John Singleton, Ross D. McDonald, Changhoon Lee, Myung-Hwan Whangbo, and Alex Plonczak**

**Two types of  $\text{HF}_2^-$  bridges** are found in the rectangular polymeric network formed from  $\text{Cu}^{\text{II}}$ ,  $\text{FHF}^-$ , and pyrazine (pyz). The resulting linear  $\text{Cu}-(\text{FHF})_2\text{-Cu}$  chains are linked into 2D sheets by pyz units. The  $\text{FHF}^-$  coordination in the  $\text{Cu}-(\text{FHF})_2\text{-Cu}$  chains mimics motifs found in transition metal/azide systems. In their Communication on page 1573 ff. J. L. Manson et al. also show that stronger spin exchange occurs along the  $\text{HF}_2^-$  bridges than the  $\text{Cu-pyz-Cu}$  bridges.

