


Total Synthesis of Aspergillide A and B Based on the Transannular Oxy-Michael Reaction**


Makoto Kanematsu, Masahiro Yoshida, and Kozo Shishido*

The aspergillides A, B, and C (**1**, **2**, and **3**) comprise a novel class of 14-membered macrolides that were isolated by



Kusumi and co-workers from the marine-derived fungus *Aspergillus ostianus* strain 01F313 that was cultured in a medium composed of bromine-modified artificial sea water.^[1] Their structures were determined by extensive spectroscopic studies, and their absolute configurations were established by X-ray crystallography (for **1** and **2**) and the modified Mosher method (for **3**).^[2] These compounds contain tri-substituted tetrahydro and dihydropyran units and exhibit cytotoxicity against mouse lymphocytic leukemia cells (L1210) with LD₅₀ values of 2.1, 71.0, and 2.0 μg mL⁻¹, respectively. Because of their intriguing structural features and biological profile, these polyketides have attracted much attention in the synthetic community as targets for total synthesis. To date, four total syntheses of **1**,^[3] six of **2**,^[3c,4] and two of **3**^[5] have been reported. Herein, we describe our total synthesis of the aspergillides A (**1**) and B (**2**) from a common macrolide intermediate by employing an interesting transannular oxy-Michael reaction^[6] as the key step.


Our strategy for the synthesis of the aspergillides A and B, illustrated in Scheme 1, proposes the formation of the trisubstituted pyran moiety through a base-mediated transannular oxy-Michael reaction from the 14-membered macrolactone **4**. It was thought that stereochemical control at the newly generated stereogenic center at C3 could be achieved by the choice of reaction conditions. The macrolactone **4** could be assembled by using a sequential cross-metathesis



Scheme 1. Retrosynthetic analysis. Bn = benzyl, HWE = Horner–Wadsworth–Emmons, TBS = *tert*-butyldimethylsilyl.

and intramolecular Horner–Wadsworth–Emmons reaction of the acyclic precursors **5**, which could be accessed from the chiral building block **7**^[7] and compound **6**, which was known in the literature.^[8] Two stereogenic centers at the future C4 and C7 positions in **5** would be created by taking advantage of the convex nature of **7**.

The optically pure enone **7** with a bicyclo[3.2.1]octane framework, prepared from 2-furfural by a six-step sequence, served as the starting material for the synthesis of the alkenyl alcohol **14** (Scheme 2). Attempted transformation of **7** into **10** using the Wharton transposition^[7,9] gave unsatisfactory results (≈20% yield), prompting us to find another route. Reduction with NaBH₄ followed by the Hata reaction^[10] of



Scheme 2. Synthesis of the alkenyl alcohol **14**. Reagents and conditions: a) NaBH₄, CeCl₃·7H₂O, MeOH, RT, 1 h, 98%; b) PhSSPh, *n*Bu₃P, pyridine, 60°C, 16 h, 86%; c) *m*CPBA, CH₂Cl₂, -78°C, 1.5 h; then (MeO)₃P, EtOH, reflux, 6 h, 71%; d) TBSCl, imidazole, DMAP, CH₂Cl₂, RT, 1 h, 94%; e) H₂, Pd(OH)₂-C, THF, 60°C, 8 h, 99%; f) MsCl, Et₃N, CH₂Cl₂, RT, 1 h; g) LiI, THF, reflux, 8 h, 94% (over 2 steps); h) Zn, EtOH, reflux, 3 h, 94%; i) LiBH₄, THF, 0°C, 2 h, quant.; j) PivCl, Et₃N; then TESCl, DMAP, CH₂Cl₂, RT, 1 h; k) LiBH₄, THF, 0°C, 3 h, 96% (over 2 steps, based on recovered diol **13**). DMAP = 4-dimethylaminopyridine, *m*CPBA = *m*-chloroperoxybenzoic acid, Ms = methanesulfonyl, Piv = pivaloyl, TES = triethylsilyl, THF = tetrahydrofuran.


[*] M. Kanematsu, Dr. M. Yoshida, Prof. Dr. K. Shishido
Graduate School of Pharmaceutical Sciences
The University of Tokushima
1-78-1 Shomachi, Tokushima 770-8505 (Japan)
Fax: (+81) 88-633-9575
E-mail: shishido@ph.tokushima-u.ac.jp

[**] This work was supported financially by a Grant-in-Aid for the Program for Promotion of Basic and Applied Research for Innovations in the Bio-oriented Technology Research Advancement Institution (BRAIN).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201007327>.

the resulting **8** produced the inverted sulfide **9**, which was treated with *m*CPBA and (MeO)₃P in ethanol at reflux^[11] to give the alcohol **10** in 60 % yield from **7**. After protection of the alcohol moiety as the TBS ether, sequential hydrogenation, debenzylization, mesylation, and iodination gave the iodide **11**, which was reduced with zinc in ethanol at reflux to afford the hemiacetal **12** in good overall yield as a 1.5:1 mixture of diastereoisomers at the acetal carbon atom. Reduction with LiBH₄ gave the diol **13**, the primary and the secondary alcohols of which were sequentially protected as the pivalate and the TES ether, respectively. The pivalate was reduced with LiBH₄ to give the alkenyl alcohol **14**.

Cross-metathesis^[12] of **14** with the phosphonoacetate **6**,^[8] prepared from (*R*)-2-methyloxirane through a two-step sequence, in the presence the second-generation Grubbs catalyst (5 mol %) in methylene chloride at reflux provided the coupled product as the *E* alkene (> 20:1) in 98 % yield (Scheme 3). Oxidation of **16** with Dess–Martin periodinane and a subsequent intramolecular Horner–Wadsworth–



Scheme 3. Synthesis of the macrolactone **4**. DBU = 1,8-diazabicyclo-[5.4.0]undec-7-ene, DMP = Dess–Martin periodinane, PPTS = pyridinium 4-toluenesulfonate.

Emmons reaction^[13] gave the macrolactone **17** in 78 % yield over the two steps. Selective removal of the TES ether was realized by treatment with PPTS to afford the requisite **4**, a substrate ready for the transannular cyclization.


With the macrolactone in hand, we next investigated the crucial transannular oxy-Michael reaction (Table 1). Treatment of **4** with DBU in acetonitrile at reflux provided in 50 % yield a 1:6 mixture of the cyclized *syn*- and *anti*-pyrans **18** and **19**, which were separable by column chromatography (entry 1). When LiCl was used with DBU, the reaction proceeded faster, and the cycloadduct was obtained quantitatively and with 7:1 *syn* selectivity (entry 2). Encouraged by this result, we next proceeded to examine more closely the reaction conditions. It turned out that the best result was obtained by treatment of **4** with DBU and LiCl in acetonitrile at room temperature for 1.5 hours, and provided a quantitative amount of the *syn*-adduct **18** as a single product (entry 3). This result can be explained by invoking the lithium-chelated transition-state **20**, in which the conformation is quite similar to the crystal structure^[2] of aspergillide A, as shown in

Table 1: Transannular oxy-Michael reaction of **4**.

Entry	Reaction conditions	Yield [%] ^[a]	18/19
1	DBU (10 equiv), CH ₃ CN, reflux, 10 h	50	1:6
2	DBU (10 equiv), LiCl (10 equiv) CH ₃ CN, reflux, 2 h	quant.	7:1
3	DBU (10 equiv), LiCl (10 equiv), CH ₃ CN, RT, 1.5 h	quant.	1:0
4	KH (1.1 equiv), THF, 0 °C, 0.5 h	90	1.6:1
5	KH (1.1 equiv), THF, 0 °C, 2 h	91	0:1
6	KH (1.1 equiv), [18]crown-6 (5 equiv), THF, 0 °C, 0.5 h	96	0:1

[a] Yield of isolated product.


Scheme 4. From these results it was thought that the *anti*-isomer **19** might be the thermodynamic product. Consequently, for the selective formation of **19**, we examined the



Scheme 4. Transition state for the conversion of **4** into **18**.

reaction under the conditions described in entry 1. Prolonged reaction times did not result in increased yields of the desired *anti* adduct; rather decomposition was observed. Treatment of **4** with KH in THF at 0 °C for 0.5 hours produced the pyrans in a ratio of 1.6:1 in 90 % yield (entry 4). However, prolonged exposure (2 h) of **4** to the same reaction conditions resulted in the exclusive formation of **19** in comparable yield (entry 5). It was also determined that the reaction could be accelerated (≈ 0.5 h) by the addition of [18]crown-6 to give the *anti*-adduct **19** in 96 % yield (entry 6). Thus, we have demonstrated that the diastereoselectivity of the addition could be altered completely by changing the reaction conditions.


To prove the thermodynamic preference, interconversion between **18** and **19** was attempted (Scheme 5). Exposure of the *syn*-adduct **18** to the conditions shown in entry 6 provided the *anti*-isomer **19** in 94 % yield;^[14] on the other hand, treatment of **19** under the conditions of entry 2 for 12 hours



Scheme 5. Attempted interconversion.

resulted in 90 % recovery of the starting material. From these results, the *anti*-isomer **19** was found to be thermodynamically favored.

Both macrocyclic pyrans **18** and **19** were independently treated with acidic conditions (3M aqueous HCl in THF at room temperature for 3 h) to give quantitatively the aspergillides A (**1**) and B (**2**), which were spectroscopically identical to the natural products (Scheme 6 and the Support-



Scheme 6. Completion of the total syntheses and attempted conversions.

ing Information). Furthermore, the attempted conversion of natural aspergillide A into aspergillide B under the same conditions as for **18** was also successful. In addition, exposure of **1** to the isolation conditions (SiO₂ in MeOH/CHCl₃ (1:3) at room temperature) did not produce **2** at all, which indicated that aspergillide B is not an artifact.

In summary, the total synthesis of the aspergillides A and B has been accomplished in a longest linear sequence of 17 steps in 30 % and 28 % yield, respectively, from readily accessible bicyclic chiral building blocks. The unique features of this work include the first application of a highly efficient and diastereoselective transannular oxy-Michael reaction for the construction of the *syn* and *anti* tetrahydropyrans, which are the penultimate intermediates for the natural products. From a common 14-membered macrolactone, this synthesis is the first demonstration of the successful conversion of the *syn* pyran into the *anti* isomer, including the conversion of the natural aspergillide A into aspergillide B by employing basic conditions for a short period of time. The transannular oxy-Michael strategy developed here would be applicable to the synthesis of other related natural products.

Received: November 22, 2010

Revised: January 4, 2010

Published online: February 15, 2011

Keywords: diastereoselectivity · macrocycles · Michael addition · polyketides · total synthesis

- [1] K. Kito, R. Ookura, S. Yoshida, M. Namikoshi, T. Ooi, T. Kusumi, *Org. Lett.* **2008**, *10*, 225–228.
- [2] R. Ookura, K. Kito, Y. Saito, T. Kusumi, T. Ooi, *Chem. Lett.* **2009**, *38*, 384.
- [3] Syntheses of aspergillide A: a) T. Nagasawa, S. Kuwahara, *Tetrahedron Lett.* **2010**, *51*, 875–877; b) S. Diaz-Oltra, C. A. Angulo-Pachon, J. Murga, M. Carda, J. A. Marco, *J. Org. Chem.* **2010**, *75*, 1775–1778; c) H. Fuwa, H. Yamaguchi, M. Sasaki, *Org. Lett.* **2010**, *12*, 1848–1851; H. Fuwa, H. Yamaguchi, M. Sasaki, *Tetrahedron* **2010**, *66*, 7492–7503; d) G. Sabitha, D. V. Reddy, A. S. Rao, J. S. Yadav, *Tetrahedron Lett.* **2010**, *51*, 4195–4198.
- [4] Syntheses of aspergillide B: a) S. M. Hande, J. Uenishi, *Tetrahedron Lett.* **2009**, *50*, 189–192; b) S. Diaz-Oltra, C. A. Angulo-Pachon, M. N. Kneeteman, J. Murga, M. Carda, J. A. Marco, *Tetrahedron Lett.* **2009**, *50*, 3783–3785; c) T. Nagasawa, S. Kuwahara, *Biosci. Biotechnol. Biochem.* **2009**, *73*, 1893–1894; d) J. Liu, K. Xu, J. He, L. Zhang, X. Pan, X. She, *J. Org. Chem.* **2009**, *74*, 5063–5066; e) A. J. Mueller, M. P. Jennings, *Tetrahedron Lett.* **2010**, *51*, 4260–4262.
- [5] Syntheses of aspergillide C: a) T. Nagasawa, S. Kuwahara, *Org. Lett.* **2009**, *11*, 761–764; b) J. D. Panarese, S. P. Waters, *Org. Lett.* **2009**, *11*, 5086–5088.
- [6] L. Furst and C. N. Boddy have reported the scope of transannular oxy-Michael additions for the synthesis of medium-sized bicyclic systems; presented at the Regional Meeting of the American Chemical Society, Binghamton, NY, United States, October 5–7, **2006** (abstract pages NRM-149).
- [7] a) M. Takeuchi, T. Taniguchi, K. Ogasawara, *Synthesis* **1999**, 341–354; b) T. Yoshimura, F. Yakushiji, S. Kondo, X. Wu, M. Shindo, K. Shishido, *Org. Lett.* **2006**, *8*, 475–478.
- [8] Y. Wang, D. Romo, *Org. Lett.* **2002**, *4*, 3231–3234.
- [9] P. S. Wharton, D. H. Bohlen, *J. Org. Chem.* **1961**, *26*, 3615–3616.
- [10] I. Nakagawa, K. Aki, T. Hata, *J. Chem. Soc. Perkin Trans. 1* **1983**, 1315–1318.
- [11] a) P. Bickart, F. W. Carson, J. Jacobus, E. G. Miller, K. Mislow, *J. Am. Chem. Soc.* **1968**, *90*, 4869–4876; b) D. A. Evans, G. C. Andrews, C. L. Sims, *Acc. Chem. Res.* **1974**, *7*, 147–155; c) E. N. Prilezhaeva, *Russ. Chem. Rev.* **2001**, *70*, 897–920.
- [12] The second-generation Grubbs catalyst: a) M. Scholl, S. Ding, C. W. Lee, R. H. Grubbs, *Org. Lett.* **1999**, *1*, 953–956; b) T. M. Trnka, J. P. Morgan, M. S. Sanford, T. E. Wilhelm, M. Scholl, T. L. Choi, S. Ding, M. W. Day, R. H. Grubbs, *J. Am. Chem. Soc.* **2003**, *125*, 2546–2558; cross-metathesis: A. K. Chatterjee, T. L. Choi, D. P. Sanders, R. H. Grubbs, *J. Am. Chem. Soc.* **2003**, *125*, 11360–11370.
- [13] M. A. Blanchette, W. Choy, J. T. Davis, A. P. Esserfeld, S. Masamune, W. R. Roush, T. Sakai, *Tetrahedron Lett.* **1984**, *25*, 2183–2186.
- [14] It has been mentioned in reference [3a] that an isomerization of the *syn* pyran into the *anti* isomer was observed in trace amounts during the macrolactonization.