

Corrigenda

Catalytic Selective Cyclizations of
Aminocyclopropanes: Formal Synthesis
of Aspidospermidine and Total Synthesis
of Goniomitine

F. De Simone, J. Gertsch,
J. Waser* 5767–5770

Angew. Chem. Int. Ed. **2010**, 49

DOI 10.1002/anie.201001853

During recent investigations on the antiproliferative effect of goniomitine, different results were obtained than originally reported in this Communication (10.1002/anie.201001853). The new values are given in the updated version of Table 2 below. The reasons for the higher activities (150–450 nm) observed originally are still not clear at this point, as no difference in the sample purity was observed.

Table 2: Antiproliferative activity of goniomitine.

Cell lines	IC ₅₀ [μM] ^[a]
A549	47.8 ± 4.3
MCF-7	52.9 ± 3.8
–	–
PC-3M	32.0 ± 2.4
MDCK	29.3 ± 4.0
MDR-1-MDCK	73.2 ± 3.5

[a] IC₅₀ values for inhibition of human tumor cell growth; A549 (lung), MCF-7 (breast), [–], PC-3M (prostate), MDCK (canine kidney). MDR-1-MDCK is a human P-glycoprotein 170 (P-gp170)-overexpressing multi-drug-resistant cell line.