

A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie


50
YEARS

International Edition



www.angewandte.org

2011–50/20



Endocyclic P–P bonds ...

... are found in carbaborane-substituted 1,2-diphosphetanes, which are obtained in high yield from diastereomeric mixtures of 1,2-bis(chlorophosphanyl)-1,2-dicarba-*closo*-dodecaboranes(12). In their Communication on page 4701 ff., E. Hey-Hawkins and co-workers show that the air- and water-stable racemic 1,2-diphosphetane depicted undergoes a ring-opening reaction with elemental iodine to give exclusively *rac*-1,2-bis(iodo-*tert*-butylphosphanyl)-1,2-dicarba-*closo*-dodecaborane(12).

WILEY-VCH

Back Cover

**Anika Kreienbrink, Menyhárt B. Sárosi, Evgeny G. Rys,
Peter Lönnecke, and Evamarie Hey-Hawkins***

Endocyclic P–P bonds are found in carbaborane-substituted 1,2-diphosphetanes, which are obtained in high yield from diastereomeric mixtures of 1,2-bis(chlorophosphanyl)-1,2-dicarba-*closo*-dodecaboranes(12). In their Communication on page 4701 ff., E. Hey-Hawkins and co-workers show that the air- and water-stable racemic 1,2-diphosphetane depicted undergoes a ring-opening reaction with elemental iodine to give exclusively *rac*-1,2-bis(iodo-*tert*-butylphosphanyl)-1,2-dicarba-*closo*-dodecaborane(12).

