

A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie


50
YEARS

International Edition

GDCh

www.angewandte.org

2011–50/28



Peptide amphiphiles ...

... and oppositely charged polymers can self-assemble into highly organized membranes at the interface of two aqueous solutions. In their Communication on page 6324 ff., S. I. Stupp and co-workers report how the biopolymer was sprayed into a peptide amphiphile solution to template the formation of cell-like microcapsules. The image shows SEM micrographs of a group of filamentous microcapsules, a single capsule (blue), the surface nanofilaments (yellow), and a membrane cross section (sepia).

 WILEY-VCH

Back Cover

**Joshua E. Goldberger, Eric J. Berns, Ronit Bitton,
Christina J. Newcomb, and Samuel I. Stupp***

Peptide amphiphiles and oppositely charged polymers can self-assemble into highly organized membranes at the interface of two aqueous solutions. In their Communication on page 6324 ff., S. I. Stupp and co-workers report how the biopolymer was sprayed into a peptide amphiphile solution to template the formation of cell-like microcapsules. The image shows SEM micrographs of a group of filamentous microcapsules, a single capsule (blue), the surface nanofilaments (yellow), and a membrane cross section (sepia).

