
## A Journal of the Gesellschaft Deutscher Chemiker ANGELVANGE STEARS International Edition GDCh A Journal of the Gesellschaft Deutscher Chemiker Chemiker Chemiker Chemiker Chemiker Chemice www.angewandte.org



## Nonacene crystals ...

... possess a bright metallic sheen and their solution absorption spectra fit well in the series of silylethyne-substituted acenes. As J. E. Anthony et al. report in their Communication on page 7013 ff., crystallographic analysis of functionalized nonacenes confirmed the nine-ringed structure and showed that  $\pi\text{-stacked}$  arrangements exist in the solid state. EPR analysis showed that the nonacenes possess radical character, and exposure of nonacene solutions to light caused decomposition to form a red-fluorescent compound.



## **Inside Cover**

## Balaji Purushothaman, Matthew Bruzek, Sean R. Parkin, Anne-Frances Miller, and John E. Anthony\*

Nonacene crystals possess a bright metallic sheen and their solution absorption spectra fit well in the series of silylethyne-substituted acenes. As J. E. Anthony et al. report in their Communication on page 7013 ff., crystallographic analysis of functionalized nonacenes confirmed the nine-ringed structure and showed that  $\pi$ -stacked arrangements exist in the solid state. EPR analysis showed that the nonacenes possess radical character, and exposure of nonacene solutions to light caused decomposition to form a red-fluorescent compound.

