

A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie


50
YEARS

International Edition

GDCh

www.angewandte.org

2011–50/33



Multiple C–H bond activations ...

... give molecular lanthanoid-transition-metal clusters from heterobimetallic organometallic complexes. In their Communication on page 7695 ff. F. R. Wagner, R. Kempe, and co-workers show there is an increase in the delocalization of the metal-metal bonding electrons suggesting a possible transition sequence from the bimetallic complex to the intermetallic phase. (Background: photomontage (by Jana Kempe) of the Festspielhaus (Bayreuth) and the Semperoper (Dresden).)

 WILEY-VCH

Inside Cover

**Mikhail V. Butovskii, Oleg L. Tok, Viktor Bezugly, Frank R. Wagner,*
and Rhett Kempe***

Multiple C–H bond activations give molecular lanthanoid–transition-metal clusters from heterobimetallic organometallic complexes. In their Communication on page 7695 ff. F. R. Wagner, R. Kempe, and co-workers show there is an increase in the delocalization of the metal–metal bonding electrons suggesting a possible transition sequence from the bimetallic complex to the intermetallic phase. (Background: photomontage (by Jana Kempe) of the *Festspielhaus* (Bayreuth) and the Semperoper (Dresden).)

