

A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

50 YEARS


International Edition

GDCh

www.angewandte.org

2011–50/34

Organocatalysis



An organocatalytic ...

... highly stereoselective addition of benzodithiolium tetrafluoroborate to aldehydes tolerates a large range of functional groups. As P. G. Cozzi and co-workers report in the Communication on page 7842 ff., the reaction occurs through the formation of a key enamine intermediate in the presence of the MacMillan catalyst. The stability of the carbenium ion and the high nucleophilicity of the enamine, illustrated by their presumed value on the Mayr scale, enable the reaction to take place even in the presence of water.

 WILEY-VCH

Inside Cover

Andrea Gualandi, Enrico Emer, Montse Guiteras Capdevila, and Pier Giorgio Cozzi*

An organocatalytic highly stereoselective addition of benzodithiolium tetrafluoroborate to aldehydes tolerates a large range of functional groups. As P. G. Cozzi and co-workers report in the Communication on page 7842 ff., the reaction occurs through the formation of a key enamine intermediate in the presence of the MacMillan catalyst. The stability of the carbenium ion and the high nucleophilicity of the enamine, illustrated by their presumed value on the Mayr scale, enable the reaction to take place even in the presence of water.

