


Alkylation

Stereoselective Alkylation of Allylic Alcohols: Tandem Ethylation and Functionalization**

Pragna Pratic Das, Ivan L. Lysenko, and Jin Kun Cha*

Transition metal catalyzed allylic substitution is used frequently in organic synthesis. A high level of regio- and stereocontrol of asymmetric allylic alkylation is made possible by a judicious choice of substrates, late-transition-metal catalysts, and chiral ligands.^[1,2] Outside the domain of late transition metal catalyzed reactions the use of group IVA metals can also be effective for allylic substitution of readily available allylic alcohols.^[3] For example, treatment of acyclic and cyclic allylic alcohols with the ethyl Grignard reagent in the presence of titanium isopropoxide provides stereoselective S_N2'-type ethylation products [Eq. (1); EI = H].^[4–6] The



synthetic utility of the low-valent titanium (Kulinkovich reagent)^[4] mediated alkylation would be greatly enhanced by an in situ introduction of a functionalized side chain (EI ≠ H). We report herein a stereoselective alkylation reaction of allylic alcohols by trapping of the presumed alkyltitanium intermediates with suitable electrophiles.

This formal S_N2' ethylation reaction by the ethyl Grignard reagent was first reported by the Kulinkovich group,^[5] and the stereochemical studies revealed the overall suprafacial stereochemical outcome is due to *syn* addition of the Kulinkovich reagent through a Ti–O tether and subsequent *syn* β elimination.^[5d,e,6] Our mechanistic analysis implicated the generation of an alkyltitanium intermediate **4** by way of β elimination of **3** at the penultimate stage, which prompted us to investigate in situ elaboration of **4** to broaden the scope of this allylation reaction (Table 1).^[7] In light of the known instability of alkyl homologues of MeTi(OiPr)₃,^[8] the temperature control of the reaction was first examined, and it was optimal to warm the reaction mixture from –78 °C up to –10 or –5 °C, to set the stage of the trapping step. The presence of a scavenger for magnesium alkoxides in the reaction mixture was necessary prior to addition of an aldehyde to prevent an aldol condensation. Among three scavengers [TMSCl, TiCl₄,

Table 1: Titanium-mediated ethylation of acyclic allylic alcohols and in situ functionalization using three different scavengers.

Entry	5 (R ²)	TMSCl ^[b]	Yield [%] ^[a] TiCl ₄ ^[c]	CITi(OiPr) ₃ ^[d]
1	5a (iPr)	52	58	70
2	5b (Ph)	56	60	73
3	5c (PhCH=CH)	52	n.d.	n.d.
4	5d (CH ₂ =CH)	56	n.d.	n.d.
5	5e (CH ₂ =CMe)	48	n.d.	69
6	5f (MeCH=CH)	52	n.d.	73
7	5g (H)	45	n.d.	48

[a] Yields of the isolated products are given for each of the three scavengers used. [b] Used 3 equiv. [c] Used 0.75 equiv. [d] Used 3 equiv. n.d. = not determined.

and CITi(OiPr)₃] examined, CITi(OiPr)₃ afforded the most satisfactory yields of the products. The reaction conditions were found to be general for aliphatic, unsaturated, and aromatic aldehydes. Halogenation and oxidation of the alkyl–titanium bond of **4** were also effected by electrophilic halogenation reagents (I₂ and CF₃SO₂Cl) and O₂, respectively, to provide **6**, **7**, and **8** (Scheme 1).

As expected, cyclic substrates are also well suited for allylic ethylation and in situ functionalization (Table 2 and Scheme 2).

Products	EI	Yield [%]
6	I	69 ^[a] (59) ^[b]
7	Cl	72 ^[a] (52) ^[b]
8	OH	65 ^[a] (55) ^[b]

Scheme 1. Halogenation and oxidation of the alkyl–titanium bond of **4**.

[a] Yields of the isolated products when using 3 equiv CITi(OiPr)₃.

[b] Yields of the isolated products in absence of CITi(OiPr)₃.

[*] Dr. P. P. Das, Dr. I. L. Lysenko, Prof. Dr. J. K. Cha
 Department of Chemistry, Wayne State University
 Detroit, MI 48236 (USA)
 E-mail: jcha@chem.wayne.edu

[**] We thank the NSF (CHE-0615604) for generous financial support.
 Supporting information for this article is available on the WWW
 under <http://dx.doi.org/10.1002/anie.201104331>.

Table 2: Titanium-mediated ethylation of cyclic allylic alcohols and in situ functionalization.

Entry	10 (R ²)	TMSCl ^[b]	Yield [%] ^[a] ClTi(OiPr) ₃ ^[b]
1	10a (<i>i</i> Pr)	42	71
2	10b (PhCH=CH)	58	76
3	10c (CH ₂ =CH)	n.d.	70
4	10d (CH ₂ =CMe)	70	78
5	10e (H)	45	56


[a] Yields of the isolated products are given for each of the three scavengers used. [b] Used 3 equiv.

Products	EI	Yield [%]
11	I	62 ^[a] (52) ^[b]
12	OH	68 ^[a] (59) ^[b]


Scheme 2. Halogenation and oxidation of the alkyl–titanium to give products **11** and **12**, respectively. [a] Yields of the isolated products when using 3 equiv ClTi(OiPr)₃. [b] Yields of the isolated products in absence of ClTi(OiPr)₃.

The construction of all-carbon quaternary stereocenters in the context of natural product synthesis was next undertaken to showcase the utility of the in situ functionalization of the presumed alkyltitanium intermediates. The enantioselective formation of quaternary stereocenters has been an active area of research.^[9] Toward this end, **13** was selected as a target molecule, which had been converted by Danishefsky and co-workers into dysidiolide,^[10f] a bioactive sesterterpene (Scheme 3).^[10,11] Starting from the known and readily available (*S*)-seudenol (**14**),^[12] the stereoselective ethylation and subsequent in situ trapping with methacrolein, provided easy access to the alcohol **15** bearing a quaternary stereocenter in 78% yield. Acetylation of alcohol **15** with subsequent chemoselective epoxidation and Pd⁰-catalyzed allylic deoxygenation,^[13] afforded the epoxide **17** as an inconsequential mixture of two diastereomers. The epoxide was next converted into the ketone **18** by standard methods. The diene **13** was then prepared by known procedures.^[10b,e] As **13** had already been converted into dysidiolide,^[10b,f] this work represents its formal synthesis.

The enone **19** was selected as the second target as a prototype of 4,4-dialkyl-2-cyclohexenones (Scheme 4). There is a paucity of general methods for enantioselectively preparing 4,4-dialkyl-2-cyclohexenones, which stands in marked contrast to more readily accessible 4-alkyl-2-cyclohexenones.^[14] Additionally, treatment of **19** with EtAlCl₂ was documented to deliver an attractive route to bicyclic ketone **20**,^[15] which has found applications in natural product synthesis.^[16,17] Under our standard reaction conditions an S_N2'-type alkylation of the enantiomerically enriched **21a**^[18a] and **21b**^[18b] and subsequent trapping with acrolein gave **23a** and



Scheme 3. Formal synthesis of (–)-dysidiolide.



Scheme 4. Enantioselective synthesis of a 4,4-dialkyl-2-cyclohexenone. a) ClTi(OiPr)₃ (1 equiv), EtMgBr (3 equiv), Et₂O, –78 → –5 °C; then ClTi(OiPr)₃ (3 equiv); CH₂=CHCHO.

23b, respectively, in 68% yield and with little loss in the *ee* value. The *Z*-alcohol **22** was prepared from nerol for comparison, and its parallel reaction proceeded unexpectedly in poor (22%) yield. Ring-closing metathesis of **23a** or **23b** with Grubbs' second-generation catalyst, followed by Dess–Martin oxidation, provided the cyclohexenone **19** cleanly. Finally, a short enantioselective synthesis of **20** was achieved by the method reported by Snider et al.^[15]

In conclusion, a straightforward, yet valuable functionalization of the primary C–Ti bond with suitable electrophiles has been incorporated into the S_N2'-type reaction of acyclic

and cyclic allylic alcohols with the ethyl Grignard reagent in the presence of a titanium alkoxide. The stereospecific and convenient introduction of a functionalized side chain by the use of inexpensive reagents is notable and broadens the scope of this alkylation. Additionally, the present method offers direct use of allylic alcohols without preactivation or derivatization. This also represents formal cross-coupling reactions between allylic alcohols and terminal olefins bearing useful functional groups with concomitant 1,3-transposition.^[3,6]

Received: June 22, 2011

Published online: August 26, 2011

Keywords: alkylation · allylic compounds · natural products · synthetic methods · titanium

- [1] For reviews, see: a) B. M. Trost, M. L. Crawley, *Chem. Rev.* **2003**, *103*, 2921; b) D. K. Leahy, P. A. Evans in *Modern Rhodium-Catalyzed Organic Reactions* (Ed.: P. A. Evans), Wiley-VCH, Weinheim, **2005**, chap. 10, pp. 191–204; c) G. Helmchen, A. Dahnz, P. Dübon, M. Schelwies, R. Weihofen, *Chem. Commun.* **2007**, 675; d) Z. Lu, S. Ma, *Angew. Chem.* **2008**, *120*, 264; *Angew. Chem. Int. Ed.* **2008**, *47*, 258; e) S. R. Harutyunyan, T. den Hartog, K. Geurts, A. J. Minnaard, B. L. Feringa, *Chem. Rev.* **2008**, *108*, 2824.
- [2] For reviews of organocopper reagents, see: a) B. H. Lipshutz, S. Sengupta, *Org. React.* **1992**, *41*, 135; b) P. Knochel, N. Millot, A. L. Rodriguez, C. E. Tucker, *Org. React.* **2001**, 58, 417.
- [3] For titanacycle-mediated stereoselective cross-coupling reactions, see: a) F. Kolundzic, G. C. Micalizio, *J. Am. Chem. Soc.* **2007**, *129*, 15112; b) J. K. Belardi, G. C. Micalizio, *J. Am. Chem. Soc.* **2008**, *130*, 16870; c) H. A. Reichard, G. C. Micalizio, *Chem. Sci.* **2011**, *2*, 573.
- [4] For reviews of the Kulinkovich reaction, see: a) O. G. Kulinkovich, A. de Meijere, *Chem. Rev.* **2000**, *100*, 2789; b) O. G. Kulinkovich, *Chem. Rev.* **2003**, *103*, 2597; c) O. G. Kulinkovich, *Russ. Chem. Bull.* **2004**, *53*, 1065; d) A. Wolan, Y. Six, *Tetrahedron* **2010**, *66*, 3097.
- [5] a) O. G. Kulinkovich, O. L. Epstein, V. E. Isakov, E. A. Khmel'nitskaya, *Synlett* **2001**, 49; b) E. A. Matyushenkov, D. G. Churikov, N. A. Sokolov, O. G. Kulinkovich, *Russ. J. Org. Chem.* **2003**, *39*, 478; c) O. G. Kulinkovich, T. A. Shevchuk, V. E. Isakov, K. N. Prokhorevich, *Russ. J. Org. Chem.* **2006**, *42*, 659; d) V. E. Isakov, O. G. Kulinkovich, *Tetrahedron Lett.* **2008**, *49*, 6959; e) O. Kulinkovich, V. Isakov, D. Kananovich, *Chem. Rec.* **2008**, *8*, 269.
- [6] I. L. Lysenko, K. Kim, H. G. Lee, J. K. Cha, *J. Am. Chem. Soc.* **2008**, *130*, 15997.
- [7] Cf; a) J. Schwartz, J. A. Labinger, *Angew. Chem.* **1976**, *88*, 402; *Angew. Chem. Int. Ed. Engl.* **1976**, *15*, 333; b) F. Sato, H. Urabe in *Titanium and Zirconium in Organic Synthesis* (Ed.: I. Marek), Wiley-VCH, New York, **2002**, pp. 319–354.
- [8] M. T. Reetz in *Organometallics in Synthesis A Manual* (Ed: M. Schlosser), Wiley, New York, **2002**, pp. 817–923.
- [9] For reviews, see: a) E. J. Corey, A. Guzman-Perez, *Angew. Chem.* **1998**, *110*, 402; *Angew. Chem. Int. Ed.* **1998**, *37*, 388; b) I. Denissova, L. Barriault, *Tetrahedron* **2003**, *59*, 10105; c) *Quaternary Stereocentres, Challenges and Solutions for Organic Synthesis* (Eds.: J. Christoffers, A. Baro), Wiley-VCH, New York, **2005**, pp. 336; d) C. J. Douglas, L. E. Overman, *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 5363; e) J. Christoffers, A. Baro, *Adv. Synth. Catal.* **2005**, *347*, 1473; f) B. M. Trost, C. Jiang, *Synthesis* **2006**, 369; g) O. Riant, J. Hannedouche, *Org. Biomol. Chem.* **2007**, *5*, 873; h) M. Bella, T. Gasperi, *Synthesis* **2009**, 1583; i) C. Hawner, A. Alexakis, *Chem. Commun.* **2010**, 46, 7295; j) J. P. Das, I. Marek, *Chem. Commun.* **2011**, 47, 4593.
- [10] (–)-Dysidiolide: a) E. J. Corey, B. E. Roberts, *J. Am. Chem. Soc.* **1997**, *119*, 12425; b) M. Takahashi, K. Dodo, Y. Hashimoto, R. Shirai, *Tetrahedron Lett.* **2000**, *41*, 2111; c) H. Miyaoka, Y. Kajiwarra, Y. Hara, Y. Yamada, *J. Org. Chem.* **2001**, *66*, 1429; d) M. E. Jung, N. Nishimura, *Org. Lett.* **2001**, *3*, 2113; (+)-dysidiolide: e) J. Boukouvalas, Y.-X. Cheng, J. Robichaud, *J. Org. Chem.* **1998**, *63*, 228; (±)-dysidiolide: f) S. R. Magnuson, L. Sepp-Lorenzino, N. Rosen, S. J. Danishefsky, *J. Am. Chem. Soc.* **1998**, *120*, 1615; g) D. Demeke, C. J. Forsyth, *Org. Lett.* **2000**, *2*, 3177; h) R. Paczkowski, C. Maichle-Mössmer, M. E. Maier, *Org. Lett.* **2000**, *2*, 3967; i) K. P. Kaliappan, P. Gowrisankar, *Tetrahedron Lett.* **2004**, *45*, 8207.
- [11] D. Brohm, N. Philippe, S. Metzger, A. Bhargava, O. Müller, F. Lieb, H. Waldmann, *J. Am. Chem. Soc.* **2002**, *124*, 13171.
- [12] a) M. Kawasaki, Y. Suzuki, S. Terashima, *Chem. Pharm. Bull.* **1985**, *33*, 5; b) K.-W. Mu, W. H. Okamura, *J. Org. Chem.* **1990**, *55*, 4025; c) S. Hashiguchi, A. Fujii, K.-J. Haack, K. Matsumura, T. Ikariya, R. Noyori, *Angew. Chem.* **1997**, *109*, 300; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 288; d) R. ter Halle, Y. Bernet, S. Billard, C. Bufferne, P. Carlier, C. Delaitre, G. Flouzat, G. Humblot, J. C. Laigle, F. Lombard, S. Wilmouth, *Org. Process Res. Dev.* **2004**, *8*, 283.
- [13] a) J. Tsuji, T. Yamakawa, *Tetrahedron Lett.* **1979**, *7*, 613; b) M. D. Alexander, S. D. Fontaine, J. J. La Clair, A. G. DiPasquale, A. L. Rheingold, M. D. Burkart, *Chem. Commun.* **2006**, 4602.
- [14] a) Y. Chi, S. H. Gellman, *Org. Lett.* **2005**, *7*, 4253; b) K. Chen, P. S. Baran, *Nature* **2009**, *459*, 824.
- [15] B. B. Snider, D. J. Rodini, J. van Straten, *J. Am. Chem. Soc.* **1980**, *102*, 5872.
- [16] Although there was no asymmetric synthesis, resolution of a racemate was reported: L. A. Paquette, T. M. Heidelberg, *Synthesis* **1998**, 495.
- [17] K. Kim, J. K. Cha, *Angew. Chem.* **2009**, *121*, 5438; *Angew. Chem. Int. Ed.* **2009**, *48*, 5334.
- [18] a) Prepared by asymmetric ethylation of geranial in 90–92% ee by the method of Kobayashi: H. Takahashi, T. Kawakita, M. Ohno, M. Yoshioka, S. Kobayashi, *Tetrahedron* **1992**, *48*, 5691; b) Prepared by CBS reduction of the corresponding methyl ketone in 83–85% ee: E. J. Corey, C. J. Helal, *Angew. Chem.* **1998**, *110*, 2092; *Angew. Chem. Int. Ed.* **1998**, *37*, 1986; c) See the Supporting Information for the determination of the ee values of **15** (**13**) and **23** (**20**).