

Allylic Alkylation


Regio- and Stereocontrolled Introduction of Secondary Alkyl Groups to Electron-Deficient Arenes through Copper-Catalyzed Allylic Alkylation**

Yusuke Makida, Hirohisa Ohmiya,* and Masaya Sawamura*


Heteroarenes are important structural motifs found in many pharmaceuticals, agrochemicals and natural products.^[1] Accordingly, the development of efficient methods for heteroarene functionalization is important. Although there are various methods for accessing functionalized heteroarenes, the C-alkylation of these species is still severely limited in scope. Indeed, conventional methods involving stoichiometric metalation of electron-deficient heteroarenes followed by trapping with alkyl halides or pseudo halides are generally difficult because heteroarylmatal species are unstable compared with arylmetals.^[2] Friedel–Crafts-type reactions also allow for alkylation, but these methods are only applicable to electron-rich heteroarenes.^[3] Recently, transition-metal-catalyzed C_{sp²}–H functionalizations of (hetero)arenes, such as alkene hydro(hetero)arylations^[4,5] or couplings with alkyl halides,^[6,7] have been introduced as new approaches for heteroarene C-alkylation. However, even with these methods the introduction of secondary alkyl groups is quite difficult.^[5,8] In particular, the stereocontrolled introduction of a secondary alkyl group remains underdeveloped, although it was achieved in the intramolecular alkene hydroarylations of Ellman, Bergman and co-workers.^[9]

Earlier, we reported organoboron-based Pd^{II}- or Cu^I-catalyzed approaches for the allylic alkylation of arenes (Scheme 1 a).^[10] High γ -regioselectivity, 1,3-*anti* or *syn* stereoselectivities, and broad functional group compatibilities are all attractive features of these approaches. Therefore, the extension of these organoboron-based approaches to the alkylation of heteroarenes might be expected. We did not take this approach, however, because we knew that α -borylheteroarenes are generally unstable and difficult to prepare. Instead, we envisioned that the base-assisted direct cupration of heteroarenes under the conditions of Daugulis might be effective and more straightforward for the catalytic

a) γ -Selective Allylic Alkylation with Arylboron Compounds



b) γ -Selective Allylic Alkylation of Heteroarenes




Scheme 1. γ -Selective allyl–aryl or allyl–heteroaryl couplings. LG = leaving group.


generation of heteroarylcopper(I) species that are reactive in the γ -selective allyl–aryl coupling (Scheme 1 b).^[11]

Herein we report a Cu-catalyzed allylic alkylation of electron-deficient heteroarenes with internal secondary allylic phosphates, which proceeded with excellent γ -regioselectivity and *E*-stereoselectivity.^[12–15] This copper catalyst system was similarly applicable to fluoroarenes. Furthermore, the reaction of enantioenriched secondary allylic phosphates proceeded with 1,3-*anti* stereoselectivity to afford the corresponding alkylated (hetero)arenes with a controlled secondary stereogenic center. Thus, this Cu-catalyzed alkylation is a straightforward method for the stereocontrolled introduction of secondary alkyl groups to electron-deficient (hetero)arenes.

Specifically, the γ -substitution reaction of 4-phenyloxazole (**1a**; 0.6 mmol) with *Z* allylic phosphate **2a** (0.5 mmol) in the presence of CuCl (10 mol %) and LiOtBu (0.5 mmol) in THF (1 mL) at 40 °C for 10 h afforded alkylated arene product **3aa** in 78 % yield (87 % conversion) with excellent regio- (**3aa/3aa'** 99:1) and stereoselectivities (*E/Z* > 99:1) [Eq. (1)].^[16,17] On the other hand, the reaction of the isomeric



substrate **2a'** proceeded with slightly decreased γ -selectivity to afford **3aa** and **3aa'** in a 5:95 ratio [Eq. (2)]. The slight



[*] Y. Makida, Prof. Dr. H. Ohmiya, Prof. Dr. M. Sawamura
Department of Chemistry, Faculty of Science
Hokkaido University, Sapporo, 060-0810 (Japan)
E-mail: ohmiya@sci.hokudai.ac.jp
sawamura@sci.hokudai.ac.jp

Homepage: <http://barato.sci.hokudai.ac.jp/~orgmet/index.php>


[**] This work was supported by Grants-in-Aid for Scientific Research (B) and for Young Scientists (B), JSPS. We thank MEXT for financial support in the form of a Global COE grant (Project No. B01: Catalysis as the Basis for Innovation in Materials Science). Y.M. thanks JSPS for scholarship support.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201200809>.

difference in the regioselectivities of the isomeric substrates indicates that the relative steric demands of the α - and γ -substituents perturb the regioselectivity to some extent. However, these results indicate that a useful level of γ -selectivity is obtainable, even when unfavorable steric effects are present.

Several observations concerning the optimum reaction conditions for the reaction between **1a** and **2a'**, [Eq. (2)] are to be noted: No reaction occurred when KOtBu was used as a base or in the absence of CuCl. The use of CuOAc, CuOTf, or CuI instead of CuCl resulted in decreased yields and regioselectivities (70%, 12:88; 73%, 13:87; 71%, 23:77, respectively). Also, allylic substrates with carbonate or acetate leaving groups were not reactive at all.

The steric effects of the α - and γ -substituents of allylic phosphates **2** (Scheme 2) on the reactivity and regioselectivity



Scheme 2. Allylic phosphates used. TIPS = triisopropylsilyl, TBDMS = *tert*-butyldimethylsilyl.

were further evaluated and the results are shown in Table 1, entries 1–5.^[18] As the γ -substituent became bulkier (Me < Bu < *i*Bu), the regioselectivity gradually decreased (99 > 96 > 92 %) with decreasing product yields (81 > 73 > 63 %; entries 1–3). On the other hand, allylic phosphates **2e** and **2f**, bearing sterically more demanding groups (*i*Pr or cyclohexyl) in place of the α -PhCH₂CH₂ substituent of **2b**, underwent the reaction with excellent γ -selectivity (> 99:1; entries 4 and 5).

Cyclic allylic phosphates also served as substrates (Table 1, entries 6 and 7). Oxazole **1a** reacted with 2-cyclohexenyl phosphate (**2g**) to provide the product in a decent yield (53 %; entry 6). The reaction of **1a** with seven-membered cyclic phosphate **2h** was more effective, providing the product in 92 % yield (entry 7).

The Cu-catalyzed allylation was applicable to a range of heteroarenes (**1**) and allylic phosphates (**2**) as shown in Table 2. Functional groups on **1** or **2**, such as silyl ethers,

Table 1: Cu-catalyzed allylic alkylation with **1a**.^[a]

Entry	Arene	Phosphate	Product ^[b]	Yield [%] ^[c]	γ/α ^[d]
1	1a	2b		81	> 99:1
2	1a	2c		73	96:4
3	1a	2d		63	92:8
4 ^[e]	1a	2e		45	> 99:1
5 ^[f]	1a	2f		61	> 99:1
6	1a	2g		53	–
7	1a	2h		92	–

[a] Reaction conditions: **1a** (0.24 mmol), **2** (0.2 mmol), CuCl (10 mol %), LiOtBu (0.2 mmol) in THF (0.4 mL) at 40 °C for 10 h. [b] Isomeric ratio (*E/Z*) > 99:1. [c] Yield of isolated product. [d] Determined by ¹H NMR or GC analysis of the crude product. [e] The reaction was carried out at 60 °C. [f] The reaction was carried out on a 1.0 mmol scale at 60 °C.

esters, CF₃, MeO, and Cl, were also compatible (Table 2, entries 1–4, 6, and 9–11).

Various azole derivatives were used (Table 2).^[19] The reaction of 4-phenyloxazole derivative **1b** with a CF₃ substituent at the *para* position of the phenyl group occurred with a high γ -selectivity (97:3; entry 3). Substitution with a MeO group (**1c**), however, decreased the γ -selectivity to 88 % (entry 4). Benzoxazole derivatives **1d** and **1e** were effective in the reaction with **2a** and afforded **3da** and **3ea**, respectively, with 96 % γ -selectivity (entries 5 and 6). Ring substitution with a Ph group was also tolerated at the 5-position of the oxazole, as shown in the reaction of **1f** (entry 7). A sulfur-containing heterocycle, 4-methylthiazole (**1g**), reacted in moderate yield (60 %) with excellent γ -regioselectivity (> 99:1; entry 8). The reaction between 2-(4-chlorophenyl)-1,3,4-oxadiazole (**1h**) and **2a** afforded **3ha** in 83 % yield with 95 % γ -regioselectivity (entry 9). The oxadiazole **1h** reacted with **2b** in 96 % γ -selectivity (entry 10).

The method was also applicable to a pyridine *N*-oxide derivative (entries 11 and 12). The reaction of 2-phenylpyridine *N*-oxide (**1i**) with **2b** occurred at the 6-position of the pyridine to afford the corresponding product **3ib** in high yield (92 %) with excellent γ -selectivity (> 99:1; entry 11). Replacing the γ -Me substituent of **2b** with a sterically more demanding butyl group (**2c**) did not affect the excellent γ -selectivity (> 99:1; entry 12).

The copper-catalyzed method has also been extended to the reaction of fluoroarenes (Table 3). Pentafluorobenzene (**1j**) reacted efficiently with **2b** to afford **3jb** in 93 % yield with 90 % γ -selectivity (entry 1). The reaction of **1j** with the α -isopropyl allylic phosphate **2e** showed excellent regioselectivity (87 % yield, γ/α > 99:1; entry 2). The cyclic allylic

Table 2: Cu-catalyzed allylic alkylation with various heteroarenes.^[a]

Entry	Arene	Phosphate	Product ^[b]	Yield [%] ^[c]	γ/α ^[d]
1	1a	2i	3ai	61	> 99:1
2	1a	2j	3aj	80	> 99:1
3	1b	2b	3bb	82	97:3
4	1c	2b	3cb	55	88:12
5	1d	2a	3da	77	96:4
6	1e	2a	3ea	50	96:4
7	1f	2b	3fb	52 ^[f]	> 99:1
8 ^[e]	1g	2b	3gb	60	> 99:1
9	1h	2a	3ha	83	95:5
10	1h	2b	3hb	86	96:4
11	1i	2b	3ib	92	> 99:1
12	1i	2c	3ic	60	> 99:1

[a] Reaction conditions: **1** (0.24 mmol), **2** (0.2 mmol), CuCl (10 mol %), LiOtBu (0.2 mmol) in THF (0.4 mL) at 40 °C for 10 h. [b] Isomeric ratio (*E/Z*) > 99:1. [c] Yield of isolated product. [d] Determined by ¹H NMR or GC analysis of the crude product. [e] The reaction was carried out at 60 °C. [f] Yield determined by NMR analysis. Significant amounts of unidentified compounds were also detected.

phosphate **2g** coupled with **1j** to give the product in good yield (entry 3). The tetrafluoroarenes **1k** and **1l**, bearing CF₃ and MeO groups, respectively, and 2,3,5,6-tetrafluoropyridine (**1m**) also served as suitable fluoroarene substrates (entries 4–6).

The alkylation of azole derivatives with enantioenriched allylic phosphates proceeded with 1,3-*anti* stereochemistry, allowing the stereocontrolled introduction of secondary alkyl groups (Table 4, entries 1–3). Specifically, the alkylation of 4-phenyloxazole (**1a**) with (*S*)-(Z)-**2b** (98% *ee*), which has α -PhCH₂CH₂ and γ -Me substituents, proceeded with 90% *anti* stereoselectivity, affording (*R*)-(E)-**3ab** with 79% *ee* (entry 1). More efficient chirality transfer occurred in the reaction of **1a** with (*S*)-(Z)-**2a'** (96% *ee*), which has α -Me and γ -Bu substituents, giving (–)-(E)-**3aa'** with 85% *ee* (*antisyn* 94:6; entry 2).^[20,21] The reaction of the oxadiazole **1h** with (*S*)-(Z)-**2a'** (96% *ee*) afforded (–)-(E)-**3ha'** with 81% *ee* (*anti*/

Table 3: Cu-catalyzed allylic alkylation with various fluoroarenes.^[a]

Entry	Arene	Phosphate	Product ^[b]	Yield [%] ^[c]	γ/α ^[d]
1	1j	2b	3jb	93	90:10
2	1j	2e	3je	87	> 99:1
3	1j	2g	3jg	78	–
4	1k	2b	3kb	88	94:6
5	1l	2b	3lb	67	92:8
6 ^[e]	1m	2e	3me	73	> 99:1

[a] Reaction conditions: **1** (0.24 mmol), **2** (0.2 mmol), CuCl (10 mol %), LiOtBu (0.2 mmol) in THF (0.4 mL) at 40 °C for 10 h. [b] Isomeric ratio (*E/Z*) > 99:1. [c] Yield of isolated product. [d] Determined by ¹H NMR or GC analysis of the crude product. [e] The reaction was carried out in toluene.

syn 92:8; entry 3). These results demonstrate that, despite some erosion of enantiomeric purity, the copper-catalyzed allylic alkylation allows the construction of a stereogenic center at the position α to electron-deficient heteroaromatic rings, which is rare and difficult to obtain by other methods.^[9]

As shown in Table 4, entries 4–7, the reaction of fluoroarenes with enantioenriched allylic phosphates also proceeded with 1,3-*anti* stereochemistry. Specifically, the reactions of pentafluorobenzene (**1j**) or 2,3,5,6-tetrafluoropyridine (**1m**) with enantioenriched (*S*)-(Z)-**2e** (99% *ee*), which has α -iPr and γ -Me groups gave enantioenriched (*R*)-(E)-**3je** (99% *ee*) and (–)-(E)-**3me** (95% *ee*) with 1,3-*anti* stereochemistry (*antisyn* > 98:2), respectively (Table 2, entries 4 and 5).^[20] The reaction of the fluoroarene **1k** with (*S*)-(Z)-**2b** (98% *ee*) also gave excellent chirality transfer (92% *ee*, *antisyn* 97:3; entry 6). The reaction of pentafluorobenzene (**1j**) with (*S*)-(Z)-**2k** (95% *ee*), which has a siloxy-methyl group at the γ -position, afforded (+)-(E)-**3jk** with 90% *ee* (*antisyn* 97:3; entry 7).

The active organocopper species is likely a monoorgano-heterocuprate ([ArCuOtBu][–]) rather than a neutral organo-copper(I) species (L_nCuAr), because LiOtBu is not basic enough to lithiate the (hetero)arenes and LiOtBu is present in an excess relative to copper during the catalytic reaction.^[22] This assumption is also supported by the fact that the γ -regioselectivities were generally high, but not always perfect. This rationale comes from our previous observation that the Cu-catalyzed allyl-alkyl coupling between allylic phosphates and alkylboranes occurred with perfect γ -selec-

Table 4: Chirality transfer.^[a]

Entry	Arene	Phosphate	Product ^[b]	Yield [%] ^[c]	γ/α ^[d]	<i>ee</i> [%] ^[e] (<i>anti/syn</i>)
1	1a	(<i>S</i>)-(<i>Z</i>)- 2b 98% <i>ee</i>		40	> 99:1	79 (90:10)
2	1a	(<i>S</i>)-(<i>Z</i>)- 2a' 96% <i>ee</i>		58	95:5	85 (94:6)
3	1h	(<i>S</i>)-(<i>Z</i>)- 2a' 96% <i>ee</i>		87	88:12	81 (92:8)
4	1j	(<i>S</i>)-(<i>Z</i>)- 2e 99% <i>ee</i>		87	> 99:1	99 (>99:1)
5	1m	(<i>S</i>)-(<i>Z</i>)- 2e 99% <i>ee</i>		73	> 99:1	95 (98:2)
6	1k	(<i>S</i>)-(<i>Z</i>)- 2b 98% <i>ee</i>		88	94:6	92 (97:3)
7	1j	(<i>S</i>)-(<i>Z</i>)- 2k 95% <i>ee</i>		91	> 99:1	90 (97:3)

[a] The reaction was carried out with **1** (0.24 mmol), **2** (0.2 mmol), CuCl (10 mol %), LiOtBu (0.2 mmol) in toluene (entries 1–3 and 5; 0.4 mL) or THF (entries 4, 6, and 7; 0.4 mL) at 40°C for 10 h. [b] Isomeric ratio (*E/Z*) > 99:1. [c] Yield of isolated product. [d] Determined by ¹H NMR or GC analysis of the crude product. [e] The *ee* values were determined by chiral-phase HPLC analysis.

([ArCuOtBu][−], **A**). Subsequently, **A** forms diastereomeric π -complexes **B** (path a) or **B'** (path b) with an allylic phosphate (**2**), in which the copper atom is *anti* to the leaving group (OP). Owing to the difference in the strength of the *trans* effect, isomer **B** is more reactive than **B'**. Thus, the oxidative addition of **B** through transition state **C**(TS) is predominant. This oxidative addition affords (σ -enyl)copper(III) species **D** (enyl[$\sigma+\pi$] complex), which has a C ^{γ} –Cu σ bond. Finally, reductive elimination of **D** results in forming the γ -coupling product **3- γ** . The occasional incomplete γ -selectivity suggests the minor involvement of path b, in which the diastereomeric π -complex **B'** leads to enyl[$\sigma+\pi$] complex **D'**, which forms the α -coupling product **3- α** .

In summary, we have developed a Cu-catalyzed allylic alkylation reaction of azoles (oxazoles, an oxadiazole, and a thiazole), a pyridine *N*-oxide derivative, and fluoroarenes with internal secondary allylic phosphates that proceeds under mild reaction conditions with excellent γ and *E*-selectivities. The reaction with enantioenriched allylic phosphates proceeded with 1,3-*anti* stereoselectivity to generate an allylic stereogenic center at the position α to the aromatic ring. Accordingly, the Cu-catalyzed reaction is a straightforward and powerful method for the stereocontrolled introduction of secondary alkyl groups to electron-deficient arenes.


Received: January 30, 2012

Published online: March 13, 2012

Keywords: allylic alkylation · azoles · copper · heteroarenes · regioselectivity

tivity, in which neutral alkylcopper(I) species were reasonable active intermediates.^[10f]

According to DFT studies by Nakamura et al. on the mechanism of the reaction between monomethylheterocuprates (Li[MeCuX]) and allyl acetate,^[23] we propose the mechanism shown in Scheme 3 for the present Cu-catalyzed reaction. The catalytic cycle is initiated by a LiOtBu-assisted C–H cupration of a (hetero)arene to produce a heterocuprate


Scheme 3. Possible mechanism.

- [1] a) T. Eicher, S. Hauptmann, *The Chemistry of Heterocycles*, Wiley-VCH, Weinheim, **2003**; b) V. S. C. Yeh, *Tetrahedron* **2004**, *60*, 11995–12042; c) Z. Jin, *Nat. Prod. Rep.* **2011**, *28*, 1143.
- [2] a) R. Chinchilla, C. Najera, M. Yus, *Chem. Rev.* **2004**, *104*, 2667–2722; b) A. Krasovskiy, V. Krasovskaya, P. Knochel, *Angew. Chem.* **2006**, *118*, 3024–3027; *Angew. Chem. Int. Ed.* **2006**, *45*, 2958–2961, and references therein.
- [3] a) M. Bandini, E. Emer, S. Tommasi, A. Umani-Ronchi, *Eur. J. Org. Chem.* **2006**, 3527–3544; b) T. B. Poulsen, K. A. Jorgensen, *Chem. Rev.* **2008**, *108*, 2903–2915; c) M. Bandini, A. Eichholzer, *Angew. Chem.* **2009**, *121*, 9786–9824; *Angew. Chem. Int. Ed.* **2009**, *48*, 9608–9644.
- [4] For recent reviews on C_{sp²}–H alkylations through transition-metal-catalyzed hydro(hetero)arylations of alkenes, see: a) J. C. Lewis, R. G. Bergman, J. A. Ellman, *Acc. Chem. Res.* **2008**, *41*, 1013–1025; b) F. Kakiuchi, T. Kochi, *Synthesis* **2008**, 3013–3039; c) D. A. Colby, R. G. Bergman, J. A. Ellman, *Chem. Rev.* **2010**, *110*, 624–655.
- [5] For the secondary benzylation of (hetero)arenes by transition-metal-catalyzed (with Ir, Ni, or Co) hydroarylations of vinylarenes, see: a) Y. Uchamaru, *Chem. Commun.* **1999**, 1133–1134; b) Y. Nakao, N. Kashiwara, K. S. Kanyiva, T. Hiyama, *J. Am. Chem. Soc.* **2008**, *130*, 16170–16171; c) Y. Nakao, N. Kashiwara, K. S. Kanyiva, T. Hiyama, *Angew. Chem.* **2010**, *122*, 4553–4556; *Angew. Chem. Int. Ed.* **2010**, *49*, 4451–4454; d) Y. Nakao, Y. Yamada, N. Kashiwara, T. Hiyama, *J. Am. Chem. Soc.* **2010**, *132*,

- 13666–13668; e) T. Mukai, K. Hirano, T. Satoh, M. Miura, *J. Org. Chem.* **2009**, *74*, 6410–6413; f) K. Gao, N. Yoshikai, *J. Am. Chem. Soc.* **2011**, *133*, 400–402; For the secondary alkylation of phenols by the Re-catalyzed hydroarylation of alkenes, see: g) Y. Kuninobu, T. Matsuki, K. Takai, *J. Am. Chem. Soc.* **2009**, *131*, 9914–9915.
- [6] For a review on transition-metal-catalyzed C_{sp}²–H alkylations of (hetero)arenes with alkyl halides, see: L. Ackermann, *Chem. Commun.* **2010**, 46, 4866–4877.
- [7] For transition-metal-catalyzed (with Ni, Pd, or Co) C_{sp}²–H alkylations of (hetero)arenes with primary alkyl halides, see: a) O. Vechorkin, V. Proust, X. Hu, *Angew. Chem.* **2010**, *122*, 3125–3128; *Angew. Chem. Int. Ed.* **2010**, *49*, 3061–3064; b) T. Yao, K. Hirano, T. Satoh, M. Miura, *Chem. Eur. J.* **2010**, *16*, 12307–12311; c) Q. Chen, L. Ilies, E. Nakamura, *J. Am. Chem. Soc.* **2011**, *133*, 428–429; d) Q. Chen, L. Ilies, N. Yoshikai, E. Nakamura, *Org. Lett.* **2011**, *13*, 3232–3234; e) L. Ackermann, B. Punji, W. Song, *Adv. Synth. Catal.* **2011**, *353*, 3325–3329; for Ru-catalyzed C_{sp}²–H alkylations of arenes with primary or secondary alkyl halides, see: f) L. Ackermann, P. Novák, R. Vicente, N. Hofmann, *Angew. Chem.* **2009**, *121*, 6161–6164; *Angew. Chem. Int. Ed.* **2009**, *48*, 6045–6048; for transition-metal-catalyzed (with Pd or Ru) benzylations of (hetero)arenes with primary benzylic electrophiles, see: g) T. Mukai, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2010**, *12*, 1360–1363; h) D. Lapointe, K. Fagnou, *Org. Lett.* **2009**, *11*, 4160–4163; i) L. Ackermann, P. Novák, *Org. Lett.* **2009**, *11*, 4966–4969; j) C. Verrier, C. Hoarau, F. Marsais, *Org. Biomol. Chem.* **2009**, *7*, 647–650; k) L. Ackermann, S. Barfüsser, J. Pospech, *Org. Lett.* **2010**, *12*, 724–726; l) S. Fan, C.-Y. He, X. Zhang, *Chem. Commun.* **2010**, 46, 4926–4928; m) L. Ackermann, N. Hofmann, R. Vicente, *Org. Lett.* **2011**, *13*, 1875–1877; n) L. Ackermann, S. Barfüsser, C. Kornhaas, A. R. Kapdi, *Org. Lett.* **2011**, *13*, 3082–3085; for the synthesis of heteroaryloxetanes by a radical reaction, see: o) M. A. J. Duncton, M. A. Estiarte, R. J. Johnson, M. Cox, D. J. R. O'Mahony, W. T. Edwards, M. G. Kelly, *J. Org. Chem.* **2009**, *74*, 6354–6357.
- [8] Wang and co-workers achieved secondary benzylation and allylation of 1,3-azoles through a Cu-catalyzed reaction with *N*-tosylhydrazones: a) X. Zhao, G. Wu, Y. Zhang, J. Wang, *J. Am. Chem. Soc.* **2011**, *133*, 3296–3299; Hirano, Miura and co-workers reported secondary alkylation of 1,3-azoles through Co- and Ni-catalyzed reactions with *N*-tosylhydrazones: b) T. Yao, K. Hirano, T. Satoh, M. Miura, *Angew. Chem.* **2012**, *124*, 799–803; *Angew. Chem. Int. Ed.* **2012**, *51*, 775–779.
- [9] a) S. J. O'Malley, K. L. Tan, A. Watzke, R. G. Bergman, J. A. Ellman, *J. Am. Chem. Soc.* **2005**, *127*, 13496–13497; b) J. C. Rech, M. Yato, D. Duckett, B. Ember, P. V. LoGrasso, R. G. Bergman, J. A. Ellman, *J. Am. Chem. Soc.* **2007**, *129*, 490–491; c) A. Watzke, R. M. Wilson, S. J. O'Malley, R. G. Bergman, J. A. Ellman, *Synlett* **2007**, 2383–2389; d) H. Harada, R. K. Thalji, R. G. Bergman, J. A. Ellman, *J. Org. Chem.* **2008**, *73*, 6772–6779; see also Refs [4a,c].
- [10] For Pd- and Cu-catalyzed γ -selective and stereospecific allyl-aryl coupling with arylboron compounds from our group, see: a) H. Ohmiya, N. Yokokawa, M. Sawamura, *Org. Lett.* **2010**, *12*, 2438–2440; b) H. Ohmiya, Y. Makida, T. Tanaka, M. Sawamura, *J. Am. Chem. Soc.* **2008**, *130*, 17276–17277; c) H. Ohmiya, Y. Makida, D. Li, M. Tanabe, M. Sawamura, *J. Am. Chem. Soc.* **2010**, *132*, 879–889; d) D. Li, T. Tanaka, H. Ohmiya, M. Sawamura, *Org. Lett.* **2010**, *12*, 2438–2440; e) Y. Makida, H. Ohmiya, M. Sawamura, *Chem. Asian J.* **2011**, *6*, 410–414; for related studies, see also: f) H. Ohmiya, U. Yokobori, Y. Makida, M. Sawamura, *J. Am. Chem. Soc.* **2010**, *132*, 2895–2897; g) D. Li, H. Ohmiya, M. Sawamura, *J. Am. Chem. Soc.* **2011**, *133*, 5672–5675.
- [11] a) H.-Q. Do, O. Daugulis, *J. Am. Chem. Soc.* **2007**, *129*, 12404–12405; b) H.-Q. Do, O. Daugulis, *J. Am. Chem. Soc.* **2008**, *130*, 1128–1129; c) H.-Q. Do, R. M. Kashif Khan, O. Daugulis, *J. Am. Chem. Soc.* **2008**, *130*, 15185–15192; d) F. Besselièvre, S. Piguel, *Angew. Chem.* **2009**, *121*, 9717–9720; *Angew. Chem. Int. Ed.* **2009**, *48*, 9553–9556; e) T. Kawano, K. Hirano, T. Satoh, M. Miura, *J. Am. Chem. Soc.* **2010**, *132*, 6900–6901; for reviews, see: f) O. Daugulis, H.-Q. Do, D. Shabashov, *Acc. Chem. Res.* **2009**, *42*, 1074–1086; g) A. A. Kulkarni, O. Daugulis, *Synthesis* **2009**, 4087–4109.
- [12] Recently, Hirano, Miura, and co-workers reported the Cu-catalyzed C_{sp}²–H allylation of fluoroarenes with allylic phosphates. The introduction of secondary alkyl groups to fluoroarenes was specifically described for the case of 2-cycloheptenyl phosphate. Under their conditions, the reaction of an asymmetrically 1,3-disubstituted secondary allylic phosphate with pentafluorobenzene was not regiocontrolled: T. Yao, K. Hirano, T. Satoh, M. Miura, *Angew. Chem.* **2011**, *123*, 3046–3050; *Angew. Chem. Int. Ed.* **2011**, *50*, 2990–2994.
- [13] Recently, Zhang and co-workers reported a Pd and Cu co-catalyzed method for C_{sp}²–H primary allylation of fluoroarenes with allylic carbonates. A secondary alkylation product was described only as a minor by-product in the formation of a primary allylation product: S. Fan, F. Chen, X. Zhang, *Angew. Chem.* **2011**, *123*, 6040–6045; *Angew. Chem. Int. Ed.* **2011**, *50*, 5918–5923.
- [14] For reviews on Cu-catalyzed allylic substitutions, see: a) B. Breit, P. Demel in *Modern Organocopper Chemistry* (Ed.: N. Krause), Wiley-VCH, Weinheim, **2002**, pp. 188–223; b) C. A. Falcicola, A. Alexakis, *Eur. J. Org. Chem.* **2008**, 3765–3780; c) A. Alexakis, J. E. Bäckvall, N. Krause, O. Pàmies, M. Diéguez, *Chem. Rev.* **2008**, *108*, 2796–2823; d) S. R. Harutyunyan, T. den Hartog, K. Geurts, A. J. Minnaard, B. L. Feringa, *Chem. Rev.* **2008**, *108*, 2824–2852.
- [15] For Cu-catalyzed enantioselective allylic substitution reactions with arylmetal reagents, see: a) M. A. Kacprzynski, T. L. May, S. A. Kazane, A. H. Hoveyda, *Angew. Chem.* **2007**, *119*, 4638–4642; *Angew. Chem. Int. Ed.* **2007**, *46*, 4554–4558; b) K. B. Selim, K. Yamada, K. Tomioka, *Chem. Commun.* **2008**, 5140–5142; c) K. B. Selim, Y. Matsumoto, K. Yamada, K. Tomioka, *Angew. Chem.* **2009**, *121*, 8889–8891; *Angew. Chem. Int. Ed.* **2009**, *48*, 8733–8735; d) C. A. Falcicola, A. Alexakis, *Chem. Eur. J.* **2008**, *14*, 10615–10627; e) D. Polet, X. Rathgeb, C. A. Falcicola, J. B. Langlois, S. E. Hajjaji, A. Alexakis, *Chem. Eur. J.* **2009**, *15*, 1205–1206; f) F. Gao, Y. Lee, K. Mandai, A. H. Hoveyda, *Angew. Chem.* **2010**, *122*, 8548–8552; *Angew. Chem. Int. Ed.* **2010**, *49*, 8370–8374.
- [16] Allylic alcohols with a *Z* configuration are readily obtainable though *Z*-selective reductions of the corresponding propargylic alcohols.
- [17] To our surprise, the *E*-isomer of **2a** showed no reactivity under the same conditions. Partial alcoholysis of **2a** to the corresponding allylic alcohol was observed.
- [18] The reaction between **1a** and a *Z* allylic phosphate bearing an aryl group at the γ -position resulted in no reaction. *Z* Allylic phosphates bearing an aryl group at the α -position were too unstable to prepare.
- [19] Reactions with imidazole derivatives were unsuccessful.
- [20] See the Supporting Information for experiments to determine the absolute configurations of (*R*)-(*E*)-**3ab** and (*R*)-(*E*)-**3je**.
- [21] Evaluation of the product enantiomeric excess values for the reactions of azole **1a** with different reaction times and conversions confirmed that no racemization of (*R*)-(*E*)-**3ab** occurred under the reaction conditions. This suggests moderate diastereotopic face selectivity in the reaction between the heteroaryl(copper(I) species and the allylic phosphate. Studies

for improving the efficiency of 1,3-chirality transfer in reactions with heteroarenes are under way.

- [22] Treatment of **1a** with LiOtBu (1:1) in THF at 40 °C for 1 h or 10 h followed by quenching with D₂O resulted in no deuterated product. These results indicate that the heteroaryllithium does not form in a meaningful amount in the Cu-catalyzed allylic alkylation. For a discussion of the LiOtBu-assisted direct
- cupration of electron-deficient arenes, see: Ref. [11], and references therein.
- [23] a) N. Yoshikai, S.-L. Zhang, E. Nakamura, *J. Am. Chem. Soc.* **2008**, *130*, 12862–12863; b) M. Yamanaka, S. Kato, E. Nakamura, *J. Am. Chem. Soc.* **2004**, *126*, 6287–6293; for a review, see: c) N. Yoshikai, E. Nakamura, *Chem. Rev.* **2012**, DOI: 10.1021/cr200241f.