


Natural Products


Biomimetic Total Synthesis of (+)-Gelsemine**

Xuan Zhou, Tao Xiao, Yusuke Iwama, and Yong Qin*

(+)-Gelsemine, a major alkaloid component of *Gelsemium sempervirens*, was first isolated in 1876.^[1] Its intriguing structure was elucidated in 1959 by NMR spectroscopy^[2] and X-ray crystallographic analysis.^[3] To date, seven members of the gelsemium alkaloid family have been characterized (1–7).^[4] These oxindole alkaloids have a hexacyclic architecture



(+)-gelsemine (1), R¹ = H, R² = H
(+)-gelsevirine (2), R¹ = OMe, R² = H, H
(+)-21-oxogelsemine (3), R¹ = H, R² = O
(+)-21-oxogelsevirine (4), R¹ = OMe, R² = O




(+)-19-hydroxydihydrogelsemine (5)
R¹ = H, R² = OH
(+)-19-hydroxydihydrogelsevirine (6)
R¹ = OMe, R² = OH
(+)-19-acetoxydihydrogelsevirine (7)
R¹ = OMe, R² = OAc

and seven carbon stereocenters, including two quaternary stereocenters compacted into a small cage. The challenging structure of gelsemine (1) has been the target of extensive synthetic efforts around the world;^[5] within the last two decades a number of total syntheses were reported by the research groups of Johnson, Speckamp, Fukuyama, Hart, Overman, and Danishefsky.^[6] Among these total syntheses, there was only one asymmetric total synthesis accomplished by the Fukuyama group.^[6g] Furthermore, none of these total syntheses has sufficiently illustrated the possible biosynthetic pathway of the gelsemine family. Herein, we report a biomimetic total synthesis of (+)-gelsemine, in which the key

reaction involves an intramolecular enol–oxonium cyclization that assembles the cage structure in a highly efficient way. Besides, a new biosynthetic pathway for the late stage of the biosynthesis of gelsemine is proposed based on the achieved enol–oxonium cyclization procedure.^[7]

The retrosynthetic analysis of (+)-gelsemine is shown in Scheme 1. We speculated that the cage structure of (+)-gelsemine could be assembled from an intermediate such as 8 in



Scheme 1. Retrosynthetic analysis of (+)-gelsemine. TBDMS = *tert*-butyldimethylsilyl, MOM = methoxymethyl, Ts = toluene-4-sulfonyl.

a single step by acid-catalyzed enol–oxonium cyclization to form, simultaneously, the E ring, the F ring, the C7 quaternary stereocenter of spirooxindole, and the C3-stereocenter. We reasoned that the D ring could be constructed by intramolecular Michael addition of highly functionalized nitrile 9, which can be readily prepared by intermolecular aldol addition of oxindole 10 to *cis* tetra-substituted piperidinoaldehyde 11, followed by dehydration. The 15*S* configuration in 11 could be generated by stereoselective formation of the C ring by an intramolecular Michael addition of oxepinone 12. Partial hydrogenation of alkynyl acid 13 could yield the *cis* double bond in 12. The *S* configuration at C16 in 13 could be produced by regio- and stereoselective ring opening of aziridine (*R,R*)-15 induced by the hydroxy group. The stereochemistry in (*R,R*)-15 could be generated from the known ester (*R,R*)-16,^[8] which is easily prepared from D-diethyl tartrate in a four-step procedure.


Our biomimetic total synthesis of (+)-gelsemine began by converting (*R,R*)-16 into the chiral *cis* tetra-substituted piperidinoaldehyde 11 (Scheme 2). Two steps of reduction

[*] X. Zhou, T. Xiao, Prof. Dr. Y. Qin
Key Laboratory of Drug Targeting and Drug Delivery Systems of the Ministry of Education, and State Key Laboratory of Biotherapy
Sichuan University
Chengdu, 610041 (China)
E-mail: yongqin@scu.edu.cn

Y. Iwama
Graduate School of Pharmaceutical Sciences, Tohoku University
Sendai, 980-8578 (Japan)
Prof. Dr. Y. Qin
Innovative Drug Research Centre and College of Chemistry and
Chemical Engineering, Chongqing University
Chongqing, 401331 (China)

[**] We are grateful for financial supports from NSFC (20825207, 21021001, and 21132006), the National Basic Research Program of China (973 program, 2010CB833200), the State Key Laboratory of Bioorganic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, and Tohoku University Global COE Program and JSPS Predoctoral Fellowship for Y.I.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201201736>.



Scheme 2. Preparation of the chiral *cis* tetra-substituted piperidine ring. Reagents and conditions: a) LiAlH_4 , EtOH, $0^\circ\text{C} \rightarrow 25^\circ\text{C}$, 3 h, 85%; b) TsCl , Et_3N , CH_2Cl_2 , $0^\circ\text{C} \rightarrow 25^\circ\text{C}$, 3 h, 90%; c) (3,3-diethoxyprop-1-yn-1-yl)lithium **14**, THF, $0^\circ\text{C} \rightarrow 25^\circ\text{C}$, 5 h, 95%; d) MeI , K_2CO_3 , DMF, 25°C , 98%; e) MOMCl , NaH , THF, 0°C , 2 h, 95%; f) Mg , MeOH, ultrasound, 0°C , 3 h, 90%; g) acrylonitrile, MeOH, reflux, 10 h, 95%; h) TFA, CH_2Cl_2 , -10°C , 3 h, 85%; i) NaClO_2 , NaH_2PO_3 , $\text{tBuOH}/\text{H}_2\text{O}$ 1:1, 25°C , 3 h, 94%; j) Lindlar cat, H_2 , EtOH, 25°C , 10 h, 95%; k) HF, pyridine, $0^\circ\text{C} \rightarrow 25^\circ\text{C}$, 10 h, 90%; l) DCC, DMAP, CH_2Cl_2 , 25°C , 10 h, 85%; m) LiHMDS , THF, -78°C , 10 min, 66%; n) DIBAL-H, CH_2Cl_2 , -78°C , 30 min, 60%; o) HC(OMe)_3 , $p\text{-TsOH}$, MeOH, $0^\circ\text{C} \rightarrow 25^\circ\text{C}$, 2 h, 80%; p) $(\text{COCl})_2$, DMSO, CH_2Cl_2 , -60°C , 1 h, 92%. TFA = trifluoroacetic acid, DCC = dicyclohexylcarbodiimide, DMAP = 4-(dimethylamino)pyridine, HMDS = 1,1,1,3,3,3-hexamethyldisilazane, DIBAL-H = diisobutylaluminum hydride, DMSO = dimethylsulfoxide.


and tosylation of **16** provided alcohol (*R,R*)-**15** in moderate yield. The attack by **14** on **15** occurred from the top face of the aziridine ring to afford **17** as a single stereomer in 95% yield. The excellent regio- and stereoselectivity is probably due to coordination of the lithium reagent **14** with the hydroxy group and efficient blockage of the bottom face of the aziridine ring by the bulky OTBDMS group in **15**.^[9] Two steps of methylation with MeI and protection of the hydroxy group in **17** provided compound **18** in excellent yield. After removal of the tosyl group in **18** with Mg dust using ultrasound,^[10] the resulting amine reacted with acrylonitrile in MeOH at reflux to give **19** in excellent yield. Treatment of **19** with TFA at -10°C , and subsequent oxidation of the resulting aldehyde group with sodium chlorite yielded acid **13** in 85% yield in two steps. Partial hydrogenation of the alkynyl group with Lindlar catalyst and removal of the TBDMS group with HF/pyridine in **13** provided *cis* olefin **20** in excellent yield. To prepare the piperidine ring with an *S* configuration at C15 as in the (+)-gelsemine structure, an oxepinone ring was first generated by condensation of the acid group with the hydroxy group in **20** to give **12** in 85% yield. Intramolecular Michael

addition at low temperature yielded two inseparable diastereomers (15*S*,20*R*)-**21a** and (15*S*,20*S*)-**21b** in 66% yield in a 7:1 ratio. The mixture of diastereomers **21a** and **21b** became separable after the lactone group in **21** was reduced to an aldehyde with DIBAL-H at low temperature, and the resulting aldehyde group was protected with a dimethoxy group, affording the major isomer (15*S*,20*R*)-**22** in 48% yield in two steps. Oxidation of the hydroxy group in **22** provided the *cis* tetra-substituted piperidinoaldehyde **11** in 92% yield. The *cis* relationship was confirmed by a strong NOE observed between the two protons at carbon atoms C15 and C20.

After having synthesized the *cis* tetra-substituted piperidinoaldehyde **11** with the required stereochemistry, we began to prepare compound **9**, which is the precursor for forming the D ring through a Michael addition (Scheme 3). Condensation of **11** with *N*-methoxyoxindole **10** in the presence of LDA at -78°C , followed by dehydration of the resulting hydroxy group with SOCl_2 /pyridine provided two separable geometric isomers (*Z*)-**9a** and (*E*)-**9b** in a 1.5:1 ratio. The *Z* conformation of the double bond in **9a** was confirmed by a strong NOE observed between the olefin proton at C6 and the aromatic proton at C9. As expected, the mixture of **9a** and **9b** easily underwent Michael addition in the presence of LDA in THF at -78°C to yield two inseparable diastereomers **23a** and **23b** epimerized at C7 in 75% yield. The Michael addition of **9** not only formed the D ring, but also created the correct C20 quaternary carbon center with an *R* configuration. However, the addition led to an *S* configuration at C6 rather than the desired *R* configuration. The *S* configuration at C6 prevented that the oxindole moiety gained access to the oxonium cation that was formed in situ when the MOM group and the two methoxy groups in **23** were removed under acidic conditions. Indeed, initial attempts to form the E and F ring by enol-oxonium cyclization generated both olefin **24** and pyranol **25** when the mixture of **23a** and **23b** was treated with either TFA at room temperature or TsOH in CHCl_3 at reflux.

To invert the C6 *S* configuration in **23** to an *R* configuration and thereby allow construction of the E and F rings by enol-oxonium cyclization, a double bond between C6 and C7 was generated to yield two separable geometric isomers **26a** and **26b** in 72% yield in a 5:1 ratio by using a standard protocol of olefination with PhSeCl/LDA/NaIO_4 (Scheme 3). Hydrogenation of the double bond in **26** with Lindlar catalyst at 1 atm of hydrogen pressure in MeOH within three hours resulted in two inseparable diastereomers **27a** and **27b** in 90% yield in a 2:1 ratio. The hydrogenation proceeded such that the nucleophilic attack occurred from the less hindered bottom face of the double bond, thereby yielding **27a** and **27b** with C6 exclusively in the *R* configuration. In contrast, hydrogenation of **26** with the more active catalyst Pd/C under the same conditions yielded two inseparable *N*-demethoxy products **28a** and **28b** in a 3:1 ratio in quantitative yield.

With **27a** and **27b** in hand, we carried out the planned one-pot, multi-step enol-oxonium cyclization reaction cascade to efficiently assemble both the E and F rings, and simultaneously establish the C3- and C7-stereocenters. When the mixture of **27a** and **27b** was treated with stoichiometric TsOH in CH_3Cl at reflux for four hours, two separable



Scheme 3. Synthesis of (+)-gelsemine. Reagents and conditions: a) **10**, LDA, THF, -78°C , 12 h, 75%; b) SOCl_2 , pyridine, 0°C , 5 min, 90%; c) LDA, THF, -78°C , 30 min, 75%; d) condition a: TfOH , CHCl_3 , RT, 24 h or condition b: $p\text{-TsOH}$, CHCl_3 , reflux, 10 h. e) LDA, PhSeCl , THF, -78°C , 30 min; then NaIO_4 , NaHCO_3 , $\text{MeOH}/\text{THF}/\text{H}_2\text{O}$ 5:1:1, 25°C , 10 h, 70% in 2 steps; f) Lindlar cat., H_2 , MeOH , 25°C , 3 h, 90% (gave **27a** and **27b**); g) 5% Pd/C , H_2 , MeOH , 25°C , 10 h, 98% (gave **28a** and **28b**); h) $p\text{-TsOH}$, CHCl_3 , reflux, 4 h, 73%, **29a/29b** 1:10 ratio; i) 5% Pd/C , H_2 , MeOH , 25°C , 10 h, 98%; j) DIBAL-H, H_2 , -78°C , 10 h, 72%; k) Tebbe reagent, THF, pyridine, reflux, 10 h, 60%. LDA = lithium diisopropylamide, Tf = trifluoromethanesulfonyl.

diastereomers **29a** and **29b** were obtained in 73% yield in a 1:10 ratio. Apparently, the reaction proceeded via two transition states **8a** and **8b**. The natural major isomer **29b** with the desired 7S configuration was produced from the favored transition state **8b**, in which an electronic repulsion between the oxonium cation and the hydroxy group of the enol functionality is avoided. In contrast, the disfavored transition state **8a**, in which both electron repulsion and a steric effect are present, led to the unnatural minor isomer

29a with the inverse 7R configuration. Interestingly, when the same reaction conditions were applied to **28a** and **28b**, a complex mixture resulted, rather than the corresponding cyclization products **30a** and **30b**. These results indicate that, in the absence of a methoxy substituent on the oxindole nitrogen atom, an enol functionality does not efficiently form during the reaction.^[11]

After having successfully synthesized **29b**, which possesses all the ring systems and correct stereochemistry of the carbon stereocenters of the target (+)-gelsemine, we turned our effort to the final task of converting the nitrile substituent at C20 in **29b** to an ethylene substituent (Scheme 3). After removal of the methoxy group in **29b** by hydrogenation with 5% Pd/C , the nitrile group in **30b** was reduced to an aldehyde group to provide **31** in 71% yield in two steps. Initial efforts with olefination of **31** using the Wittig reagent $\text{Ph}_3\text{P}=\text{CH}_2$ failed under various conditions, because the aldehyde group in **31** was severely hindered. Fortunately, olefination of **31** with Tebbe reagent yielded (+)-gelsemine in 60% yield.^[12] The synthetic (+)-gelsemine showed ^1H and ^{13}C NMR spectra that were identical to those of a previous report^[6f] and the commercially available (+)-natural gelsemine.^[13]

In summary, the biomimetic total synthesis of (+)-gelsemine from (R,R)-aziridine **16** has been accomplished in 25 steps with approximately 1% overall yield. A multi-step, one-pot enol-oxonium cyclization cascade is used in a highly efficient way to construct, simultaneously, the E ring, F ring, C3-stereocenter, and C7 quaternary stereocenter. The current synthesis using the enol-oxonium cyclization reaction as a key step to make the cage structure has demonstrated a possible biosynthetic pathway. We also demonstrated that without a methoxy substituent on the oxindole nitrogen atom, the enol-oxonium cyclization does not work (failure of converting compound **28** to compound **30**). This result is meaningful for elucidating the function of the methoxy group in the proposed biosynthetic procedure, and to partially explain why most members of the gelsemine family have a methoxy substituent on the oxindole nitrogen atom.

Received: March 3, 2012
Published online: April 4, 2012

Keywords: asymmetric synthesis · biomimetic synthesis · indole alkaloids · natural products · total synthesis

- [1] F. L. Sonnenschein, *Ber. Dtsch. Chem. Ges.* **1876**, 9, 1182–1186.
- [2] a) H. Conroy, J. K. Chakrabarti, *Tetrahedron Lett.* **1959**, 1, 6–13; b) Y. Schun, G. A. Cordell, *J. Nat. Prod.* **1985**, 48, 969–971.
- [3] F. M. Lovell, R. Pepinsky, A. J. C. Wilson, *Tetrahedron Lett.* **1959**, 1, 1–5.
- [4] a) Y. Schun, G. A. Cordell, M. Garland, *J. Nat. Prod.* **1986**, 49, 483–487; b) Y. Schun, G. A. Cordell, *J. Nat. Prod.* **1986**, 49, 806–

- 808; c) D. Ponglux, S. Wongseripipatana, S. Subhadhirasakul, H. Takayama, M. Yokota, K. Ogata, C. Phisalaphong, N. Aimi, S.-I. Sakai, *Tetrahedron* **1988**, *44*, 5075–5094; d) L.-Z. Lin, Y. Schun, G. A. Cordell, C.-Z. Ni, J. Clardy, *Phytochemistry* **1991**, *30*, 679–683.
- [5] For model studies and synthesis of a fragment of gelsemine, see: a) R. L. Autrey, F. C. Tahk, *Tetrahedron* **1967**, *23*, 901–917; b) R. L. Autrey, F. C. Tahk, *Tetrahedron* **1968**, *24*, 3337–3345; c) R. S. Johnson, T. O. Lovett, T. S. Stevens, *J. Chem. Soc. C* **1970**, *6*, 796–800; d) I. Fleming, J. P. Michael, *J. Chem. Soc. Perkin Trans. 1* **1981**, 1549–1556; e) I. Fleming, M. A. Loreto, J. P. Michael, I. H. M. Wallace, *Tetrahedron Lett.* **1982**, *23*, 2053–2056; f) I. Fleming, M. A. Loreto, I. H. M. Wallace, J. P. Michael, *J. Chem. Soc. Perkin Trans. 1* **1986**, 349–359; g) G. Stork, M. E. Krafft, S. A. Biller, *Tetrahedron Lett.* **1987**, *28*, 1035–1038; h) R. J. Vijn, H. Hiemstra, J. J. Kok, M. Knotter, W. N. Speckamp, *Tetrahedron* **1987**, *43*, 5019–5030; i) M. M. Abelman, T. Oh, L. E. Overman, *J. Org. Chem.* **1987**, *52*, 4130–4133; j) C. Clarke, I. Fleming, J. M. D. Fortunak, P. T. Gallagher, M. C. Honan, A. Mann, C. O. Nubling, P. R. Raithby, J. J. Wolff, *Tetrahedron* **1988**, *44*, 3931–3934; k) H. Hiemstra, R. J. Vijn, W. N. Speckamp, *J. Org. Chem.* **1988**, *53*, 3882–3884; l) W. G. Earley, E. J. Jacobsen, G. P. Meier, T. Oh, L. E. Overman, *Tetrahedron Lett.* **1988**, *29*, 3781–3784; m) W. G. Earley, T. Oh, L. E. Overman, *Tetrahedron Lett.* **1988**, *29*, 3785–3788; n) J.-K. Choi, D.-C. Ha, D. J. Hart, C.-S. Lee, S. Ramesh, S. Wu, *J. Org. Chem.* **1989**, *54*, 279–290; o) I. Fleming, R. C. Moses, M. Tercel, J. Ziv, *J. Chem. Soc. Perkin Trans. 1* **1991**, 617–626; p) D. J. Hart, S. C. Wu, *Tetrahedron Lett.* **1991**, *32*, 4099–4102; q) W.-J. Koot, H. Hiemstra, W. N. Speckamp, *J. Org. Chem.* **1992**, *57*, 1059–1061; r) A. Madin, L. E. Overman, *Tetrahedron Lett.* **1992**, *33*, 4859–4862; s) L. E. Overman, M. J. Sharp, *J. Org. Chem.* **1992**, *57*, 1035–1038; t) D. J. Hart, S. C. Wu, *Heterocycles* **1993**, *35*, 135–138; u) H. Takayama, N. Seki, M. Kitajima, N. Aimi, S.-I. Sakai, *Nat. Prod. Lett.* **1993**, *2*, 271–276; v) A. P. Johnson, R. W. A. Luke, R. W. Steele, A. N. Boa, *J. Chem. Soc. Perkin Trans. 1* **1996**, 883–893; w) F. Ng, P. Chiu, S. J. Danishefsky, *Tetrahedron Lett.* **1998**, *39*, 767–770; x) M. J. Sung, C.-W. Lee, J. K. Cha, *Synlett* **1999**, 561–562; y) A. G. Avent, P. W. Byrne, C. S. Penkett, *Org. Lett.* **1999**, *1*, 2073–2075; z) J. Dijkink, J.-C. Cintrat, W. N. Speckamp, H. Hiemstra, *Tetrahedron Lett.* **1999**, *40*, 5919–5922; aa) A. J. Pearson, X. Wang, *J. Am. Chem. Soc.* **2003**, *125*, 13326–13327; ab) W. G. Earley, J. E. Jacobsen, A. Madin, G. P. Meier, C. J. O'Donnell, T. Oh, D. W. Old, L. E. Overman, M. J. Sharp, *J. Am. Chem. Soc.* **2005**, *127*, 18046–18053; ac) A. Madin, C. J. O'Donnell, T. Oh, D. W. Old, L. E. Overman, M. J. Sharp, *J. Am. Chem. Soc.* **2005**, *127*, 18054–18065; ad) S. Grecian, J. Aube, *Org. Lett.* **2007**, *9*, 3153–3156; ae) K. Tchabanenko, N. S. Simpkins, L. Male, *Org. Lett.* **2008**, *10*, 4747–4750.
- [6] For a review on the total synthesis of gelsemine, see: H. Lin, S. J. Danishefsky, *Angew. Chem.* **2003**, *115*, 38–53; *Angew. Chem. Int. Ed.* **2003**, *42*, 36–51. For individual total syntheses of racemic gelsemine, see: a) Z. Sheikh, R. Stell, A. S. Tasker, A. P. Johnson, *J. Chem. Soc. Chem. Commun.* **1994**, 763–764; b) J. K. Dutton, R. W. Steel, A. S. Tasker, V. Popsavin, A. P. Johnson, *J. Chem. Soc. Chem. Commun.* **1994**, 765–766; c) N. J. Newcombe, F. Ya, R. J. Vijn, H. Hiemstra, W. N. Speckamp, *J. Chem. Soc. Chem. Commun.* **1994**, 767–768; d) S. Atarashi, J. K. Choi, D. C. Ha, D. J. Hart, D. Kuzmich, C.-S. Lee, S. Ramesh, S. C. Wu, *J. Am. Chem. Soc.* **1997**, *119*, 6226–6241; e) T. Fukuyama, G. Liu, *J. Am. Chem. Soc.* **1996**, *118*, 7426–7427; f) A. Madin, C. J. O'Donnell, T. Oh, D. W. Old, L. E. Overman, M. J. Sharpe, *Angew. Chem.* **1999**, *111*, 3110–3112; *Angew. Chem. Int. Ed.* **1999**, *38*, 2934–2936; g) S. Yokoshima, H. Tokuyama, T. Fukuyama, *Angew. Chem.* **2000**, *112*, 4239–4241; *Angew. Chem. Int. Ed.* **2000**, *39*, 4073–4075; h) F. W. Ng, H. Lin, S. J. Danishefsky, *J. Am. Chem. Soc.* **2002**, *124*, 9812–9824.
- [7] Previously proposed biosynthetic pathway, see: a) H. Conroy, J. K. Chakrabarti, *Tetrahedron Lett.* **1959**, *1*, 6–13; b) J. Stöckigt, *Tetrahedron Lett.* **1979**, *20*, 2615–2618; c) P. Dhavadee, W. Sumphan, S. Sanan, T. Hiromitsu, Y. Masaki, O. Koreharu, P. Chada, A. Norio, S. Shinichiro, *Tetrahedron* **1988**, *44*, 5075–5094; d) Z.-J. Liu, R.-R. Liu in *The Alkaloids*, Vol. 33 (Ed.: R. H. F. Manske), Academic Press, New York, **1988**, pp. 83–140.
- [8] R. Hili, A. K. Yudin, *J. Am. Chem. Soc.* **2006**, *128*, 14772–14773.
- [9] B. Nyasse, L. Grehnb, U. Ragnarsson, *Chem. Commun.* **1997**, 1017–1018.
- [10] K. Fuji, T. Kawabata, Y. Kriyu, Y. Sugiura, *Heterocycles* **1996**, *42*, 701–722.
- [11] A similar enol functionality of *N*-methoxy oxoindole was reported in the synthesis of racemic 7-epi-20-desethylgelsedine, see: A. S. Kende, M. J. Luzzio, J. S. Mendoza, *J. Org. Chem.* **1990**, *55*, 918–924.
- [12] F. N. Tebbe, G. W. Parshall, G. S. Reddy, *J. Am. Chem. Soc.* **1978**, *100*, 3611–3613.
- [13] The standard sample of natural (+)-gelsemine was purchased from Sichuan Weikeqi BioTech Co., Ltd (China).