


# Palladium-Catalyzed Oxidative Double C–H Functionalization/Carbonylation for the Synthesis of Xanthenes\*\*

Hua Zhang, Renyi Shi, Pei Gan, Chao Liu, Anxing Ding, Qiuyi Wang, and Aiwen Lei\*

The chemical structure of xanthone constitutes the central core of a wide variety of naturally occurring and manmade compounds, which exhibit extraordinary biological and pharmaceutical properties (e.g. antibacterial, anti-inflammatory, anticancer, and antiviral).<sup>[1]</sup> The xanthone scaffold has even been described as “privileged structure”, since members of this structural class are able to interact with different types of drug targets and attracted interests across a broad spectrum of sciences from chemistry and biology to medicine (Scheme 1).<sup>[2]</sup> Among the known synthetic routes to obtain



**Scheme 1.** Strategies towards syntheses of xanthenes. X=OH, OMe.

the xanthone skeleton, the cyclodehydration of 2,2'-dihydroxybenzophenones and electrophilic cycloacylation of 2-aryloxybenzoic acids are the most popular methods (Scheme 1).<sup>[3]</sup> However, both methods involve multistep procedures under harsh reaction conditions, in which strong acids or toxic metals are often used.<sup>[4]</sup> Because of the important biological applications of xanthenes, the development of an efficient and general synthetic route with reduced waste and in fewer steps is highly desirable.

The double C–H functionalization/carbonylation of simple diaryl ethers with CO would represent a very attractive and sustainable approach towards the syntheses of xanthone, because neither of the two C–H bonds need to be prefunctionalized, and diaryl ethers can be easily synthesized from basic chemicals (Scheme 1). Transition-metal-catalyzed carbonylation of aromatic halides with CO in the presence of various nucleophiles has undergone rapid development,<sup>[5]</sup> since the pioneering work of Heck and co-workers in 1974.<sup>[6]</sup> During the past decade, increased attention has been focused on Pd-catalyzed aromatic C–H functionalization/carbonylation.<sup>[7]</sup> Very recently, Pd-catalyzed direct transformation of two C–H bonds simultaneously has attracted more and more interest.<sup>[8]</sup> However, double C–H functionalization/carbonylation remains an outstanding challenge.

Owing to the synthetic importance of the xanthone structure and our continuous interests in the Pd-catalyzed oxidative coupling<sup>[8c]</sup> and carbonylation reactions,<sup>[7b]</sup> herein, we describe the first Pd-catalyzed double C–H functionalization/carbonylation of diaryl ethers to form biologically active xanthone derivatives.

The combination of Pd(OAc)<sub>2</sub> and K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> in trifluoroacetic acid (TFA) at 50 °C gave the best result for the oxidative carbonylation of 4-tolyl ether (**1a**) under 1 atm CO (Table 1, entry 1). With these conditions, a 93 % yield of 2,7-

**Table 1:** Pd-catalyzed oxidative double C–H carbonylation of **1a**: effects of reaction parameters.<sup>[a]</sup>

Entry	Variation from “standard conditions”	Yield[%] <sup>[b]</sup>
1	none	93
2	PdCl <sub>2</sub> , instead of Pd(OAc) <sub>2</sub>	0
3	[Pd(dba) <sub>3</sub> ], instead of Pd(OAc) <sub>2</sub>	17
4	MnO <sub>2</sub> , instead of K <sub>2</sub> S <sub>2</sub> O <sub>8</sub>	87
5	BQ, instead of K <sub>2</sub> S <sub>2</sub> O <sub>8</sub>	90
6	Na <sub>2</sub> S <sub>2</sub> O <sub>8</sub> , instead of K <sub>2</sub> S <sub>2</sub> O <sub>8</sub>	47
7	Cu(OAc) <sub>2</sub> , instead of K <sub>2</sub> S <sub>2</sub> O <sub>8</sub>	34
8	O <sub>2</sub> , instead of K <sub>2</sub> S <sub>2</sub> O <sub>8</sub>	0
9	TFA/TFAA (9:1), instead of TFA	56
10	TFA/HOAc (1:9), instead of TFA	0
11	25 °C, instead of 50 °C	32
12	diphenyl ether, instead of <b>1a</b>	27

[a] Standard reaction conditions: **1a** (0.2 mmol), Pd(OAc)<sub>2</sub> (2.5 mol %), K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> (2 equiv), 1 atm CO, TFA (1.0 mL), 50 °C, 2 h. [b] The yield was determined by GC, calibrated using biphenyl as internal standard.

BQ = 1,4-benzoquinone, dba = dibenzylideneacetone, TFAA = trifluoroacetic anhydride.


[\*] H. Zhang, R. Shi, P. Gan, C. Liu, A. Ding, Q. Wang, Prof. A. Lei  
The College of Chemistry and Molecular Sciences  
Wuhan University  
Wuhan, Hubei (P.R. China)  
E-mail: aiwenlei@whu.edu.cn  
Prof. A. Lei  
State Key Laboratory for Oxo Synthesis and Selective Oxidation  
Lanzhou Institute of Chemical Physics  
Chinese Academy of Sciences  
730000 Lanzhou (China)

[\*\*] This work was supported by the National Natural Science Foundation of China (21025206, 20832003, and 20972118) and the 973 Program (2012CB725302).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201201050>.




As shown in Figure 1, zero-order kinetics for the carbonylation of different substrates were observed, thus indicating that the first C–H functionalization of diaryl ether with Pd<sup>II</sup>



**Figure 1.** Kinetic plots of oxidative double C–H carbonylation of different substrates.

was not the rate-determining step. Furthermore, the carbonylation of **1a** was conducted under different pressures of CO (Figure 2). The kinetic profiles were overlaid, indicating that the insertion of CO was not rate-determining under the




**Figure 2.** Kinetic profiles of oxidative double C–H carbonylation of **1a** with different CO pressures.

conditions. The reductive elimination of ArCOPdAr species has been reported to be a facile process;<sup>[9]</sup> therefore, we concluded that the second C–H functionalization might be the rate-determining step of this catalytic cycle.

On the basis of above results and previous studies,<sup>[10]</sup> we propose a mechanism for this double C–H functionalization/carbonylation reaction (Scheme 2). Pd(OTFA)<sub>2</sub> is formed in situ by the reaction of Pd(OAc)<sub>2</sub> with TFA. Then, the fast electrophilic palladation of **1a** with Pd(OTFA)<sub>2</sub> affords the arylpalladium species **I**, which further reacts with CO to produce the intermediate **II**. The subsequent intramolecular C–H functionalization and reductive elimination of **III** gives the xanthone **2a** and generates the Pd<sup>0</sup> species. Pd(OTFA)<sub>2</sub> is then regenerated by oxidation with K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> in the presence of TFA.

In conclusion, we have developed the first Pd-catalyzed double C–H functionalization/carbonylation of diaryl ethers to form xanthenes. By using a simple catalytic system



**Scheme 2.** Proposed reaction mechanism.

consisting of Pd(OAc)<sub>2</sub>, K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>, and TFA, a variety of diaryl ethers could be directly carbonylated to xanthenes in moderate to good yields. Moreover, various functional groups were tolerated under the optimized conditions. Notably, this transformation provides an effective and practical protocol towards the syntheses of bioactive xanthenes. Preliminary mechanism studies revealed that the second C–H functionalization might be the rate-determining step. Further studies on substrate scope and mechanism are currently underway and will be reported in due course.

## Experimental Section

**General procedure:** In an oven-dried Schlenk tube equipped with a stir bar, Pd(OAc)<sub>2</sub> (2.5 mol %), K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> (2 equiv), and diaryl ether (0.2 mmol) were combined. A balloon filled with CO was connected to the Schlenk tube by the side tube and purged three times. Then, TFA (1.0 mL) was added to the tube through a syringe. The Schlenk tube was heated at 50 °C for 2–6 h (as indicated by TLC) and then cooled to room temperature. After the gas from the balloon was released carefully, the reaction was quenched by water and extracted with CH<sub>2</sub>Cl<sub>2</sub> three times. The combined organic layers were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and the solvent was evaporated in vacuum. The desired products were obtained in the corresponding yields after purification by flash chromatography on silica gel with petroleum ether and ethyl acetate.

Received: February 8, 2012

Published online: April 4, 2012

**Keywords:** C–H activation · oxidative carbonylation · palladium · synthetic methods · xanthenes

- [1] a) V. Peres, T. J. Nagem, F. F. de Oliveira, *Phytochemistry* **2000**, 55, 683; b) M. M. M. Pinto, M. E. Sousa, M. S. J. Nascimento, *Curr. Med. Chem.* **2005**, 12, 2517; c) L. M. M. Vieira, A. Kijjoa, *Curr. Med. Chem.* **2005**, 12, 2413; d) M. Riscoe, J. X. Kelly, R. Winter in *Curr. Med. Chem.*, Vol. 12, **2005**, pp. 2539; e) H. R. El-Seedi, D. M. H. El-Ghorab, M. A. El-Barbary, M. F. Zayed, U. Goeransson, S. Larsson, R. Verpoorte, *Curr. Med. Chem.* **2009**, 16, 2581; f) M. Saleem, M. Nazir, M. S. Ali, H. Hussain, Y. S. Lee, N. Riaz, A. Jabbar, *Nat. Prod. Rep.* **2010**, 27, 238.

- [2] a) M. Pedro, F. Cerqueira, M. E. Sousa, M. S. J. Nascimento, M. Pinto, *Bioorg. Med. Chem.* **2002**, *10*, 3725; b) N. P. Minton, *Nat. Rev. Microbiol.* **2003**, *1*, 237; c) L. Gales, A. M. Damas, *Curr. Med. Chem.* **2005**, *12*, 2499; d) G. M. Tozer, C. Kanthou, B. C. Baguley, *Nat. Rev. Cancer* **2005**, *5*, 423; e) Y. Na, *J. Pharm. Pharmacol.* **2009**, *61*, 707; f) H. T. Nguyen, M.-C. Lallemant, S. Bouteffouchet, S. Michel, F. Tillequin, *J. Nat. Prod.* **2009**, *72*, 527; g) N. Pouli, P. Marakos, *Anti-Cancer Agents Med. Chem.* **2009**, *9*, 77; h) H. R. El-Seedi, M. A. El-Barbary, D. M. H. El-Ghorab, L. Bohlin, A.-K. Borg-Karlson, U. Goeransson, R. Verpoorte, *Curr. Med. Chem.* **2010**, *17*, 854.
- [3] a) M. Afzal, J. M. Al-Hassan, *Heterocycles* **1980**, *14*, 1173; b) M. E. Sousa, M. M. M. Pinto, *Curr. Med. Chem.* **2005**, *12*, 2447; c) M. M. M. Pinto, R. A. P. Castanheiro, *Curr. Org. Chem.* **2009**, *13*, 1215.
- [4] J. Zhao, R. C. Larock, *Org. Lett.* **2005**, *7*, 4273.
- [5] a) M. Beller, B. Cornils, C. D. Frohning, C. W. Kohlpaintner, *J. Mol. Catal. A* **1995**, *104*, 17; b) R. Skoda-Foldes, L. Kollar, *Curr. Org. Chem.* **2002**, *6*, 1097; c) C. F. J. Barnard, *Organometallics* **2008**, *27*, 5402; d) A. Brennfürher, H. Neumann, M. Beller, *Angew. Chem.* **2009**, *121*, 4176; *Angew. Chem. Int. Ed.* **2009**, *48*, 4114; e) X.-F. Wu, H. Neumann, M. Beller, *Chem. Soc. Rev.* **2011**, *40*, 4986.
- [6] a) A. Schoenberg, I. Bartoletti, R. F. Heck, *J. Org. Chem.* **1974**, *39*, 3318; b) A. Schoenberg, R. F. Heck, *J. Org. Chem.* **1974**, *39*, 3327.
- [7] a) B. Gabriele, G. Salerno, M. Costa, *Top. Organomet. Chem.* **2006**, *18*, 239; b) Q. Liu, H. Zhang, A. Lei, *Angew. Chem.* **2011**, *123*, 10978; *Angew. Chem. Int. Ed.* **2011**, *50*, 10788; c) Y. Fujiwara, K. Takaki, Y. Taniguchi, *Synlett* **1996**, 591; d) W. Lu, Y. Yamaoka, Y. Taniguchi, T. Kitamura, K. Takaki, Y. Fujiwara, *J. Organomet. Chem.* **1999**, *580*, 290; e) I. O. Kalinovskii, V. V. Pogorelov, A. I. Gelbshtein, N. G. Akhmetov, *Russ. J. Gen. Chem.* **2001**, *71*, 1457; f) K. Orito, A. Horibata, T. Nakamura, H. Ushito, H. Nagasaki, M. Yuguchi, S. Yamashita, M. Tokuda, *J. Am. Chem. Soc.* **2004**, *126*, 14342; g) S. Ohashi, S. Sakaguchi, Y. Ishii, *Chem. Commun.* **2005**, 486; h) R. Giri, J.-Q. Yu, *J. Am. Chem. Soc.* **2008**, *130*, 14082; i) C. E. Houlden, M. Hutchby, C. D. Bailey, J. G. Ford, S. N. G. Tyler, M. R. Gagne, G. C. Lloyd-Jones, K. I. Booker-Milburn, *Angew. Chem.* **2009**, *121*, 1862; *Angew. Chem. Int. Ed.* **2009**, *48*, 1830; j) R. Giri, J. K. Lam, J.-Q. Yu, *J. Am. Chem. Soc.* **2010**, *132*, 686; k) H. Li, G.-X. Cai, Z.-J. Shi, *Dalton Trans.* **2010**, *39*, 10442; l) B. López, A. Rodríguez, D. Santos, J. Albert, X. Ariza, J. Garcia, J. Granell, *Chem. Commun.* **2011**, *47*, 1054; m) B. Haffemayer, M. Gulias, M. J. Gaunt, *Chem. Sci.* **2011**, *2*, 312; n) H. Zhang, D. Liu, C. Chen, C. Liu, A. Lei, *Chem. Eur. J.* **2011**, *17*, 9581.
- [8] a) D. R. Stuart, K. Fagnou, *Science* **2007**, *316*, 1172; b) C. S. Yeung, V. M. Dong, *Chem. Rev.* **2011**, *111*, 1215; c) C. Liu, H. Zhang, W. Shi, A. Lei, *Chem. Rev.* **2011**, *111*, 1780; d) S. H. Cho, J. Y. Kim, J. Kwak, S. Chang, *Chem. Soc. Rev.* **2011**, *40*, 5068.
- [9] J. W. Labadie, J. K. Stille, *J. Am. Chem. Soc.* **1983**, *105*, 6129.
- [10] a) Y. Izawa, I. Shimizu, A. Yamamoto, *Bull. Chem. Soc. Jpn.* **2004**, *77*, 2033; b) Y. Hu, J. Liu, Z. Lu, X. Luo, H. Zhang, Y. Lan, A. Lei, *J. Am. Chem. Soc.* **2010**, *132*, 3153.