

Imidazolidinone-Derived Enamines: Nucleophiles with Low Reactivity**

Sami Lakhdar,* Biplab Maji, and Herbert Mayr*

Dedicated to Professor Wolfgang Beck on the occasion of his 80th birthday

During the last decade the concept of enamine activation has become a powerful tool in asymmetric synthesis. It uses chiral secondary amines as catalysts to activate saturated carbonyl compounds by conversion into enamines.^[1] Among various catalysts tested, diarylprolinol silyl ether **1b** was found to be particularly useful for the stereoselective introduction of different functionalities to the α -position of aldehydes.^[2,3]










Imidazolidinones **1c–e** (Table 1), which have been used extensively in iminium catalysis,^[4] are less effective in enamine-activated processes unless strong electrophiles are employed. Typical examples are enantioselective α -halogenations^[5] and α -alkylations of aldehydes with stabilized carbocations which were in situ generated by treatment of alcohols with acids.^[6] Mechanistic investigations, focusing on the characterization and reactivities of the intermediate enamines, are rare.^[7–9]

While the synthesis and the X-ray structure of the enamine **3b** have previously been described by Seebach et al.,^[8] we are not aware of any X-ray structures of enamines derived from imidazolidinones. Gellman et al. used ¹H NMR spectroscopy to characterize the enamine generated from imidazolidinone **1c** and 3-phenylpropanal in DMSO solution and reported its reaction with methyl vinyl ketone catalyzed by 4-ethoxycarbonylcatechol.^[9]

In order to elucidate the relationships between structure and reactivities of enamines derived from **1a–e**, we have now synthesized the enamines **3a–e**, performed X-ray analyses of **3d** and **3e**, and measured the kinetics of their reactions with the stabilized benzhydrylium ions **4a–h** (Table 2).

Enamines **3c–e**, which had not been isolated previously, were obtained by refluxing phenylacetaldehyde (**2**) and the amines **1c–e** in the presence of 1 mol % of *p*-toluenesulfonic acid in toluene under argon using a Dean–Stark apparatus filled with molecular sieves (4 Å) to remove the generated water.^[10] After evaporation of the solvent, **3e** was immedi-

Table 1: Amines **1a–e** and the corresponding phenylacetaldehyde-derived enamines **3a–e**.

$\text{HNRR}' + \text{O}=\text{CH}-\text{Ph} \xrightarrow[\text{toluene, reflux}]{p\text{-TsOH (1\%)}}$		$\text{R}'\text{N}(\text{R})=\text{CH}-\text{Ph}$	
Amine	Product		Yield [%]
	1a		3a 86 ^[a]
	1b		3b 35 ^[b]
	1c		3c 44 ^[c]
	1d		3d 32 ^[c]
	1e		3e 87 ^[d]

[a] After distillation. [b] **3b** was prepared by heating **1b** and **2** in benzene at reflux following a procedure described by Seebach (Ref. [8]). After solvent removal and crystallization from Et₂O, **3b** was obtained as a pure material. [c] After column chromatography. [d] After crystallization.

ately obtained as a crystalline material in 87% yield, while column chromatography was employed to separate the enamines **3c** and **3d** from the nonreacted imidazolidinones **1c** and **1d**, respectively.

Crystals of **3d** and **3e** suitable for X-ray analysis were grown by the vapor diffusion crystallization method in diethyl ether/*n*-pentane mixtures. As shown in Figure 1,^[11] the C–N bond between the heterocyclic ring and the *E*-configured C=C bond has *s*-trans conformation in both enamines **3d** and **3e**. Whereas the benzylic phenyl group is located over the imidazolidinone ring in **3d**, possibly because of stabilizing CH– π interactions (London dispersion interaction between the phenyl ring and the cyclopentane ring),^[12] the benzyl group is located over the C=C bond in the enamine **3e** and thus directs electrophiles to the *Si* face of the C=C bond.^[5] NMR spectroscopy showed that the conformations of **3d** and **3e**, which were observed in the crystals, also dominate in CDCl₃ solution (see the Supporting Information).

[*] Dr. S. Lakhdar, M. Sc. B. Maji, Prof. Dr. H. Mayr
Department Chemie, Ludwig-Maximilians-Universität München
Butenandtstrasse 5–13 (Haus F), 81377 München (Germany)
E-mail: sami.lakhdar@cup.uni-muenchen.de
herbert.mayr@cup.uni-muenchen.de
Homepage: <http://www.cup.uni-muenchen.de/oc/mayr>

[**] We thank the Alexander von Humboldt Foundation (research fellowship for S.L.) and the Deutsche Forschungsgemeinschaft (SFB 749) for financial support, Dr. P. Mayer for the determination of the X-ray structures, and Dr. A. R. Ofial and Dr. M. Horn for helpful discussions.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201201240>.




Figure 1. Crystal structures of the enamines **3d** (left) and **3e** (right).

From the pyramidalization parameter Δ , defined by Dunitz as the distance between the N atom and the plane formed by the three attached carbon atoms,^[13] one can derive that the almost planar sp^2 -hybridized nitrogen in **3b** ($\Delta = 0.037 \text{ \AA}$)^[8] adopts more and more sp^3 character as one moves to **3d** ($\Delta = 0.155 \text{ \AA}$) and eventually to **3e** ($\Delta = 0.293 \text{ \AA}$).

In order to quantify the nucleophilic reactivities of **3a–e** we have studied the kinetics of their reactions with the benzhydrylium ions **4a–h** (Table 2), which have been used as reference electrophiles for the construction of comprehensive nucleophilicity scales on the basis of Equation (1). Herein, nucleophiles are characterized by two parameters (nucleophilicity N and sensitivity parameter s_N) and electrophiles by one parameter (electrophilicity E).^[14]

$$\lg k_2(20^\circ\text{C}) = s_N(N + E) \quad (1)$$


As described previously,^[14b] the rates of the reactions of carbocations **4** with the enamines **3a–e** were measured photometrically by following the disappearance of the UV/

Table 2: Reference electrophiles **4a–h**.

		$E^{[a]}$
	4a	−1.36
	4b	−3.14
	4c	−3.85
	4d	−5.53
	4e	−7.02
	4f	−7.69
	4g ($n=2$)	−8.22
	4h ($n=1$)	−8.76

[a] Electrophilicity parameters E from Ref. [14b].

Vis absorbances of the diarylcarbenium ions **4** (Scheme 1), using conventional and stopped-flow instruments. All kinetic experiments were performed at 20°C in acetonitrile with a large excess of the enamines **3a–e** in order to achieve conditions for first-order kinetics.



Scheme 1. Reactions of the enamines **3** with the carbocations **4** in acetonitrile at 20°C .

The first-order rate constants k_{obs} were obtained by least-squares fitting of the function $A_t = A_0 \exp(-k_{\text{obs}}t) + C$ to the time-dependent absorbances of the electrophiles. Plots of k_{obs} versus the concentrations of the nucleophiles [**3**] were linear, as exemplified in Figure 2. The slopes of these plots gave the second-order rate constants k_2 (in $\text{M}^{-1}\text{s}^{-1}$), which are summarized in Table 3.




Figure 2. Exponential decay of the absorbance at 586 nm during the reaction of **4c**- BF_4^- ($1.60 \times 10^{-5} \text{ M}$) with **3c** ($3.90 \times 10^{-4} \text{ M}$). Inset: Plot of the rate constants k_{obs} versus [**3c**] (20°C in CH_3CN).

Plots of $\lg k_2$ versus the empirical electrophilicity parameters E are linear for all reactions studied in this investigation (Figure 3), indicating that Equation (1) can be used to determine N and s_N parameters for the enamines **3a–e** (Table 3).

One can see that the enamine **3b**, which is derived from the Hayashi–Jørgensen catalyst **1b**, is almost two orders of magnitude less reactive than **3a**, the parent compound of this series. As one of the diastereotopic faces of **3b** is completely open for electrophilic attack, the reduction of reactivity must predominantly be due to the electron-withdrawing inductive effect of the trimethylsilyloxybenzhydryl group in **3b**.

The inductive electron-withdrawing effect of the extra endocyclic amido group in the imidazolidinone derivatives **3c** and **3d**, the pyramidalization of the enamine nitrogen, and the steric shielding of both faces of the $\text{C}=\text{C}$ bond by the two alkyl groups at the 2-position of the imidazolidinone ring reduce

Table 3: Second-order rate constants k_2 for the reactions of the carbocations **4a–h** with the enamines **3a–e** (acetonitrile, 20 °C).

Enamine	R ⁺	k_2 [M ⁻¹ s ⁻¹]	N , s_N
3a	4e	1.38×10^5	12.25, 0.99
	4f	3.48×10^4	
	4g	9.94×10^3	
	4h	2.64×10^3	
3b	4d	1.48×10^5	10.56, 1.01
	4e	2.33×10^3	
	4h	7.94×10^1	
3c	4a	4.73×10^6	7.20, 1.14
	4b	7.80×10^4	
	4c	2.24×10^3	
	4d	1.15×10^2	
3d	4b	2.77×10^5	7.92, 1.07
	4c	7.26×10^3	
	4d	4.93×10^2	
3e	4b	3.46×10^2	5.80, 0.87
	4c	2.56×10^1	
	4d	2.13	





Figure 3. Plots of $\lg k_2$ for the reactions of **3a–c**, and **3e** with the benzhydrylium ions **4** in CH₃CN at 20 °C versus the corresponding electrophilicity parameters E . (Correlation for **3d** is omitted for the sake of clarity; it is shown on page S19 of the Supporting Information).

the nucleophilicities of these enamines by another two to three orders of magnitude relative to **3b**.

As the C=C bond of **3e** has one open face, its low nucleophilicity (10^2 times less than that of **3c** and **3d**) must be due to the enhanced pyramidalization of the enamine nitrogen in **3e** (X-ray structure, Figure 1), which strongly reduces the electron density in the C=C bond.^[15]

While the interpretation of the NMR chemical shifts of the β -protons in **3a–e** is problematic because of the anisotropy of the phenyl groups, the ¹³C NMR chemical shifts show the lower electron densities at the β carbon of the imidazolidinone-derived enamines **3c–e** (Table 4).

Table 4: NMR chemical shifts (CDCl₃) of the enamines **3a–e**.

enamine		
	$\delta(\text{C}^\beta\text{-H})$ [ppm]	$\delta(\text{C}^\beta)$ [ppm]
3a	5.18	97.4
3b	5.02 ^[a]	97.2 ^[a]
3c	5.47	101.9
3d	5.47	102.1
3e	4.76	102.9

[a] From Ref. [8b].

The fact that **3b** has significantly higher nucleophilicity than **3c–e** may explain why **1b** is a more suitable catalyst than **1c–e** for most enamine-activated reactions.

By using the N and s_N values of **3b** (Table 3) and the E values of Michael acceptors,^[14] Equation (1) allows one to predict whether a reaction may take place at room temperature. Hayashi's observation that **1b** catalyzes Michael additions of aldehydes to β -nitrostyrenes^[16] is in line with a calculated second-order rate constant ($k_{\text{calcd}} = 4.8 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$) for the reaction of **3b** with β -nitrostyrene ($E = -13.85$).^[17] It has been shown, however, that the initially generated zwitterions from the enamines and nitrostyrene collapse with formation of cyclobutanes, the ring opening of which is the rate-determining step of the catalytic cycle.^[7e,18] For that reason, a sufficiently fast reaction of the enamine with the Michael acceptor is only one of the criteria that have to be fulfilled for a catalytic cycle to proceed.

The more than 100 times higher nucleophilicity of **3b** compared with **3c,d** may also rationalize the fact that **1b** and not **1c–e** are usually employed as catalysts for Mannich-type reactions of imines with aldehydes.^[19] In line with the low nucleophilicity of **3c–e** Gellman et al reported that **1c**-catalyzed conjugate additions of aldehydes to enones require Brønsted acids as cocatalysts to activate the enones.^[9]

In summary, we have described the first X-ray structures of enamines derived from imidazolidinones. Nucleophilic reactivities of these enamines have been determined from the kinetics of their reactions with diarylcarbenium ions **4**, which showed that the enamine **3b**, derived from the Hayashi–Jørgensen catalyst, is 10^3 to 10^5 times more nucleophilic than the enamines derived from the imidazolidinones **1c–e**.

Received: February 14, 2012

Published online: April 25, 2012

Keywords: enamines · kinetics · linear free energy relationships · nucleophilicity · pyramidalization

- [1] a) B. List, *Acc. Chem. Res.* **2004**, *37*, 548–557; b) W. Notz, F. Tanaka, C. F. Barbas III, *Acc. Chem. Res.* **2004**, *37*, 580–591; c) S. Mukherjee, J. W. Yang, S. Hoffmann, B. List, *Chem. Rev.* **2007**, *107*, 5471–5569; d) C. F. Barbas III, *Angew. Chem.* **2008**, *120*, 44–50; *Angew. Chem. Int. Ed.* **2008**, *47*, 42–47; e) A. Dondoni, A. Massi, *Angew. Chem.* **2008**, *120*, 4716–4739; *Angew. Chem. Int. Ed.* **2008**, *47*, 4638–4660; f) P. Melchiorre,

- M. Marigo, A. Carlone, G. Bartoli, *Angew. Chem.* **2008**, *120*, 6232–6265; *Angew. Chem. Int. Ed.* **2008**, *47*, 6138–6171; g) S. Bertelsen, K. A. Jørgensen, *Chem. Soc. Rev.* **2009**, *38*, 2178–2189; h) M. Nielsen, D. Worgull, T. Zweifel, B. Gschwend, S. Bertelsen, K. A. Jørgensen, *Chem. Commun.* **2011**, *47*, 632–649.
- [2] a) A. Mieglo, C. Palomo, *Chem. Asian J.* **2008**, *3*, 922–948; b) C. Palomo, A. Mielgo, *Angew. Chem.* **2006**, *118*, 8042–8046; *Angew. Chem. Int. Ed.* **2006**, *45*, 7876–7880.
- [3] For a recent review on the impact of diaryl prolinol silyl ethers in organocatalysis, see: K. L. Jensen, G. Dickmeiss, H. Jiang, L. Albrecht, K. A. Jørgensen, *Acc. Chem. Res.* **2012**, *45*, 248–264.
- [4] a) G. Lelais, D. W. C. MacMillan, *Aldrichimica Acta* **2006**, *39*, 79–87; b) A. Erkkilä, I. Majander, P. M. Pihko, *Chem. Rev.* **2007**, *107*, 5416–5470.
- [5] a) M. P. Brochu, S. P. Brown, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2004**, *126*, 4108–4109; b) D. D. Steiner, N. Mase, C. F. Barbas III, *Angew. Chem.* **2005**, *117*, 3772–3776; *Angew. Chem. Int. Ed.* **2005**, *44*, 3706–3710; c) T. D. Beeson, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2005**, *127*, 8826–8828.
- [6] For asymmetric S_N1 -type α -alkylations of aldehydes with carbocations, see: a) R. R. Shaikh, A. Mazzanti, M. Petrini, G. Bartoli, P. Melchiorre, *Angew. Chem.* **2008**, *120*, 8835–8838; *Angew. Chem. Int. Ed.* **2008**, *47*, 8707–8710; b) P. G. Cozzi, F. Benfatti, L. Zoli, *Angew. Chem.* **2009**, *121*, 1339–1342; *Angew. Chem. Int. Ed.* **2009**, *48*, 1313–1316; c) F. Benfatti, M. G. Capdevila, L. Zoli, E. Benedetto, P. G. Cozzi, *Chem. Commun.* **2009**, 5919–5921; d) F. Benfatti, E. Benedetto, P. G. Cozzi, *Chem. Asian J.* **2010**, *5*, 2047–2052; e) M. G. Capdevila, F. Benfatti, L. Zoli, M. Stenta, P. G. Cozzi, *Chem. Eur. J.* **2010**, *16*, 11237–11241; f) B. Zhang, S.-K. Xiang, L.-H. Zhang, Y. Cui, N. Jiao, *Org. Lett.* **2011**, *13*, 5212–5215; g) A. Gualandi, E. Emer, M. G. Capdevila, P. G. Cozzi, *Angew. Chem.* **2011**, *123*, 7988–7992; *Angew. Chem. Int. Ed.* **2011**, *50*, 7842–7846; h) L. Tak-Tak, H. Dhiman, P. I. Dalko, *Angew. Chem.* **2011**, *123*, 12350–12352; *Angew. Chem. Int. Ed.* **2011**, *50*, 12146–12147.
- [7] a) M. B. Schmid, K. Zeitler, R. M. Gschwind, *Angew. Chem.* **2010**, *122*, 5117–5123; *Angew. Chem. Int. Ed.* **2010**, *49*, 4997–5003; b) M. Wiesner, G. Upert, G. Angelici, H. Wennemers, *J. Am. Chem. Soc.* **2010**, *132*, 6–7; c) M. B. Schmid, K. Zeitler, R. M. Gschwind, *J. Org. Chem.* **2011**, *76*, 3005–3015; d) M. B. Schmid, K. Zeitler, R. M. Gschwind, *J. Am. Chem. Soc.* **2011**, *133*, 7065–7074; e) K. Patora-Komisarska, M. Benohoud, H. Ishikawa, D. Seebach, Y. Hayashi, *Helv. Chim. Acta* **2011**, *94*, 719–745; f) M. B. Schmid, K. Zeitler, R. M. Gschwind, *Chem. Sci.* **2011**, *2*, 1793–1803.
- [8] a) D. Seebach, U. Grošelj, D. M. Badine, W. B. Schweizer, A. K. Beck, *Helv. Chim. Acta* **2008**, *91*, 1999–2034; b) U. Grošelj, D. Seebach, D. M. Badine, W. B. Schweizer, A. K. Beck, I. Krossing, P. Klose, Y. Hayashi, T. Uchimaru, *Helv. Chim. Acta* **2009**, *92*, 1225–1259.
- [9] T. J. Peelen, Y. Chi, S. H. Gellman, *J. Am. Chem. Soc.* **2005**, *127*, 11598–11599.
- [10] This procedure is a modification of the method described in: K. L. Brown, L. Damm, J. D. Dunitz, A. Eschenmoser, R. Hobi, C. Kratky, *Helv. Chim. Acta* **1978**, *61*, 3108–3135.
- [11] CCDC 877482 (**3d**) and 877481 (**3e**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [12] For a detailed analysis of the dispersion interactions in α,β -unsaturated iminium ions derived from imidazolidinones, see: D. Seebach, U. Grošelj, W. B. Schweizer, S. Grimme, C. Mück-Lichtenfeld, *Helv. Chim. Acta* **2010**, *93*, 1–16.
- [13] Reviews on pyramidalization indices: a) F. K. Winkler, J. D. Dunitz, *J. Mol. Biol.* **1971**, *59*, 169–182; b) J. D. Dunitz, *X-Ray Analysis and Structure of Organic Molecules*, Cornell University Press, London, **1979**.
- [14] a) H. Mayr, M. Patz, *Angew. Chem.* **1994**, *106*, 990–1010; *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 938–957; b) H. Mayr, T. Bug, M. F. Gotta, N. Hering, B. Irrgang, B. Janker, B. Kempf, R. Loos, A. R. Ofial, G. Remennikov, H. Schimmel, *J. Am. Chem. Soc.* **2001**, *123*, 9500–9512; c) H. Mayr, B. Kempf, A. R. Ofial, *Acc. Chem. Res.* **2003**, *36*, 66–77; d) H. Mayr, A. R. Ofial, *Pure Appl. Chem.* **2005**, *77*, 1807–1821; e) H. Mayr, A. R. Ofial, *J. Phys. Org. Chem.* **2008**, *21*, 584–595; f) Database for N , s_N , and E parameters and references to their origin: <http://www.cup.uni-muenchen.de/oc/mayr/DBintro.html>.
- [15] a) G. Deslongchamps, P. Deslongchamps, *Org. Biomol. Chem.* **2011**, *9*, 5321–5333; b) see also discussion in Ref. [10].
- [16] Y. Hayashi, H. Gotoh, T. Hayashi, M. Shoji, *Angew. Chem.* **2005**, *117*, 4284–4287; *Angew. Chem. Int. Ed.* **2005**, *44*, 4212–4215.
- [17] I. Zenz, H. Mayr, *J. Org. Chem.* **2011**, *76*, 9370–9378.
- [18] a) J. Burés, A. Armstrong, D. G. Blackmond, *J. Am. Chem. Soc.* **2011**, *133*, 8822–8825; b) J. Burés, A. Armstrong, D. G. Blackmond, *J. Am. Chem. Soc.* **2012**, *134*, 6741–6750.
- [19] I. Ibrahim, A. Córdova, *Chem. Commun.* **2006**, 1760–1762.