

H<sub>2</sub> Activation

# Hydrogen Activation by an Intramolecular Boron Lewis Acid/ Zirconocene Pair\*\*


Santhosh Kumar Podiyanachari, Roland Fröhlich, Constantin G. Daniliuc, Jeffrey L. Petersen, Christian Mück-Lichtenfeld, Gerald Kehr, and Gerhard Erker\*

Small-molecule activation is a very important feature in stoichiometric synthetic chemistry as well as in catalysis. The cleavage and activation of dihydrogen is a typical example.<sup>[1]</sup> In many molecular systems hydrogen activation is carried out by oxidative addition of the H<sub>2</sub> molecule to a single transition-metal center, the Wilkinson catalyst being a typical example.<sup>[2]</sup> Alternatively transition-metal centers can split dihydrogen heterolytically with the assistance of a proton accepting main-group-element Lewis base, variants of the Noyori catalysts being typical examples.<sup>[3]</sup> We have now prepared a system featuring an inverse pair of functionalities, namely a bifunctional Group 4 metallocene/borane system where the main-group-element component may serve as a hydride acceptor. Herein we describe the synthesis of this new zirconocene/RB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> pair and its reaction with dihydrogen under mild conditions.

We first treated the aldimido zirconocene complex **1**<sup>[4]</sup> with trimethylsilyl ethynyl lithium. Product **2** (see Scheme 1) was isolated in over 90% yield as a yellow solid. It was characterized by X-ray diffraction (see the Supporting Information) and shown to contain a close to linear “metalla-2-aza allene” type unit.

Subsequent treatment of compound **2** with one molar equivalent of “Piers’ borane” [HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>]<sup>[5]</sup> gave complex **4** (isolated in over 80% yield; see Scheme 1 and Figure 1). Apparently, compound **4** is the product of a 1,1-hydroboration reaction<sup>[6,7]</sup> of HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> with the pendant trimethylsilyl acetylide  $\sigma$ -ligand. This reaction is typically initiated by  $\sigma$ -alkynyl ligand transfer from zirconium to the boron Lewis acid.<sup>[7]</sup> Hydride shift from boron to carbon followed by coordination of the imino nitrogen atom to the boron Lewis acid then straightforwardly leads to **4**.

We then investigated the formally related, but in some essential aspects decidedly different reaction of the  $\sigma$ -alkenyl/



Scheme 1. Preparation of complexes **2** and **4**; Cp = C<sub>5</sub>H<sub>5</sub>.




Figure 1. Molecular structure of compound **4**. Selected bond lengths [Å] and angles [°]: Zr1–C1 2.165(4), Zr1–C2 2.513(4), Zr1–N1 2.277(3), C1–C2 1.338(5), B1–C2 1.693(5), Si1–C1 1.872(4), N1–C3 1.269(4), B1–N1 1.578(5); C1–Zr1–N1 101.8(1), B1–N1–Zr1 93.2(2), C1–Zr1–C2 32.1(1), N1–Zr1–C2 69.7(1), C2–C1–Si1 121.5(3), C2–C1–Zr1 88.4(2), B1–C2–Zr1 82.5(2).<sup>[19]</sup>

[\*] Dr. S. K. Podiyanachari, Dr. R. Fröhlich,<sup>[†]</sup> Dr. C. G. Daniliuc,<sup>[†]</sup> Dr. C. Mück-Lichtenfeld,<sup>[††]</sup> Dr. G. Kehr, Prof. Dr. G. Erker  
Organisch-Chemisches Institut, Westfälische Wilhelms-Universität  
Correnstrasse 40, 48149 Münster (Germany)  
E-mail: erker@uniso-muenster.de

Prof. Dr. J. L. Petersen<sup>[†]</sup>  
Bennett Department of Chemistry, West Virginia University  
P.O. Box 6045, Morgantown, WV 26506 (USA)


[†] X-ray crystal structure analyses

[††] Computational chemistry


[\*\*] Financial support from the Deutsche Forschungsgemeinschaft is gratefully acknowledged.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201202218>.

$\sigma$ -alkynyl zirconocene complex **5** with HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>.<sup>[8]</sup> Complex **5** was prepared by hydrozirconation of *tert*-butylacetylene<sup>[9]</sup> followed by treatment with the Li–C≡C–SiMe<sub>3</sub> reagent (see Scheme 2). It was then treated with one molar equivalent of HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> in benzene at room temperature (12 h). Workup gave the product **7** as a dark red solid in 86% yield. The X-ray crystal structure analysis revealed that one equivalent of *tert*-butylethene had apparently been eliminated during the reaction leaving the substituted 2-boryl-1-silylacetylene zirconocene complex **7** behind (see Figure 2). Both the acetylenic carbon atoms are strongly bonded to zirconium (Zr1–C1: 2.275(2), Zr1–C2: 2.390(2), C1–C2: 1.281(3) Å). The C1–



**Scheme 2.** Synthesis of compound **7**.



**Figure 2.** A view of the molecular structure of compound **7** featuring a weak Zr...F contact. Selected bond lengths [Å] and angles [°]: Zr1–C1 2.275(2), Zr1–C2 2.390(2), Si1–C1 1.857(2), Zr1–F36 2.512(1), C1–C2 1.281(3), C2–B1 1.455(3); C1–Zr1–C2 31.7(6), C1–Zr1–F36 121.5(5), B1–C2–Zr1 94.6(1).<sup>[19]</sup>


Si1 vector is bent away from the zirconocene unit (angle C2–C1–Si1: 136.3(2)°). Most remarkable is that the C2–B1 vector is slightly leaning toward the zirconocene moiety (angle C1–C2–B1: 162.61(2)°). A close inspection of the coordination sphere of the central zirconium atom revealed a weak interaction with a fluorine atom of one of the C<sub>6</sub>F<sub>5</sub> groups at boron Zr1–F36: 2.512(1) Å.<sup>[10]</sup> The Zr1...B1 separation is 2.896(2) Å.

In solution complex **7** features <sup>1</sup>H and <sup>13</sup>C NMR signals of the Cp<sub>2</sub>Zr core at δ = 5.42 (s, 10H) ppm and δ = 110.7 ppm, respectively. The acetylene ligand <sup>13</sup>C NMR resonances occur at δ = 173.9 (≡C[Si]) and δ = 178.6 ppm ([B]C≡). Compound **7** shows a <sup>11</sup>B NMR resonance at δ = 15.5 ppm and the <sup>29</sup>Si NMR signal of the –SiMe<sub>3</sub> group at δ = –6.0 ppm (with a corresponding <sup>1</sup>H NMR singlet at δ = 0.16 ppm). In [D<sub>8</sub>]THF solution the spectrum of compound **7** features three sharp <sup>19</sup>F NMR signals at δ = –130.2 (4F, *ortho*), –160.9 (2F, *para*), –166.3 ppm (4F, *meta*) for the C<sub>6</sub>F<sub>5</sub> substituents at boron. In the non-coordinating solvent [D<sub>8</sub>]toluene we observed decoalescence of the averaged *o*-C<sub>6</sub>F<sub>5</sub> <sup>19</sup>F NMR resonance at δ = –145.2 ppm (4F, 363 K) to a broad “normal” *o*-C<sub>6</sub>F<sub>5</sub> signal at δ = –126.7 ppm (2F, 233 K), and a broad *o*-

C<sub>6</sub>F<sub>5</sub> signal at δ = –167.1 ppm (2F, 233 K; the spectra are shown in the Supporting Information). The δ = –167.1 ppm resonance is an averaged <sup>19</sup>F NMR signal of the bridging Zr–F–C(Ar) with the remaining non-bridged *o*-F resonance of the one *o*-C<sub>6</sub>F<sub>5</sub> ring. Consequently, this signal starts to decoalesce upon further lowering the temperature without reaching a fully resolved <sup>19</sup>F spectrum (temperature limit 193 K). Finally, the pair of *m*-F atoms of the two C<sub>6</sub>F<sub>5</sub> rings at boron give rise to a pair of signals at δ = –158.7, –163.2 ppm (each 2F, 193 K).


We can safely assume that the initial step of the reaction of **5** with HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> is again acetylide transfer from zirconium to boron<sup>[7]</sup> (to generate **6**). In contrast to the nitrogen case described above (see Scheme 1) the intermediate **6** apparently disfavors entering into the 1,1-hydroboration route but chooses stabilization by hydride transfer and reductive coupling to eventually yield **7** and *tert*-butylethene (see Scheme 2).

Compound **7** reacted rapidly with dihydrogen at ambient conditions in benzene solution (1.5 bar H<sub>2</sub> pressure, RT, 15 min) to give the H<sub>2</sub> addition product **9** (see Scheme 3). The



**Scheme 3.** Preparation of the hydrogen-addition product **9**.

hydrido zirconocene/hydrido borate complex **9** was isolated as a pale yellow solid in over 80 % yield. The X-ray crystal structure analysis (Figure 3) shows the presence of a [Cp<sub>2</sub>ZrH<sup>+</sup>] cation moiety to which a [HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>]<sup>–</sup>–C≡C–SiMe<sub>3</sub> anion is π-coordinated. The resulting structure features a C1–C2 bond length of 1.265(5) Å (there are two chemically equivalent crystallographically independent molecules in the




**Figure 3.** A projection of the molecular structure of the zwitterionic hydrogen-addition product **9**. Selected bond lengths [Å] and angles [°]: Zr1–C1 2.516(4), Zr1–C2 2.327(3), Zr1–H1 1.86(4), Zr1–H01 2.04(3), B1–C2 1.554(5), B1–H01 1.24(3), C1–C2 1.265(5), C1–H1 1.54(4); C1–Zr1–C2 29.9(1), C1–Zr1–H1 37.5(1), C2–Zr1–H01 60.3(8), C2–C1–Si1 143.2(3), C2–C1–Zr1 66.7(2), Si1–C1–Zr1 149.4(2).<sup>[19]</sup>

crystal; the values of molecule **A** are given here) which is slightly shorter than found in its precursor **7** (see above). In **9** the C1–Si1 vector is markedly pointing away from the zirconocene unit (angle C2–C1–Si1: 143.2(3)°) and there seems to be a weak interaction of the zirconium-hydride with C1. The C2–B1 vector is slightly oriented toward the metallocene (angle C1–C2–B1: 169.8(3)°) and the hydride at B is oriented toward the Zr atom; it is probably unsymmetrically bridging. Complex **9** shows a pronouncedly unsymmetrical coordination of the substituted acetylene ligand to the [Cp<sub>2</sub>ZrH<sup>+</sup>] moiety with a short Zr1–C2 bond (2.327(3) Å) and a markedly longer Zr1–C1 linkage (2.516 (4) Å), as is often observed in related alkene or alkyne units coordinated to alkyl zirconocene cations.<sup>[11]</sup>


In solution complex **9** shows a <sup>1</sup>H NMR Cp singlet at δ = 4.94 (<sup>13</sup>C: δ = 104.1 ppm). The B–H–Zr hydrogen atom was detected at δ = –2.29 ppm, the Zr–H–C hydrogen atom at δ = 3.80 ppm (d, <sup>2</sup>J<sub>HH</sub> = 5.5 Hz).<sup>[12,13]</sup> We observe the large chemical shift difference of the alkyne ligand carbon NMR signals ([Si]C≡: δ = 79.8 ppm (<sup>1</sup>J<sub>CH</sub> ≈ 44 Hz); ≡C[B]: δ = 151.4 ppm) that is typical for the unsymmetrical bonding situation of an acetylene ligand π-coordinated to a [Cp<sub>2</sub>ZrR<sup>+</sup>] cation moiety.<sup>[11]</sup> We also detect a single set of three <sup>19</sup>F NMR signals at δ = –129.5 (o), –157.2 (p), and –162.9 ppm (m) with a small δ<sub>p,m</sub> shift difference [<sup>11</sup>B NMR: δ = –22.8 ppm (d, <sup>1</sup>J<sub>BH</sub> ≈ 60 Hz)].

Our study shows that the borane HB(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> readily and selectively abstracts a σ-acetylide ligand from a zirconocene complex. The resulting zwitterionic intermediate then has several competing choices for stabilization. In the imido zirconocene case (**3**) the rarely observed 1,1-hydroboration reaction is preferred, whereas in the alkenyl zirconocene analogue (**6**) hydride transfer to zirconium followed by reductive coupling prevails to yield the unique product **7**. Although π-alkene and π-alkyne zirconocenes are known to sometimes exhibit a pronounced metallacyclopentene σ-complex character,<sup>[14]</sup> the system **7** formally may be regarded as a Cp<sub>2</sub>Zr<sup>II</sup> derivative. Two pathways for the reaction of **7** with dihydrogen can thus be formulated. Pathway (b) in Scheme 4 describes a “conventional stepwise route” where both the zirconocene and the borane functionalities act separately in two consecutive steps, namely oxidative H–H addition to zirconium (to give **11**) followed by hydride abstraction by the borane to generate **8**.

DFT calculations (TPSS-D3/def2-TZVP)<sup>[15]</sup> have been performed on the intermediates and the transition structure of H<sub>2</sub> cleavage in Scheme 4. We have tried to identify a product (**11**) of oxidative H<sub>2</sub> addition to the zirconocene complex **7** (pathway (b)) without success. However, pathway (a) in Scheme 4 has been found to be a facile and exothermic process (Figure 4 and Table 1): Side-on coordination of H<sub>2</sub> at the metal center is enthalpically favored by –6.1 kcal mol<sup>–1</sup> at 298 K. Coordination of H<sub>2</sub> at the side of the boron atom (**10a**) is preferred over the alternative coordination at the silyl-substituted side (**10b**, –0.8 kcal mol<sup>–1</sup>). The transition structure of H<sub>2</sub> cleavage (see Figure 4) does not require extensive geometrical changes of **10a** except the elongation of the H–H bond to 1.043 Å, accompanied by an almost vanishing enthalpic barrier (ΔH<sup>‡</sup>(298 K) = 0 kcal



**Scheme 4.** Mechanistic pathways for the formation of zwitterionic compound **9**.



**Figure 4.** Calculated transition-state structure of the **7** + H<sub>2</sub> → **8** reaction.

mol<sup>–1</sup>, ΔH<sup>‡</sup>(220 K) = 0.1 kcal mol<sup>–1</sup>) of the activation. Rearrangement of the zwitterionic intermediate **8** to its isomer **9b** is exothermic again by –4.7 kcal mol<sup>–1</sup> (at 298 K). Interestingly, our DFT optimizations in the gas phase identified two isomers of **9**: Besides the structure found in the solid state (**9b**), a second, slightly more stable isomer exhibits a long H1–Zr1 and a shorter C1–H1 bond (**9a**) [see the Supporting Information].


**Table 1:** DFT calculated (TPSS-D3/def2-TZVP) relative energies (in kcal mol<sup>–1</sup>) of intermediates and the H<sub>2</sub> cleavage transition structure in the formation of compound **9** from **7**.<sup>[a]</sup>

| Structure         | ΔE    | ΔH(298 K) <sup>[b]</sup> | ΔH(220 K) <sup>[b]</sup> |
|-------------------|-------|--------------------------|--------------------------|
| <b>10a</b>        | –9.8  | –6.1                     | –5.9                     |
| <b>10b</b>        | –4.9  | –0.8                     | –0.7                     |
| <b>TS(10a-8)</b>  | –8.8  | –6.1                     | –5.8                     |
| <b>8</b>          | –12.3 | –8.4                     | –8.1                     |
| <b>9a</b>         | –17.6 | –13.3                    | –12.9                    |
| <b>9b (= “9”)</b> | –17.0 | –13.1                    | –12.8                    |

[a] Referenced to the sum of energies of complex **7** and H<sub>2</sub>. [b] Includes ZPVE, H<sub>vib</sub>, H<sub>rot</sub>, H<sub>trans</sub> at the given temperature and 1 atm.

It seems that the synergistic oxidative addition pathway (a)<sup>[16]</sup> involving both the formally low-oxidation state zirconium and the borane Lewis acid in the actual step of heterolytic cleavage of the H–H molecule leading to **8** (and subsequently **9**) is favored over the conventional stepwise mechanism pathway (b).

It is interesting to note that the heterolytic splitting of dihydrogen by a variety of “frustrated Lewis pairs”,<sup>[17]</sup> such as the recently reported very reactive hydrogen activating system **12** (see Scheme 5),<sup>[18]</sup> resembles the synergistic pathway (a) in Scheme 4 conceptually. It seems that currently



**Scheme 5.** Formation of zwitterionic compound **13**.

examples are increasingly found where transition-metal complexes and main-group-element-derived frustrated Lewis pairs show some remarkably similar chemical behavior.

Received: March 20, 2012

Revised: May 22, 2012

Published online: July 30, 2012

**Keywords:** boron · frustrated Lewis pairs · hydrogen activation · oxidative addition · zirconium

- [1] a) J. G. de Vries, C. J. Elsevier, *The Handbook of Homogeneous Hydrogenation* (Eds.: J. G. de Vries, C. J. Elsevier), Wiley-VCH, Weinheim, **2007**, and references therein.
- [2] J. A. Osborn, F. H. Jardine, J. F. Young, G. Wilkinson, *J. Chem. Soc. A* **1966**, 1711–1732.
- [3] a) K.-J. Haack, S. Hashiguchi, A. Fujii, T. Ikariya, R. Noyori, *Angew. Chem.* **1997**, *109*, 297–300; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 285–288; b) S. Hashiguchi, A. Fujii, K.-J. Haack, K. Matsumura, T. Ikariya, R. Noyori, *Angew. Chem.* **1997**, *109*, 300–303; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 288–290; c) H. Doucet, T. Ohkuma, K. Murata, T. Yokozawa, M. Kozawa, E. Katayama, A. F. England, T. Ikariya, R. Noyori, *Angew. Chem.* **1998**, *110*, 1792–1796; *Angew. Chem. Int. Ed.* **1998**, *37*, 1703–1707.
- [4] G. Erker, W. Frömberg, J. L. Atwood, W. E. Hunter, *Angew. Chem.* **1984**, *96*, 72–73; *Angew. Chem. Int. Ed. Engl.* **1984**, *23*, 68–69.
- [5] a) D. J. Parks, R. E. von H. Spence, W. E. Piers, *Angew. Chem.* **1995**, *107*, 895–897; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 809–811; b) R. E. von H. Spence, D. J. Parks, W. E. Piers, M.-A. MacDonald, M. J. Zaworotko, S. J. Rettig, *Angew. Chem.* **1995**, *107*, 1337–1340; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1230–1233; c) W. E. Piers, T. Chivers, *Chem. Soc. Rev.* **1997**, *26*, 345–354; d) R. E. von H. Spence, W. E. Piers, Y. Sun, M. Parvez, L. R. MacGillivray, M. J. Zaworotko, *Organometallics* **1998**, *17*, 2459–2469; e) D. J. Parks, W. E. Piers, G. P. A. Yap, *Organometallics* **1998**, *17*, 5492–5503.
- [6] a) J. Ugoletti, G. Dierker, G. Kehr, R. Fröhlich, S. Grimme, G. Erker, *Angew. Chem.* **2008**, *120*, 2662–2665; *Angew. Chem. Int. Ed.* **2008**, *47*, 2622–2625; b) J. Ugoletti, G. Kehr, R. Fröhlich, G. Erker, *Chem. Commun.* **2010**, *46*, 3016–3018; c) J. Ugoletti, G. Dierker, R. Fröhlich, G. Kehr, G. Erker, *J. Organomet. Chem.* **2011**, *696*, 1184–1188.
- [7] See also: a) B. Wrackmeyer, *Coord. Chem. Rev.* **1995**, *145*, 125–156; b) B. Wrackmeyer, G. Kehr, S. Willbold, *J. Organomet. Chem.* **1999**, *590*, 93–103; c) C. Chen, T. Voss, R. Fröhlich, G. Kehr, G. Erker, *Org. Lett.* **2011**, *13*, 62–65; d) C. Chen, G. Kehr, R. Fröhlich, G. Erker, *J. Am. Chem. Soc.* **2010**, *132*, 13594–13595; e) C. Chen, F. Eweiner, B. Wibbeling, R. Fröhlich, S. Senda, Y. Ohki, K. Tatsumi, S. Grimme, G. Kehr, G. Erker, *Chem. Asian J.* **2010**, *5*, 2199–2208.
- [8] G. Erker, G. Kehr, R. Fröhlich, *Adv. Organomet. Chem.* **2004**, *51*, 109–162.
- [9] J. Schwartz, J. A. Labinger, *Angew. Chem.* **1976**, *88*, 402–409; *Angew. Chem. Int. Ed. Engl.* **1976**, *15*, 333–340.
- [10] a) B. Temme, G. Erker, J. Karl, H. Luftmann, R. Fröhlich, S. Kotila, *Angew. Chem.* **1995**, *107*, 1867–1869; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 1755–1757; b) J. Karl, G. Erker, R. Fröhlich, *J. Am. Chem. Soc.* **1997**, *119*, 11165–11173; c) M. Dahlmann, G. Erker, R. Fröhlich, O. Meyer, *Organometallics* **2000**, *19*, 2956–2967; d) N. Kleigrewe, T. Brackmeyer, G. Kehr, R. Fröhlich, G. Erker, *Organometallics* **2001**, *20*, 1952–1955.
- [11] a) A. S. Guam, R. F. Jordan, *Organometallics* **1991**, *10*, 3470–3479; b) A. S. Guam, R. F. Jordan, *J. Org. Chem.* **1992**, *57*, 5994–5999; c) E. J. Stobenau III, R. F. Jordan, *J. Am. Chem. Soc.* **2006**, *128*, 8162–8175.
- [12] See for a comparison: P. Arndt, W. Baumann, A. Spannenberg, U. Rosenthal, V. V. Burlakov, V. B. Shur, *Angew. Chem.* **2003**, *115*, 1455–1458; *Angew. Chem. Int. Ed.* **2003**, *42*, 1414–1418.
- [13] See also: a) A. Ohff, P. Kosse, W. Baumann, A. Tillack, R. Kempe, H. Görls, V. V. Burlakov, U. Rosenthal, *J. Am. Chem. Soc.* **1995**, *117*, 10399–10400; b) M.-F. Fan, Z. Lin, *Organometallics* **1997**, *16*, 494–496; c) N. Peulecke, A. Ohff, P. Kosse, A. Tillack, A. Spannenberg, R. Kempe, W. Baumann, V. V. Burlakov, U. Rosenthal, *Chem. Eur. J.* **1998**, *4*, 1852–1861.
- [14] a) U. Rosenthal, V. V. Burlakov, P. Arndt, W. Baumann, A. Spannenberg, *Organometallics* **2005**, *24*, 456–471; b) J. Ugoletti, G. Kehr, R. Fröhlich, S. Grimme, G. Erker, *J. Am. Chem. Soc.* **2009**, *131*, 1996–2007; c) E. D. Jemmis, S. Roy, V. V. Burlakov, H. Jiao, M. Klahn, S. Hansen, U. Rosenthal, *Organometallics* **2010**, *29*, 76–81.
- [15] a) J. Tao, J. P. Perdew, V. N. Staroverov, G. E. Scuseria, *Phys. Rev. Lett.* **2003**, *91*, 146401; b) F. Weigend, R. Ahlrichs, *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297–3305; c) S. Grimme, J. Antony, S. Ehrlich, H. Krieg, *J. Chem. Phys.* **2010**, *132*, 154104; d) S. Grimme, S. Ehrlich, L. Goerigk, *J. Comput. Chem.* **2011**, *32*, 1456–1465; e) TURBOMOLE V6.3 2011, a development of University of Karlsruhe and Forschungszentrum Karlsruhe GmbH, 1989–2007, TURBOMOLE GmbH, since 2007; available from <http://www.turbomole.com>.
- [16] See for a comparison: W. Hill Harman, J. C. Peters, *J. Am. Chem. Soc.* **2012**, *134*, 5080–5082, and references therein.
- [17] D. W. Stephan, G. Erker, *Angew. Chem.* **2010**, *122*, 50–81; *Angew. Chem. Int. Ed.* **2010**, *49*, 46–76.
- [18] a) P. Spies, G. Erker, G. Kehr, K. Bergander, R. Fröhlich, S. Grimme, D. W. Stephan, *Chem. Commun.* **2007**, 5072–5074; See also: b) P. Spies, S. Schwendemann, S. Lange, G. Kehr, R. Fröhlich, G. Erker, *Angew. Chem.* **2008**, *120*, 7654–7657; *Angew. Chem. Int. Ed.* **2008**, *47*, 7543–7546; c) B.-H. Xu, G. Kehr, R. Fröhlich, B. Wibbeling, B. Schirmer, S. Grimme, G. Erker, *Angew. Chem.* **2011**, *123*, 7321–7324; *Angew. Chem. Int. Ed.* **2011**, *50*, 7183–7186.
- [19] CCDC 888636 (1), 888637 (2), 888638 (4), 888639 (7), and 888640 (9) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).