

Enantioselective Total Synthesis of the Reported Structures of (–)-9-*epi*-Presilphiperfolan-1-ol and (–)-Presilphiperfolan-1-ol: Structural Confirmation and Reassignment and Biosynthetic Insights**

Allen Y. Hong and Brian M. Stoltz*

In memory of Michael W. Day

Presilphiperfolanols (also known as prebotrydials) are members of a family of natural products that are important biosynthetic precursors to diverse sesquiterpenoids (Figure 1).^[1–4] To date, four members of the group have been isolated: presilphiperfolan-1-ol (**1**),^[5] 9-*epi*-presilphiperfolan-1-ol (**2**),^[6] presilphiperfolan-8-ol (**3**),^[2] and presilphiperfolan-9-ol (**4**),^[7] with **1** and **2** being the most recently disclosed (Figure 1A).

The original proposal for the biosynthesis of the presilphiperfolane core and its rearrangement to related sesquiterpenes was reported independently by Bohlmann et al.^[1,2] and Hanson.^[3a] Subsequent studies by multiple research groups^[3b,4,8,9] have provided further insight into this biosynthetic pathway. Farnesyl pyrophosphate (FPP) is commonly believed to undergo enzyme-catalyzed cyclization to caryophyllenyl cation **6**,^[9] which undergoes further rearrangements to the C(9)- and C(8)-presilphiperfolanyl cations **7** and **8** (Figure 1B). Hydration of these cations affords the natural products **3** and **4** as single stereoisomers. To date, existing biosynthetic proposals have not accounted for the formation of **1** and **2**, which possess an unusual C(9)-epimeric relationship. Neither **3** nor **4** have known naturally occurring epimers. From the tricyclic cation **8**, further C–C bond migrations can lead to diverse sesquiterpenoid natural products (Figure 1C).^[1–4]




Figure 1. Presilphiperfolanyl (prebotrydial) cations as key intermediates in the biosynthesis of diverse sesquiterpenes.

In addition to their central biosynthetic role, the presilphiperfolanols possess antimycobacterial properties (e.g., **2**)^[10] and insect antifeedant properties (e.g., **4**).^[11] Furthermore, unnatural presilphiperfolane analogs have demonstrated antifungal activity.^[12]

Although the presilphiperfolanols have garnered broad scientific interest in the natural product,^[13] biosynthesis,^[3,4,9,12,14,15] organic synthesis,^[16,17] computation,^[8] and fragrance^[7a,16,18] literature since their discovery, they have proven to be challenging targets for total synthesis. To date, only (±)-presilphiperfolan-9-ol (**4**) has been prepared by total synthesis,^[16] and no asymmetric routes toward any of the presilphiperfolanols have been reported. Within this context,

[*] A. Y. Hong, Prof. B. M. Stoltz

Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology
1200 E. California Blvd, MC 101-20, Pasadena, CA 91125 (USA)
E-mail: stoltz@caltech.edu

[**] The authors thank NIH-NIGMS (R01GM080269-01), Roche, Abbott Laboratories, Amgen, Boehringer Ingelheim, the Gordon and Betty Moore Foundation, and Caltech for awards and financial support. Prof. Suzana G. Leitão (Universidade Federal do Rio de Janeiro) generously provided NMR spectra for natural **2**. Dr. Lawrence Henling is gratefully acknowledged for X-ray crystallographic structure determination. The Bruker KAPPA APEXII X-ray diffractometer used in this study was purchased through an NSF CRIF:MU award to Caltech (CHE-0639094). Prof. Sarah Reisman, Dr. Scott Virgil, Dr. Douglas C. Behenna, Robert J. Ely and Fang Gao (Boston College), and Jessica Y. Wu (Harvard University) are acknowledged for helpful discussions and suggestions. Dr. David VanderVelde and Dr. Scott Ross are acknowledged for NMR assistance.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201205276>.

our group became interested in developing an enantioselective synthesis to enable further exploration of their biosynthetic relationships and biological activity.

The structurally complex presilphiperfolanols are distinguished by their rare, compact tricyclic terpenoid core, which bears five contiguous stereocenters, two all-carbon quaternary centers, and a tertiary alcohol (Figure 1 A). In addition to these readily apparent structural features, considerable ring strain is built into the tricyclic system,^[4c,19] allowing these compounds to undergo thermodynamically favorable skeletal rearrangements leading to numerous terpenes (Figure 1 C).^[1–4] Our goal was to develop an efficient and general asymmetric route to access various members of the presilphiperfolanol family. Here, we describe the first asymmetric total syntheses of the reported structures of presilphiperfolanols **1** and **2**. Our investigation has confirmed the structure of **2** and prompted us to reassign the structure of **1**. Finally, in the context of this reassignment, we propose a new biosynthetic route to account for our observations.

Retrosynthetic analysis suggested that tricycle **9** could serve as an intermediate for the divergent synthesis of several members of the family (Figure 2). Two of the three rings could be forged simultaneously in an intramolecular [4+2] cyclo-





Figure 2. Synthetic strategy toward the presilphiperfolane core.

addition (via **10**), thereby avoiding a more conventional sequential annulation strategy. Monocyclic precursor **11** could be prepared from an α -quaternary vinylogous ester **12** by a two-carbon ring contraction. The strategic application of our recently reported methodology for the synthesis of γ -quaternary acylcyclopentenones and α -quaternary vinylogous esters would provide a solid starting point for our synthesis.^[20]

Our synthetic efforts began with acylation/alkylation of commercial 3-isobutoxycycloheptanone **14** using isoprenol-derived carbamate **15**^[21,22] and methyl iodide under basic conditions (Scheme 1). This improved protocol enabled efficient access to β -ketoester **13** in a single synthetic operation and avoided employing the corresponding cyanofomate.^[23] With our desired functionality installed, we were poised to evaluate our asymmetric Pd-catalyzed decarboxylative alkylation^[20,24] on this novel substrate type containing a 2-vinyl allyl fragment. To our delight, treatment of β -ketoester **13** with catalytic $\text{Pd}_2(\text{pmdba})_3$ and (*S*)-*t*Bu-PHOX^[25,26] in toluene proceeded smoothly on 10 g scale, affording isoprenylated vinylogous ester **12** in 91 % yield and 95 % *ee*.^[27,28] Subsequent LiAlH_4 reduction afforded an intermediate β -hydroxyketone, which underwent a base-




Scheme 1. Gram-scale synthesis of acylcyclopentene **11** and IMDA. HMDS = hexamethyldisilazane, pmdba = 4,4'-methoxydibenzylideneacetone, TFE = 2,2,2-trifluoroethanol, TBS = *tert*-butyl dimethylsilyl, NOE = nuclear Overhauser effect.

mediated two-carbon ring contraction to provide multigram quantities of acylcyclopentene **11** in 92 % yield over two steps.^[20] After straightforward silyl dienol ether formation, we were able to examine our bicyclization strategy (via **10a**: R = OTBS, R' = CH₂). The intramolecular Diels–Alder (IMDA) reaction^[29] proceeded efficiently with microwave irradiation despite the lack of an activated dienophile,^[30] affording tricycle **16** in 80 % yield over two steps as a single diastereomer. The *cis* relationship of the C(7) and C(8) hydrogens was determined by NOESY experiments on cycloadduct **16**. Unfortunately, in order to proceed toward targets **1** and **2**, a *trans* relationship was required.

With a general and concise strategy to the tricyclic core, we next pursued the installation of the naturally occurring substituents and stereochemistry present in the strained presilphiperfolanols. Our revised strategy aimed to introduce the *gem*-dimethyl group at an earlier stage to potentially improve IMDA diastereoselectivity toward the desired *trans* product (via acylcyclopentene **17**, Scheme 2). Inspection of the possible IMDA transition states of the silyl dienol ether **10b** suggested that nonbonding interactions from the C(6) steric bulk could help favor the desired C(7) stereochemistry (Scheme 2B).^[31] In **TS-A**, the steric elements would be oriented away from the bond-forming centers while cycloaddition through **TS-B** could only proceed with severe steric clash between the cyclopentene ring and the *gem*-dimethyl functionality. Additionally, the new *gem*-dimethyl groups should decrease the C(5)–C(6)–C(7) bond angle while also providing greater conformational bias^[32] for the desired cyclization with this uncommon IMDA substrate type.^[33]

Our efforts toward this end commenced with a regioselective 1,4-hydroboration/oxidation of the sterically hindered 2-substituted 1,3-diene **18** after carbonyl protection of **11** (Scheme 2 A). While Pd,^[34a] Fe,^[34b] and Ni^[34c,d] catalysts are known to effect this transformation, literature precedent suggested that the regioselective formation of linear product **19a** would be most favorable with iron catalysis. To our surprise, the Ritter $\text{FeCl}_2(\text{py-imine})$ catalyst system provided a disappointing 1.1:1 ratio of linear:branched products.^[34b,35]




Scheme 2. Revised synthetic approach to presilpiperfolane core. cod = cyclooctadiene, DMAP = 4-dimethylaminopyridine, DMDO = dimethyldioxirane, TBAF = tetrabutylammonium fluoride.

Investigation of Suzuki–Miyaura conditions with $\text{Pd}(\text{PPh}_3)_4$ and HBPIn did not provide any reactivity.^[34a,35] Finally, we turned to Morken's $\text{Ni}(\text{cod})_2/\text{PCy}_3$ catalyst system.^[34c,35] Gratifyingly, we were able to obtain the separable allylic alcohols **19a** and **19b** in 81 % combined yield in a 3.5:1 ratio (linear:branched) on multigram scale with 5 mol % catalyst loading. Doubling the catalyst loading led to a decrease in reaction time and a small improvement to 88 % combined yield. To our knowledge, this is the first application of a regioselective metal-catalyzed 1,4-hydroboration in total synthesis.

Subsequent phosphorylation and regioselective Cu-catalyzed allylic substitution^[36] provided the requisite *gem*-dimethyl acylcyclopentene **17** in 80 % yield over two steps (Scheme 2B). After silyl dienol ether formation, the IMDA bicyclization proceeded smoothly under thermal or microwave-assisted conditions to provide a mixture of inseparable diastereomers **20a** and **20b**. A highly diastereoselective Rubottom oxidation of these products with DMDO^[37] provided α -hydroxyketones **21a** and **21b** in a 1:2 ratio as determined by ^1H NMR spectroscopy. Chromatographic separation gave **21a** in 27 % yield and **21b** in 61 % yield over three steps. The structure of tricycle **21b** was confirmed by single-crystal X-ray analysis.^[38]

With α -hydroxyketone **21a** in hand, Wittig methylenation proceeded efficiently to give allylic alcohol **22** in 90 % yield (Scheme 3). While PtO_2 -catalyzed hydrogenation of olefin **22** afforded a separable mixture of targets **1** and **2**, a diastereo-



Scheme 3. Synthesis of the reported structures of 9-*epi*-presilpiperfolan-1-ol (**2**) and presilpiperfolan-1-ol (**1**).

selective synthesis of both target molecules could also be achieved. Tertiary alcohol-directed hydrogenation with Crabtree's catalyst^[39] produced the reported structure of presilpiperfolan-1-ol (**1**) in 92 % yield and the structure was verified by single-crystal X-ray diffraction.^[38] Alternatively, efficient formation of the reported structure of 9-*epi*-presilpiperfolan-1-ol (**2**) was accomplished in 88 % yield over three steps by silylation^[40] of tertiary allylic alcohol **22** followed by PtO_2 -catalyzed hydrogenation and desilylation with TBAF.

Upon completing the synthesis of **1** and **2**, we compared our spectral data with the reported data for the isolated compounds. Although the ^1H and ^{13}C NMR spectra of synthetic 9-*epi*-presilpiperfolan-1-ol (**2**) were in excellent agreement with reported data, the spectra for synthetic presilpiperfolan-1-ol (**1**) clearly were not.^[35] In particular, the C(15) methyl hydrogens of synthetic **1** showed a ^1H NMR resonance at 0.94 ppm compared the corresponding resonance of reported **1**^[5] at 0.89 ppm. From the work of Joseph-Nathan and Leitão, the structure of reported 9-*epi*-presilpiperfolan-1-ol (**2**) was unambiguously established by X-ray crystallography.^[6b] The structure of reported presilpiperfolan-1-ol (**1**) isolated by König was assigned based on NMR data.^[5,41,42] Overall, the spectral data of reported **1** more closely matches synthetic and reported **2**^[6a] than synthetic **1**.

Given the significant discrepancy between our data and König's data for **1**, we sought to rationalize the formation of both **1** and **2** in the context of proposed mechanisms for presilpiperfolanol biosynthesis.^[1–4,8] While no proposals for the formation of **1** and **2** have been published, it is reasonable that presilpiperfolanols **1–4** are commonly derived from caryophyllenyl cation **6** (Figure 1B). Upon rearrangement to C(9)-presilpiperfolanyl cation **7**, a divergent pathway can be envisioned (Figure 3). A simple 1,2-*syn* hydride migration to tertiary C(1) cation **23** followed by hydration leads to 9-*epi*-presilpiperfolan-1-ol (**2**) with 9 β -methyl stereochemistry without invoking the intermediacy of unfavorable secondary carbocations.^[8] In contrast to this pathway, the formation of 9 α -methyl-oriented presilpiperfolan-1-ol (**1**) seems unlikely since there is no obvious pathway from C(9) cation **7** to C(1) cation **24**. Based on our NMR spectral data, X-ray crystal structure for **2**, and new biosynthetic proposals, we believe the reported structure of natural presilpiperfolan-1-ol (**1**) is




Figure 3. Biosynthetic proposal for the formation of **2**, and structural revision of **1**.

misassigned at C(9) and most likely is not a natural product. The nearly identical chemical shifts of reported **1** and **2** (^1H and ^{13}C NMR) suggest that the true structure of presilphiperfolan-1-ol (**1**) may actually be the same as 9-*epi*-presilphiperfolan-1-ol (**2**) with the 9β -methyl configuration (Figure 3). Since the reassignment can potentially lead to confusion, we suggest that the natural product structure represented by **2** be referred to as presilphiperfolan-1-ol or the more descriptive name 9β -presilphiperfolan-1 α -ol in the future discussions.

In summary, the first total syntheses of the reported structures of presilphiperfolanols **1** and **2** have been achieved in 13–15 steps and 7.9–8.6% overall yield from commercially available vinyllogous ester **14**. This work constitutes the first asymmetric total synthesis of any member of the presilphiperfolanol family. In addition to demonstrating the synthetic utility of functionality-rich acylcyclopentenones **11** and **17** generated from our Pd-catalyzed allylic alkylation/ring contraction methodology, we exploited a range of methods to concisely construct the fully substituted carbocyclic core of our targets. Analysis of spectral data and evaluation of likely presilphiperfolanol biosynthetic pathways prompted us to reassign the structure of reported **1** to **2**. Our synthetic route provides access to **1**, **2**, and synthetic analogs, enabling investigation of their biological activity. Extension of the synthetic strategy to the remaining members of the family, investigations of the biosynthetic pathway, and biomimetic rearrangement to other natural products are currently in progress.

Received: July 4, 2012

Published online: August 22, 2012

Keywords: asymmetric catalysis · natural products · structure determination · terpenoids · total synthesis

[1] F. Bohlmann, J. Jakupovic, *Phytochemistry* **1980**, *19*, 259–265.

- [2] F. Bohlmann, C. Zdero, J. Jakupovic, H. Robinson, R. M. King, *Phytochemistry* **1981**, *20*, 2239–2244.
- [3] For representative investigations of presilphiperfolane biosynthesis by isotopic incorporation studies, see the following and references therein: a) J. R. Hanson, *Pure Appl. Chem.* **1981**, *53*, 1155–1162; b) C.-M. Wang, R. Hopson, X. Lin, D. E. Cane, *J. Am. Chem. Soc.* **2009**, *131*, 8360–8361.
- [4] For reports describing the rearrangement of presilphiperfolanols to related sesquiterpene natural products, see: a) R. M. Coates, Z. Ho, M. Klobus, S. R. Wilson, *J. Am. Chem. Soc.* **1996**, *118*, 9249–9254; b) P. Weyerstahl, H. Marschall, I. Seelmann, J. Jakupovic, *Eur. J. Org. Chem.* **1998**, 1205–1212; c) C. E. Davis, B. C. Duffy, R. M. Coates, *J. Org. Chem.* **2003**, *68*, 6935–6943.
- [5] a) S. Melching, W. A. König, *Phytochemistry* **1999**, *51*, 517–523; b) “Isolierung, Strukturaufklärung und stereochemische Untersuchungen neuer sesquiterpenoider Verbindungen aus vier Chemotypen des Lebermooses *Conocephalum conicum*”: S. Melching, Ph.D. Thesis, Universität Hamburg, April **1999**.
- [6] a) S. C. Pinto, G. G. Leitão, H. R. Bizzo, N. Martinez, E. Dellacassa, F. M. dos Santos, Jr., F. L. P. Costa, M. B. de Amorim, S. G. Leitão, *Tetrahedron Lett.* **2009**, *50*, 4785–4787; b) P. Joseph-Nathan, S. G. Leitão, S. C. Pinto, G. G. Leitão, H. R. Bizzo, F. L. P. Costa, M. B. de Amorim, N. Martinez, E. Dellacassa, A. Hernández-Barragán, N. Pérez-Hernández, *Tetrahedron Lett.* **2010**, *51*, 1963–1965.
- [7] a) P. Weyerstahl in *Newer Trends in Essential Oils and Flavours* (Eds.: K. L. Dhar, R. K. Thappa, S. G. Agarwal), Tata McGraw-Hill, New Delhi, **1993**, pp. 24–41; b) J. A. Marco, J. F. Sanz-Cervera, M. D. Morante, V. Garcia-Lliso, J. Vallès-Xirau, J. Jakupovic, *Phytochemistry* **1996**, *41*, 837–844.
- [8] For computational studies exploring the presilphiperfolanol biosynthetic pathway, see: a) S. C. Wang, D. J. Tantillo, *Org. Lett.* **2008**, *10*, 4827–4830; b) J. E. Barquera-Lozada, G. Cuevas, *J. Org. Chem.* **2011**, *76*, 1572–1577.
- [9] For studies describing presilphiperfolane biosynthetic gene clusters, see: C. Pinedo, C.-M. Wang, J.-M. Pradier, B. Dalmais, M. Choquer, P. Le Pêcheur, G. Morgant, I. G. Collado, D. E. Cane, M. Viaud, *ACS Chem. Biol.* **2008**, *3*, 791–801.
- [10] S. C. Pinto, G. G. Leitão, D. R. de Oliveira, H. R. Bizzo, D. F. Ramos, T. S. Coelho, P. E. A. Silva, M. C. S. Lourenço, S. G. Leitão, *Nat. Prod. Commun.* **2009**, *4*, 1675–1678.
- [11] A. González-Coloma, F. Valencia, N. Martín, J. F. Hoffmann, L. Hutter, J. A. Marco, M. Reina, *J. Chem. Ecol.* **2002**, *28*, 117–129.
- [12] a) I. G. Collado, J. Aleu, A. J. Macías-Sánchez, R. Hernández-Galán, *J. Chem. Ecol.* **1994**, *20*, 2631–2644; b) I. G. Collado, J. Aleu, A. J. Macías-Sánchez, R. Hernández-Galán, *J. Nat. Prod.* **1994**, *57*, 738–746.
- [13] Presilphiperfolanol-derived natural products with oxidized core structures have been discovered. See: a) F. Bohlmann, J. Ziesche, R. K. Gupta, *Phytochemistry* **1982**, *21*, 1331–1334; b) A. H. Mericli, F. Mericli, J. Jakupovic, F. Bohlmann, X. A. Dominguez, H. S. Vega, *Phytochemistry* **1989**, *28*, 1149–1153; c) J.-L. Yang, L.-L. Liu, Y.-P. Shi, *Tetrahedron Lett.* **2009**, *50*, 6315–6317.
- [14] To our knowledge, the rearrangement of β -caryophyllene or β -caryophyllene derivatives in biomimetic reactions has not produced any of the presilphiperfolanol natural products. For representative studies, see the following and references therein: a) L. Fitjer, A. Malich, C. Paschke, S. Kluge, R. Gerke, B. Rissom, J. Weiser, M. Noltemeyer, *J. Am. Chem. Soc.* **1995**, *117*, 9180–9189; b) S. Shankar, R. M. Coates, *J. Org. Chem.* **1998**, *63*, 9177–9182.
- [15] The rearrangement of isocaryophyllene with Brønsted or Lewis acids has yielded unnatural isomeric presilphiperfolanol derivatives. For representative studies, see the following and references therein: a) K. Gollnick, G. Schade, A. F. Cameron, C. Hannaway, J. S. Roberts, J. M. Robertson, *J. Chem. Soc. Chem.*

- Commun.* **1970**, 248–249; b) T. M. Khomenko, I. Y. Bagryan-skaya, Y. V. Gatilov, D. V. Korchagina, V. P. Gatilova, Z. V. Dubovenko, V. A. Barkhash, *Zh. Org. Khim.* **1985**, *21*, 677–678. *Russ. J. Org. Chem. (Engl. Transl.)* **1985**, *21*, 614–615; c) also see ref. [12].
- [16] P. Weyerstahl, H. Marschall, M. Schulze, I. Schwöpe, *Liebigs Ann.* **1996**, 799–807.
- [17] “Carbocycle Construction in Terpenoid Synthesis. The Total Synthesis of (±)-Sarcodonin G and (±)-1-Epi-9-Norpresilphiperfolan-9-one”: M. W. Gilbert, Ph.D. Thesis, University of British Columbia, June **2002**.
- [18] a) P. Weyerstahl, H. Marschall, M. Schröder, H.-C. Wahlburg, V. K. Kaul, *Flavour Fragrance J.* **1997**, *12*, 315–325; b) C. Menut, G. Lamaty, P. Weyerstahl, H. Marschall, I. Seelmann, P. H. Amvam Zollo, *Flavour Fragrance J.* **1997**, *12*, 415–421.
- [19] E. Osawa, K. Aigami, N. Takaiishi, Y. Inamoto, Y. Fujikura, Z. Majerski, P. v. R. Schleyer, E. M. Engler, M. Farcasiu, *J. Am. Chem. Soc.* **1977**, *99*, 5361–5373.
- [20] a) A. Y. Hong, M. R. Krout, T. Jensen, N. B. Bennett, A. M. Harned, B. M. Stoltz, *Angew. Chem.* **2011**, *123*, 2808–2812; *Angew. Chem. Int. Ed.* **2011**, *50*, 2756–2760; b) A. Y. Hong, N. B. Bennett, M. R. Krout, T. Jensen, A. M. Harned, B. M. Stoltz, *Tetrahedron* **2011**, *67*, 10234–10248; c) N. B. Bennett, A. Y. Hong, B. M. Stoltz, *Org. Biomol. Chem.* **2012**, *10*, 56–59.
- [21] Carbamate **15** was prepared from commercial 2-methyl-2-vinyl-oxirane by modification of reported procedures. See: a) Y. Satake, T. Nishikawa, T. Hiramatsu, H. Araki, M. Isobe, *Synthesis* **2010**, 1992–1998; b) S. T. Heller, R. Sarpong, *Org. Lett.* **2010**, *12*, 4572–4575.
- [22] Substituted allylic carbamates have been used by Trost to achieve *O*-acylation of enolates under different reaction conditions. See: B. M. Trost, J. Xu, *J. Org. Chem.* **2007**, *72*, 9372–9375.
- [23] L. N. Mander, S. P. Sethi, *Tetrahedron Lett.* **1983**, *24*, 5425–5428.
- [24] a) D. C. Behenna, B. M. Stoltz, *J. Am. Chem. Soc.* **2004**, *126*, 15044–15045; b) J. T. Mohr, D. C. Behenna, A. M. Harned, B. M. Stoltz, *Angew. Chem.* **2005**, *117*, 7084–7087; *Angew. Chem. Int. Ed.* **2005**, *44*, 6924–6927; c) D. C. Behenna, J. T. Mohr, N. H. Sherden, S. C. Marinescu, A. M. Harned, K. Tani, M. Seto, S. Ma, Z. Novák, M. R. Krout, R. M. McFadden, J. L. Roizen, J. A. Enquist, Jr., D. E. White, S. R. Levine, K. V. Petrova, A. Iwashita, S. C. Virgil, B. M. Stoltz, *Chem. Eur. J.* **2011**, *17*, 14199–14223.
- [25] a) P. von Matt, A. Pfaltz, *Angew. Chem.* **1993**, *105*, 614–615; *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 566–568; b) J. Sprinz, G. Helmchen, *Tetrahedron Lett.* **1993**, *34*, 1769–1772; c) G. J. Dawson, C. G. Frost, J. M. J. Williams, S. J. Coote, *Tetrahedron Lett.* **1993**, *34*, 3149–3150.
- [26] a) K. Tani, D. C. Behenna, R. M. McFadden, B. M. Stoltz, *Org. Lett.* **2007**, *9*, 2529–2531; b) M. R. Krout, J. T. Mohr, B. M. Stoltz, *Org. Synth.* **2009**, *86*, 181–193.
- [27] For examples of the asymmetric alkylation of vinylogous esters or thioesters in the synthesis of natural products from our laboratory, see: a) D. E. White, I. C. Stewart, R. H. Grubbs, B. M. Stoltz, *J. Am. Chem. Soc.* **2008**, *130*, 810–811; b) S. R. Levine, M. R. Krout, B. M. Stoltz, *Org. Lett.* **2009**, *11*, 289–292; c) K. V. Petrova, J. T. Mohr, B. M. Stoltz, *Org. Lett.* **2009**, *11*, 293–295.
- [28] For a related example of the asymmetric alkylation of vinylogous thioesters, see: B. M. Trost, R. N. Bream, J. Xu, *Angew. Chem.* **2006**, *118*, 3181–3184; *Angew. Chem. Int. Ed.* **2006**, *45*, 3109–3112.
- [29] For selected IMDA reviews, see: a) B. R. Bear, S. M. Sparks, K. J. Shea, *Angew. Chem.* **2001**, *113*, 864–894; *Angew. Chem. Int. Ed.* **2001**, *40*, 820–849; b) K. C. Nicolaou, S. A. Snyder, T. Montagnon, G. Vassilikogiannakis, *Angew. Chem.* **2002**, *114*, 1742–1773; *Angew. Chem. Int. Ed.* **2002**, *41*, 1668–1698; c) K.-i. Takao, R. Munakata, K.-i. Tadano, *Chem. Rev.* **2005**, *105*, 4779–4807; d) M. Juhl, D. Tanner, *Chem. Soc. Rev.* **2009**, *38*, 2983–2992.
- [30] A related IMDA cycloaddition for the construction of an analogous tricyclic system with an activated alkyne has been reported. See: L. Evanno, A. Deville, B. Bodo, B. Nay, *Tetrahedron Lett.* **2007**, *48*, 4331–4333.
- [31] Relevant IMDA studies performed by Jung and Corey demonstrated that the diastereoselectivity of normal-demand Type I IMDA reactions can be reversed by substituting enone dienophiles with their corresponding sterically demanding cyclic ketal derivatives. See: a) M. E. Jung, K. M. Halweg, *Tetrahedron Lett.* **1981**, *22*, 3929–3932; b) E. J. Corey, P. A. Magriotis, *J. Am. Chem. Soc.* **1987**, *109*, 287–289.
- [32] E. H. Krenske, E. W. Perry, S. V. Jerome, T. J. Maimone, P. S. Baran, K. N. Houk, *Org. Lett.* **2012**, *14*, 3016–3019.
- [33] Our IMDA substrate does not fit neatly into the Type I or Type II IMDA classification, making prediction of the reaction outcome less straightforward.
- [34] a) M. Satoh, Y. Nomoto, N. Miyaura, A. Suzuki, *Tetrahedron Lett.* **1989**, *30*, 3789–3792; b) J. Y. Wu, B. Moreau, T. Ritter, *J. Am. Chem. Soc.* **2009**, *131*, 12915–12917; c) R. J. Ely, J. P. Morken, *J. Am. Chem. Soc.* **2010**, *132*, 2534–2535; d) R. J. Ely, J. P. Morken, *Org. Synth.* **2011**, *88*, 342–352.
- [35] See Supporting Information for additional details and experimental procedures.
- [36] C. A. Luchaco-Cullis, H. Mizutani, K. E. Murphy, A. H. Hov-eyda, *Angew. Chem.* **2001**, *113*, 1504–1508; *Angew. Chem. Int. Ed.* **2001**, *40*, 1456–1460.
- [37] R. W. Murray, M. Singh, *Org. Synth.* **1997**, *74*, 91–96.
- [38] CCDC 889569 (**1**) and 889570 (**21b**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [39] R. H. Crabtree, M. W. Davis, *J. Org. Chem.* **1986**, *51*, 2655–2661.
- [40] N. Kotoku, Y. Sumii, M. Kobayashi, *Org. Lett.* **2011**, *13*, 3514–3517.
- [41] Unfortunately, attempts to obtain samples or spectra of naturally isolated presilphiperfolan-1-ol (**1**) were unsuccessful.
- [42] An X-ray crystal structure for a synthetic compound with the same structure as our synthetic **1** (prepared from isocaryophyllene) was reported by Khomenko and co-workers (see Ref. [15b]). Our NMR data for synthetic **1** matches the spectral data for Khomenko’s synthetic **1** but not the spectral data for König’s reported natural **1**.