

Modular Synthesis of Phenanthridine Derivatives by Oxidative Cyclization of 2-Isocyanobiphenyls with Organoboron Reagents**


Mamoru Tobisu,* Keika Koh, Takayuki Furukawa, and Naoto Chatani*

Homolytic aromatic substitution (HAS) by an aryl radical and related processes^[1,2] have recently attracted considerable attention as alternatives to transition-metal-catalyzed C–H bond arylation^[3] for the construction of biaryl motifs. These HAS reactions have several advantages over the conventional methodology, including unique chemo- and regioselectivity as well as avoiding the requirement for precious-metal catalysts and ligands. Mechanistically, it has been proposed that the HAS reaction proceeds by the addition of an aryl radical to an arene, followed by the oxidative re-aromatization of the resulting cyclohexadienyl-type radical.^[1,2] Although there has been only limited application of the intermolecular HAS to the synthesis of biaryl systems because of issues associated with low levels of efficiency and selectivity, the application in intramolecular settings largely alleviates these issues and allows for straightforward access to biaryl moieties embedded in polycyclic system (Scheme 1a). It was envisioned that the


utility of this intramolecular HAS could be considerably expanded if a radical generated by the intermolecular addition of an external radical could undergo a HAS type cyclization (Scheme 1b). Herein, we report the development of a reaction guided by this hypothesis for the synthesis of phenanthridine framework.^[4]

To realize the intramolecular HAS reaction depicted in Scheme 1b, an isocyano group was selected as a radical acceptor (A in Scheme 1b) because intermolecular radical addition to isocyanide, forming an imido radical, occurs at a sufficiently fast rate to maintain the efficiency of the overall process.^[5] Furthermore, the potential for an imido radical intermediate to participate in a HAS reaction is partly supported by the report that a fragment radical derived from an initiator (such as, AIBN) can induce the radical cyclization of isocyanide by a HAS process.^[6]

After examining several radical precursors (see Supporting Information for details), the phenyl radical generated from phenylboronic acid and a manganese salt^[7] was identified as promoting the desired cyclization (Scheme 2). Thus, the reaction of isocyanide **1a** with phenylboronic acid (**2a**) in



Scheme 1. C–H bond functionalization by intramolecular homolytic aromatic substitution, see text for details.



Scheme 2. Manganese(III) mediated annulation of **1a** with **2a**.

[*] K. Koh, T. Furukawa, Prof. Dr. N. Chatani
Department of Applied Chemistry, Faculty of Engineering, Osaka University
Suita, Osaka 565-0871 (Japan)
E-mail: chatani@chem.eng.osaka-u.ac.jp

Dr. M. Tobisu
Center for Atomic and Molecular Technologies, Graduate School of Engineering
Osaka University, Suita, Osaka 565-0871 (Japan)
E-mail: tobisu@chem.eng.osaka-u.ac.jp

[**] This work was supported, in part, by a Grant-in-Aid for Scientific Research on Innovative Areas of “Molecular Activation Directed toward Straightforward Synthesis” from MEXT (Japan). We thank Prof. Kei Ohkubo and Prof. Shunichi Fukuzumi for obtaining EPR spectra. We also thank the Instrumental Analysis Center of the Faculty of Engineering at Osaka University for their assistance with HRMS.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201206115>.

the presence of $[\text{Mn}(\text{acac})_3]$ (3 equiv) gave phenanthridine **3aa** in 77% yield under relatively mild conditions (80 °C, 1 h).^[8] The only byproduct detected in the reaction mixture was phenanthridine **4a** (7%).^[9] The use of more than two equivalents of the manganese salt was required for the complete conversion of **1a**.^[10] This result is particularly relevant to the reaction mechanism.

As revealed in Table 1, a broad range of arylboronic acids were successfully coupled with isocyanide **1b** to give the corresponding phenanthridine bearing substituents adjacent to the ring nitrogen atom.^[11] Ethers, esters, fluorides, chlorides, and bromides are readily tolerated (entries 2–6).^[12] A sterically hindered 2-tolyl group was also incorporated without significant loss in the yield (entry 7). Fused arene and heteroaromatic systems, including an indole, a pyridine, and a thiophene, were also compatible with the reaction conditions (entries 8–11), which represented a significant outcome

bilities,^[1] it was envisaged that the re-aromatization proceeded via cationic intermediate **F** through single-electron oxidation by Mn^{III} followed by deprotonation, on the basis of the observation that more than two equivalents of Mn^{III} were required to drive the reaction to completion. Several experiments were performed to provide support for the proposed mechanism. One such experiment involved the addition of 2,2,6,6-tetramethyl-1-piperidinoxyl (TEMPO) to effectively quench the reaction, and the putative alkyl radical intermediate was intercepted by TEMPO (See Supporting Information). In a separate experiment, the use of 5-hexenylboronic acid (**2q**) exclusively afforded phenanthridine bearing a cyclopentylmethyl group, indicating that a rapid 5-exo process preceded the intermolecular addition to isocyanide (See Supporting Information).^[15]

To further demonstrate the value of the current two-component cyclization, we implemented this method in the synthesis of phenanthridine derivative **8**, which displays in vivo antitumor activity (Scheme 4).^[16] Our synthetic scheme features a late-stage introduction of a C6 substituent



Scheme 4. Application in the synthesis of **8**.

as well as amide at C4, allowing for rapid access to the product diversity required for structure–activity relationship studies.

In conclusion, we have developed a novel bimolecular coupling of 2-isocyanobiaryls with organoboronic acids by a formal HAS. The method enables the rapid divergent synthesis of phenanthridine and its π -extended analogues bearing aryl, heteroaryl and alkyl groups at the C6 position from readily accessible starting materials and a promoter. Further efforts are currently underway to develop synthetic methods capable of exploiting HAS and related processes in the transformation of C–H bonds.

Received: July 31, 2012

Revised: September 8, 2012

Published online: October 10, 2012

Keywords: boronic acid · homolytic aromatic substitution · isocyanide · manganese · phenanthridine

- [1] Selected Reviews: a) R. Bolton, G. H. Williams, *Chem. Soc. Rev.* **1986**, *15*, 261; b) A. Studer in *Radicals in Organic Synthesis*, Vol. 2, 1st ed. (Eds.: P. Renaud, M. P. Sibi), Wiley-VCH, Weinheim, **2001**, pp. 62–80; c) W. R. Bowman, J. M. D. Storey, *Chem. Soc. Rev.* **2007**, *36*, 1803; d) S. E. Vaillard, B. Schulte, A.

Studer in *Modern Arylation Methods* (Ed.: L. Ackermann), Wiley-VCH, Weinheim, **2009**, pp. 475–511; e) S. Yanagisawa, K. Itami, *ChemCatChem* **2011**, *3*, 827; f) A. Studer, D. P. Curran, *Angew. Chem.* **2011**, *123*, 5122; *Angew. Chem. Int. Ed.* **2011**, *50*, 5018.

- [2] Recent leading works on HAS and related processes for (hetero)biaryl synthesis: a) K.-i. Fujita, M. Nonogawa, R. Yamaguchi, *Chem. Commun.* **2004**, 1926; b) D. P. Curran, A. I. Keller, *J. Am. Chem. Soc.* **2006**, *128*, 13706; c) S. Yanagisawa, K. Ueda, T. Taniguchi, K. Itami, *Org. Lett.* **2008**, *10*, 4673; d) F. d. r. Vallée, J. J. Mousseau, A. B. Charette, *J. Am. Chem. Soc.* **2010**, *132*, 1514; e) W. Liu, H. Cao, A. Lei, *Angew. Chem.* **2010**, *122*, 2048; *Angew. Chem. Int. Ed.* **2010**, *49*, 2004; f) C.-L. Sun, H. Li, D.-G. Yu, M. Yu, X. Zhou, X.-Y. Lu, K. Huang, S.-F. Zheng, B.-J. Li, Z.-J. Shi, *Nat. Chem.* **2010**, *2*, 1044; g) I. B. Seiple, S. Su, R. A. Rodriguez, R. Gianatassio, Y. Fujiwara, A. L. Sobel, P. S. Baran, *J. Am. Chem. Soc.* **2010**, *132*, 13194; h) E. Shirakawa, K.-i. Itoh, T. Higashino, T. Hayashi, *J. Am. Chem. Soc.* **2010**, *132*, 15537; i) W. Liu, H. Cao, H. Zhang, H. Zhang, K. H. Chung, C. He, H. Wang, F. Y. Kwong, A. Lei, *J. Am. Chem. Soc.* **2010**, *132*, 16737; j) Y. Fujiwara, V. Domingo, I. B. Seiple, R. Gianatassio, M. Del Bel, P. S. Baran, *J. Am. Chem. Soc.* **2011**, *133*, 3292; k) W. Liu, H. Cao, J. Xin, L. Jin, A. Lei, *Chem. Eur. J.* **2011**, *17*, 3588; l) J. W. Lockner, D. D. Dixon, R. Risgaard, P. S. Baran, *Org. Lett.* **2011**, *13*, 5628; m) C.-L. Sun, Y.-F. Gu, W.-P. Huang, Z.-J. Shi, *Chem. Commun.* **2011**, 47, 9813; n) N. Uchiyama, E. Shirakawa, R. Nishikawa, T. Hayashi, *Chem. Commun.* **2011**, 47, 11671; o) G.-P. Yong, W.-L. She, Y.-M. Zhang, Y.-Z. Li, *Chem. Commun.* **2011**, 47, 11766; p) B. S. Bhakuni, A. Kumar, S. J. Balkrishna, J. A. Sheikh, S. Konar, S. Kumar, *Org. Lett.* **2012**, *14*, 2838; q) W.-C. Chen, Y.-C. Hsu, W.-C. Shih, C.-Y. Lee, W.-H. Chuang, Y.-F. Tsai, P. P.-Y. Chen, T.-G. Ong, *Chem. Commun.* **2012**, 48, 6702; r) L.-P. B. Beaulieu, R. S. Roman, F. Vallée, A. B. Charette, *Chem. Commun.* **2012**, 48, 8249.
- [3] Selected Reviews on catalytic C–H bond functionalization: a) F. Kakiuchi, T. Kochi, *Synthesis* **2008**, 3013; b) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem.* **2009**, *121*, 5196; *Angew. Chem. Int. Ed.* **2009**, *48*, 5094; c) O. Daugulis, H.-Q. Do, D. Shabashov, *Acc. Chem. Res.* **2009**, *42*, 1074; d) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147; e) D. A. Colby, R. G. Bergman, J. A. Ellman, *Chem. Rev.* **2010**, *110*, 624; f) K. Hirano, M. Miura, *Synlett* **2011**, 294; g) C.-L. Sun, B.-J. Li, Z.-J. Shi, *Chem. Rev.* **2011**, *111*, 1293; h) L. Ackermann, *Chem. Rev.* **2011**, *111*, 1315.
- [4] Synthesis of cyclic compounds other than phenanthridines by two- or multi-component radical cascades that involve intramolecular HAS: a) R. Leardini, D. Nanni, G. F. Pedulli, A. Tundo, G. Zanardi, *J. Chem. Soc. Perkin Trans. 1* **1986**, 1591; b) D. P. Curran, H. Liu, *J. Am. Chem. Soc.* **1991**, *113*, 2127; c) D. P. Curran, H. Liu, *J. Am. Chem. Soc.* **1992**, *114*, 5863; d) S. Araneo, F. Fontana, F. Minisci, F. Recupero, A. Serri, *Tetrahedron Lett.* **1995**, *36*, 4307; e) D. P. Curran, S.-B. Ko, H. Josien, *Angew. Chem.* **1995**, *107*, 2948; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 2683; f) R. Leardini, D. Nanni, P. Pareschi, A. Tundo, G. Zanardi, *J. Org. Chem.* **1997**, *62*, 8394; g) H. Josien, S.-B. Ko, D. Bom, D. P. Curran, *Chem. Eur. J.* **1998**, *4*, 67; h) C. M. Camaggi, R. Leadrini, D. Nanni, G. Zanardi, *Tetrahedron* **1998**, *54*, 5587; i) S. Bertrand, N. Hoffmann, J.-P. Pete, V. Bulach, *Chem. Commun.* **1999**, 2291; j) A. Dickschat, A. Studer, *Org. Lett.* **2010**, *12*, 3972.
- [5] a) D. Nanni in *Radicals in Organic Synthesis Vol. 2*, 1st ed. (Eds.: P. Renaud, M. P. Sibi), Wiley-VCH, Weinheim, **2001**, pp. 44–60; b) M. Minozzi, D. Nanni, P. Spagnolo, *Curr. Org. Chem.* **2007**, *11*, 1366.
- [6] a) D. Nanni, P. Pareschi, C. Rizzoli, P. Sgarabotto, A. Tundo, *Tetrahedron* **1995**, *51*, 9045; imidoyl radical generated from

- imine is also reported to participate in HAS; b) R. Leardini, A. Tundo, G. Zanardi, *Synthesis* **1985**, 107.
- [7] A. S. Demir, Ö. Reis, M. Emrullahoglu, *J. Org. Chem.* **2003**, 68, 578.
- [8] Reviews on synthesis of heterocycles from isocyanides: a) J. Campo, M. Garcia-Valverde, S. Marcaccini, M. J. Rojo, T. Torroba, *Org. Biomol. Chem.* **2006**, 4, 757; b) A. V. Lygin, A. de Meijere, *Angew. Chem.* **2010**, 122, 9280; *Angew. Chem. Int. Ed.* **2010**, 49, 9094; c) M. Tobisu, N. Chatani, *Chem. Lett.* **2011**, 40, 330.
- [9] **4a** was formed in 6% yield when the reaction was conducted in the absence of **2a**. The same cyclization was also promoted by AlCl₃, in place of [Mn(acac)₃], indicating that **4a** is formed by a Lewis acid mediated pathway. M. Tobisu, S. Yamaguchi, N. Chatani, *Org. Lett.* **2007**, 9, 3351.
- [10] [Mn(acac)₃] from TCI Co., Inc. costs approximately \$0.82 per gram.
- [11] Catalytic functionalization of C6 position of preformed phenanthridine: a) M. Tobisu, I. Hyodo, N. Chatani, *J. Am. Chem. Soc.* **2009**, 131, 12070; b) I. Hyodo, M. Tobisu, N. Chatani, *Chem. Asian J.* **2012**, 7, 1357.
- [12] Electron-rich arylboronic acids reacted faster than electron-deficient ones.
- [13] a) S. C. Dickerman, W. M. Feigenbaum, M. Fryd, N. Milstein, G. B. Vermont, I. Zimmerman, J. F. W. McOmie, *J. Am. Chem. Soc.* **1973**, 95, 4624; b) A. Bravo, H.-R. Bjørsvik, F. Fontana, L. Liguori, A. Mele, F. Minisci, *J. Org. Chem.* **1997**, 62, 7128.
- [14] L.-C. Campeau, M. Parisien, A. Jean, K. Fagnou, *J. Am. Chem. Soc.* **2006**, 128, 581.
- [15] Involvement of organomanganese species in all bond-forming steps cannot fully be excluded at this point. For our work on catalytic reactions that involve imido-yl-metal species, see: a) M. Tobisu, S. Imoto, S. Ito, N. Chatani, *J. Org. Chem.* **2010**, 75, 4835; b) M. Tobisu, H. Fujihara, K. Koh, N. Chatani, *J. Org. Chem.* **2010**, 75, 4841. EPR measurement of the frozen reaction mixture indicated the formation of an organic radical. See Supporting Information for details.
- [16] G. J. Atwell, B. C. Baguley, W. A. Denny, *J. Med. Chem.* **1988**, 31, 774.