


Synthetic Methods

Activation of a C(sp³)–H Bond by a Transient σ -Alkylpalladium(II) Complex: Synthesis of Spirooxindoles Through a Palladium-Catalyzed Domino Carbopalladation/C(sp³)–C(sp³) Bond-Forming Process**

Tiffany Piou, Luc Neuville,* and Jieping Zhu*

Catalytic C–H bond activation is an area of considerable current interest.^[1] Whereas a plethora of methods for the metal-catalyzed functionalization of arenes and heteroarenes have been developed to allow the formation of C(sp²)–C(sp²) bonds,^[2] methods dealing with C(sp²)–C(sp³) bond formation through activation of the less reactive C(sp³)–H bond are still an emerging research field.^[3–6] An even less exploited sub-field is the creation of a C(sp³)–C(sp³) bond through activation of a C(sp³)–H by a transient σ -alkylmetal complex.^[7] In connection with our interest in the development of domino processes involving C–H functionalization as a key step^[8] we became interested in the spirocyclization of **1** for the synthesis of spirooxindoles **2**.^[9] The underlying principle is shown in Scheme 1. Intramolecular carbopalladation of **1** would give intermediate **A**, which has a neopentyl-type σ -alkylpalladium(II) function. The latter, being sufficiently stable and ideally positioned, should be able to activate the nearby C(sp³)–H bond, thus leading to six-membered palladacycle **B**. Reductive elimination from **B** would then afford spirooxindole **2** and regenerate the Pd⁰ species.^[10,8] The realization of this unprecedented domino process is the subject of the present communication.

The easily accessible *N*-(2-bromophenyl)-2-(2,6-dimethylphenyl)-*N*-methylacrylamide (**1a**) was used as a substrate to test the feasibility of our projected domino process (Table 1).^[11] Initial efforts employed the catalytic system reported by Guyonnet and Baudoin.^[40] Under these conditions, the desired spirooxindole **2a** was indeed obtained,^[12] albeit accompanied by a significant amount of quinolinone **3a**, which resulted from a 6-*endo*-trig cyclization (**2a/3a** = 1:2,



Scheme 1. Domino carbopalladation/C(sp³)–C(sp³) bond-forming process. Ligands on Pd are omitted for clarity.

Table 1: Survey of reaction conditions for the cyclization of **1a**.^[a]

Entry	Ligand	Base	Additive	2a/3a ^[b]	Yield [%] ^[c]
1 ^[d]	PCy ₃ ·HBF ₄	Na ₂ CO ₃ /K ₂ CO ₃	PivOH	1:2	n.d.
2	PCy ₃ ·HBF ₄	Cs ₂ CO ₃	–	1:2	n.d.
3	PtBu ₃ ·HBF ₄	Cs ₂ CO ₃	–	2.2:1	n.d.
4	PPh ₃	Cs ₂ CO ₃	–	2.2:1	67
5	P(2-furyl) ₃	Cs ₂ CO ₃	–	2.5:1	71
6	XantPhos	Cs ₂ CO ₃	–	2.8:1	73
7	dppe	Cs ₂ CO ₃	–	3.6:1	n.d.
8 ^[e]	dppp	Cs ₂ CO ₃	–	12:1	85
9 ^[f]	dppp	Cs ₂ CO ₃	–	3.3:1	n.d.
10 ^[g]	dppp	Na ₂ CO ₃	–	n.d.	n.d.
11	dppp	K ₂ CO ₃	–	32.7:1	94
12	dppp	RbCO ₃	–	17.1:1	85

[a] **1a** (1.0 equiv), Pd(OAc)₂ (0.1 equiv), ligand (0.2 equiv), and base (1.3 equiv) in mesitylene (0.15 M) at 140°C, under argon for 1–2 h. [b] Determined by ¹H NMR analysis of the crude material. [c] Yield of isolated **2a**. [d] Reaction was performed using PivOH (0.3 equiv), Na₂CO₃ (0.3 equiv), and K₂CO₃ (1.3 equiv). [e] *n*Bu₄NCl (1.0 equiv) was used. [f] Reaction was performed at 120°C. [g] Incomplete conversion after 24 h. Cy = cyclohexyl, dppe = ethylenebis(diphenylphosphine), dppp = 1,3-bis(diphenylphosphino)propane, n.d. = not determined, OAc = acetate, Xantphos = 4,5-bis(diphenylphosphino)-9,9-dimethyl-xanthene.

entry 1, Table 1). The predominant formation of **3a** was surprising, as a 5-*exo*-trig cyclization is generally overwhelmingly favored over a 6-*endo*-trig cyclization in the Heck reaction.^[13] Systematic studies of the reaction conditions

[*] T. Piou, Prof. Dr. J. Zhu
Ecole Polytechnique Fédérale de Lausanne, EPFL-SB-ISIC-LSPN
BCH 5304, 1015 Lausanne (Switzerland)
E-mail: jieping.zhu@epfl.ch
Homepage: http://lspn.epfl.ch
Dr. L. Neuville
Centre de Recherche de Gif, Laboratoire International Associé,
Institut de Chimie des Substances Naturelles, CNRS
91198 Gif-sur-Yvette Cedex (France)
E-mail: luc.neuville@icsn.cnrs-gif.fr

[**] Financial support from EPFL (Switzerland), Swiss National Science Foundation (SNF), Swiss National Centres of Competence in Research (NCCR) and ICSN, CNRS (France) are gratefully acknowledged. T.P. thanks ICSN and EPFL for a doctoral fellowship. We thank Dr. Pascal Mieville for helpful discussion on NMR experiments.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201206267>.

revealed that the ligand, the base, and the temperature exerted a dramatic influence on the regioselectivity of the reaction. Some representative results are summarized in Table 1.^[14] Overall, the optimized conditions consisted of Pd(OAc)₂ (OAc = acetate; 0.1 equiv), 1,3-bis(diphenylphosphino)propane (dppp; 0.2 equiv), K₂CO₃ (1.3 equiv) in mesitylene (0.15 M) at 140 °C. Under these conditions, the domino 5-*exo* carbopalladation/C(sp³)–C(sp³) bond-forming process occurred smoothly to provide spirooxindole **2a** in 94 % yield (entry 11). Attempts to render this reaction enantioselective through the use of chiral bidentate ligands such as (*R,R*)-BINAP or (*S,S*)-DIOP failed to produce any significant chiral induction (see the Supporting Information).

With the optimized conditions in hand, the scope of the domino process was next evaluated. The influence of the haloaniline subunit on the reaction outcome was examined first (Table 2). Whereas the iodoanilide (**1b**, X = I) afforded the spirooxindole **2a** in 92 % yield, the chloroanilide (**1c**, X = Cl) failed to undergo the spirocyclization (Table 2, entry 1 vs. 2). The use of a tertiary amide was mandatory to ensure the occurrence of the domino reaction, as treatment of secondary amide **1d** failed to produce the spirocycle (entry 3). Cycliza-

tions of *N*-benzyl, *N*-*para*-methoxybenzyl and *N*-phenyl anilide took place smoothly to provide the spirooxindoles (**2e–g**) in excellent yields (entries 4–6). In these cases, the direct aromatic C(sp²)–H activation was not observed.^[15] The *N*-2-(trimethylsilyl)ethoxy)methyl group was compatible as shown in the synthesis of **2h** (entry 7). Electron-donating or electron-withdrawing groups were well tolerated (entries 8–12), and substrates bearing substituents *meta*- and *para*- to the halide furnished spirooxindoles in excellent yields (entries 13–14). In contrast, the presence of a substituent *ortho* to the halide preferentially underwent a 6-*endo*-Heck cyclization to afford the quinolinone **3p** in 60 % yield, together with a 37 % yield of the desired spirooxindole **2p** (entry 15). Finally, heterocycles such as benzopyrazine underwent cyclization without difficulty to afford **2q** in excellent yield (entry 16).

Variation of the α -aryl substituent on the acrylate was next evaluated (Table 3). Substituents such as *tert*-butyl, methyl, and methoxy groups were tolerated, providing the corresponding spirooxindoles (**2r–2t**) in excellent yields (Table 3, entries 1–3). On the other hand, 2-(*o*-tolyl)acrylamide (**1u**) failed to produce the desired oxindole, **2u** (entry 4). In the case of unsymmetrical 2,5-disubstituted derivative **1v**, the cyclization proceeded chemoselectively on the methyl group to produce compound **2v** in 97 % yield (entry 5). However, when anilide **1w**, which bears a 2,5-diethylbenzene moiety, was submitted to the reaction conditions, cyclization by activation of the methylene C(sp³)–H group took place to provide spirooxindole **2w** as a mixture of two diastereomers (68 %, d.r. = 2:1, entry 6). This result is

Table 2: Reaction scope: varying the aniline.^[a]

Entry	Substrate (1)	Product (2 or 3)	Yield [%] ^[b]
1 ^[c]			92
2 ^[d,e]		2a	trace
3		2d , R = H	0
4		2e , R = Ph	99
5		2f , R = Bn	96
6		2g , R = PMB	89
7		2h , R = SEM	85
8		2i , R = CO ₂ Me	85
9		2j , R = F	78
10		2k , R = CN	79
11		2l , R = OMe	96
12		2m , R = Me	98
13		2n , R ² = H	97
14		2o , R ² = Me	97
15 ^[f]		3p	60
16		2q	97

[a] **1** (1.0 equiv), Pd(OAc)₂ (0.1 equiv), dppp (0.2 equiv), K₂CO₃ (1.3 equiv) in mesitylene ($c = 0.15$ M) at 140 °C under argon for 1–5 h. [b] Yield of isolated product. [c] X = I. [d] 24 h. [e] X = Cl. [f] Oxindole **2p** (structure not shown) was isolated in 37 % yield. Bn = benzyl, PMB = *p*-methoxybenzyl ether, SEM = 2-(trimethylsilyl)ethoxymethyl.


Table 3: Reaction scope: varying the α -aryl substituent.^[a]

Entry	Substrate (1)	Product (2)	Yield [%] ^[b]
1		2r , R = OMe	92
2		2s , R = <i>t</i> Bu	91
3		2t	97
4		2u	0
5		2v	97
6		2w ^[c]	68
7		2x + 2x' ^[d]	89

[a] **1** (1.0 equiv), Pd(OAc)₂ (0.1 equiv), dppp (0.2 equiv), K₂CO₃ (1.3 equiv) in mesitylene (0.15 M) at 140 °C under argon for 1–5 h. [b] Yield of isolated product. [c] d.r. = 2:1. [d] **2x**/**2x'** = 1:3.

a rare example of $C(sp^3)-H$ bond activation/ $C(sp^3)-C(sp^3)$ bond formation of a methylene group. When the phenyl group was replaced by a naphthalene derivative (**1x**), activation of the naphthyl $C(sp^2)-H$ bond dominated over the methyl $C(sp^3)-H$ bond, which is consistent with the notion that activation of $C(sp^2)-H$ bonds is generally favored over $C(sp^3)-H$ bonds.


Although bidentate ligands have been used in Pd-catalyzed $C(sp^2)-H$ bond activation for the formation of aryl-aryl bonds,^[16] bulky and electron-rich monodentate ligands are generally the ligand of choice for $C(sp^3)-H$ activation processes. Considering the reversibility of the carbopalladation step, an appealing rationale for the beneficial effect of dppp in the present domino transformation could be that it chelates strongly to Pd and therefore: 1) favors the formation of the less sterically congested neopentyl-type σ -alkyl Pd^{II} complex (5-*exo* cyclization) over the tertiary σ -alkyl Pd^{II} complex **C** (6-*endo* cyclization; Scheme 2) owing



Scheme 2. Regioselectivity of the domino process.

to the increased effective size of metal coordination sphere;^[17] 2) disfavors the β -hydride elimination as the 14-electron Pd^{II} complex **C** has no vacant coordination site available to accommodate the β -hydride elimination (Scheme 2, step 5);^[18] 3) enforce the *cis* geometry of the two alkyl groups in intermediate **B**, facilitating therefore the reductive elimination process.^[19] The low selectivity displayed by XantPhos, might indicate that dissociation was occurring with this ligand.^[20] Indeed, the similar selectivity observed with XantPhos and with a monophosphine ligand was in agreement with this assumption (entries 3–6, Table 1).

A kinetic study determined that the reaction is zero order in **1a** (see the Supporting Information), indicating a fast oxidative addition in which the Pd catalyst is saturated as a Pd^{II}-aryl species, and ruled out the possibility of oxidative addition being the rate-determining step. To gain further insight into the reaction mechanism, the kinetic isotope effect was examined next (Scheme 3).^[21] Submitting [**D**₃]-**1a** to our standard spirocyclization conditions afforded [**D**₃]-**2a**/[**D**₂]-**2a** in a ratio of 3.4:1, hence with an intramolecular KIE value of 3.4. This result indicated that the C–H cleavage is directly



Scheme 3. Kinetic isotope effect experiments.

involved in the $C(sp^3)-H$ activation step leading to palladacycle **B**. An intermolecular competition experiment using an equimolar amount of **1a** and its hexadeuterated derivative [**D**₆]-**1a** provided an intermolecular KIE value of 1.2. More significantly, side-by-side kinetic experiments using **1a** and [**D**₆]-**1a** provided an intermolecular KIE (k_H/k_D , initial rate constant) value of 0.90. The inverse KIE observed in this last experiment might indicate that reductive elimination, rather than C–H activation, was the turnover-limiting step of the domino process, and that the observed inverse intermolecular KIE is in fact a secondary kinetic isotope effect. To the best of our knowledge, this is the first time an inverse KIE was observed for a palladium-catalyzed $C(sp^3)-H$ functionalization reaction.^[22] By invoking the Curtin–Hammett principle,^[23] these mechanistic considerations also explain the failure encountered in developing an enantioselective variant of this domino process.

As the biphosphine-chelated Pd intermediate is essential for the efficacy of the present transformation, we speculated that the intramolecular base-assisted concerted metalation-deprotonation (CMD) mechanism may not be operating in our case, as this would implicate a pentacoordinated Pd^{II} intermediate. Alternatively, an intermolecular base-assisted CMD mechanism could be advanced to account for the reaction outcome. The fact that compound **4** (which would result from deuterium scrambling) was not observed when [**D**₆]-**1a** was subjected to the spirocyclization conditions also allowed us to speculate that the C–H activation was irreversible (Scheme 3).

In summary, we have reported an efficient one-pot synthesis of spirooxindoles using a novel palladium(0)-catalyzed domino carbopalladation/ $C(sp^3)-C(sp^3)$ bond-forming process. The strategically positioned double bond allows the remote $C(sp^3)-H$ bond to be activated by a transient σ -alkylmetal complex. We believe that this reaction provides a new means for the elaboration of domino reactions by including a $C(sp^3)-C(sp^3)$ bond-forming

reaction as an elementary step, and offers an unusual bond disconnection logic.

Experimental Section

Anilide **1** (0.10 mmol), Pd(OAc)₂ (0.01 mmol, 0.1 equiv), dppp (0.02 mmol, 0.2 equiv), and K₂CO₃ (0.13 mmol, 1.3 equiv) were added to a screw-cap vial equipped with a magnetic stir bar. The vial was purged with argon for 10 min. Degassed mesitylene (0.50 mL) was added and the mixture was stirred at 140 °C until TLC monitoring indicated the complete disappearance of the starting material. After being cooled to room temperature, ethyl acetate was added and the reaction mixture was filtered through a short pad of Celite. The filtrate was concentrated under reduced pressure and the resulting crude product was purified by flash column chromatography (SiO₂, petroleum ether/ethyl acetate; see the Supporting Information for details) to provide the desired spirooxindole **2**.

Received: August 5, 2012

Published online: October 10, 2012

Keywords: C–H activation · domino reaction · heterocycles · palladium · spirooxindoles

- [1] C–H Activation in *Top. Curr. Chem.* (Eds.: J.-Q. Yu, Z. Shi), Vol. 292, Springer, Berlin, **2010**.
- [2] For recent reviews on C–H activation, see: a) A. R. Dick, M. S. Sanford, *Tetrahedron* **2006**, 62, 2439–2463; b) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* **2007**, 107, 174–238; c) B.-J. Li, S.-D. Yang, Z.-J. Shi, *Synlett* **2008**, 949–957; d) R. Giri, B.-F. Shi, K. M. Engle, N. Maugel, J.-Q. Yu, *Chem. Soc. Rev.* **2009**, 38, 3242–3272; e) T.-S. Mei, L. Kou, S. Ma, K. M. Engle, J.-Q. Yu, *Synthesis* **2012**, 1778–1791; f) J. Wencel-Delord, T. Dröge, F. Liu, F. Glorius, *Chem. Soc. Rev.* **2011**, 40, 4740–4761; for C–H functionalization in natural product synthesis, see: g) L. McMur-ray, F. O'Hara, M. J. Gaunt, *Chem. Soc. Rev.* **2011**, 40, 1885–1898; h) W. R. Gutekunst, P. S. Baran, *Chem. Soc. Rev.* **2011**, 40, 1976–1991.
- [3] a) O. Baudoin, *Chem. Soc. Rev.* **2011**, 40, 4902–4911; b) H. Li, B.-J. Li, Z.-J. Shi, *Catal. Sci. Technol.* **2011**, 1, 191–206; for a review on the cross-coupling of C(sp³)–H with organometallic reagents, see: c) M. Wasa, K. M. Engle, J.-Q. Yu, *Isr. J. Chem.* **2010**, 50, 605–616.
- [4] For selected recent examples of palladium(0)-catalyzed C(sp³)–H activation, see: a) G. Dyker, *Angew. Chem.* **1992**, 104, 1079–1081; *Angew. Chem. Int. Ed. Engl.* **1992**, 31, 1023–1025; b) O. Baudoin, A. Herrbach, F. Gueritte, *Angew. Chem.* **2003**, 115, 5914–5918; *Angew. Chem. Int. Ed.* **2003**, 42, 5736–5740; c) H. Ren, P. Knochel, *Angew. Chem.* **2006**, 118, 3541–3544; *Angew. Chem. Int. Ed.* **2006**, 45, 3462–3465; d) C.-G. Dong, Q.-S. Hu, *Angew. Chem.* **2006**, 118, 2347–2350; *Angew. Chem. Int. Ed.* **2006**, 45, 2289–2292; e) M. Lafrance, S. I. Gorelsky, K. Fagnou, *J. Am. Chem. Soc.* **2007**, 129, 14570–14571; f) T. Watanabe, S. Oishi, N. Fujii, H. Ohno, *Org. Lett.* **2008**, 10, 1759–1762; g) M. Chaumontet, R. Piccardi, N. Audic, J. Hitce, J.-L. Peglion, E. Clot, O. Baudoin, *J. Am. Chem. Soc.* **2008**, 130, 15157–15166; h) A. Salcedo, L. Neuville, J. Zhu, *J. Org. Chem.* **2008**, 73, 3600–3603; i) M. Wasa, K. M. Engle, J. Q. Yu, *J. Am. Chem. Soc.* **2009**, 131, 9886–9887; j) S. Rousseaux, M. Davi, J. Sofack-Kreutzer, C. Pierre, C. E. Kefalidis, E. Clot, K. Fagnou, O. Baudoin, *J. Am. Chem. Soc.* **2010**, 132, 10706–10716; k) C. Pierre, O. Baudoin, *Org. Lett.* **2011**, 13, 1816–1819; l) J. Pan, M. Su, S. L. Buchwald, *Angew. Chem.* **2011**, 123, 8806–8810; *Angew. Chem. Int. Ed.* **2011**, 50, 8647–8651; m) C. Tsukano, M. Okuno, Y. Takemoto, *Angew. Chem.* **2012**, 124, 2817–2820; *Angew. Chem. Int. Ed.* **2012**, 51, 2763–2766; n) S. Rousseaux, B. Liégault, K. Fagnou, *Chem. Sci.* **2012**, 3, 244–248; o) M. Guyonnet, O. Baudoin, *Org. Lett.* **2012**, 14, 398–401.
- [5] For palladium(II)-catalyzed C(sp²)–H activation, see: a) B. Liégault, K. Fagnou, *Organometallics* **2008**, 27, 4841–4843; b) J. J. Neumann, S. Rakshit, T. Dröge, F. Glorius, *Angew. Chem.* **2009**, 121, 7024–7027; *Angew. Chem. Int. Ed.* **2009**, 48, 6892–6895; c) M. Wasa, K. M. Engle, J.-Q. Yu, *J. Am. Chem. Soc.* **2010**, 132, 3680–3681; d) E. J. Yoo, M. Wasa, J.-Q. Yu, *J. Am. Chem. Soc.* **2010**, 132, 17378–17380; e) K. J. Stowers, K. C. Fortner, M. S. Sanford, *J. Am. Chem. Soc.* **2011**, 133, 6541–6544; f) P. Novák, A. Correa, J. Gallardo-Donaire, R. Martin, *Angew. Chem.* **2011**, 123, 12444–12447; *Angew. Chem. Int. Ed.* **2011**, 50, 12236–12239; g) Y. Ano, M. Tobisu, N. Chatani, *J. Am. Chem. Soc.* **2011**, 133, 12984–12986; h) G. He, G. Chen, *Angew. Chem.* **2011**, 123, 5298–5302; *Angew. Chem. Int. Ed.* **2011**, 50, 5192–5196; i) E. T. Nadres, O. Daugulis, *J. Am. Chem. Soc.* **2012**, 134, 7–10; j) A. Iglesias, R. Alvarez, A. R. de Lera, K. Muñiz, *Angew. Chem.* **2012**, 124, 2268–2271; *Angew. Chem. Int. Ed.* **2012**, 51, 2225–2228.
- [6] For enantioselective C(sp³)–H functionalization, see: a) M. Nakanishi, D. Katayev, C. Besnard, E. P. Kündig, *Angew. Chem.* **2011**, 123, 7576–7579; *Angew. Chem. Int. Ed.* **2011**, 50, 7438–7441; b) S. Anas, A. Cordi, H. B. Kagan, *Chem. Commun.* **2011**, 47, 11483–11485; c) T. Saget, S. Lemouzy, N. Cramer, *Angew. Chem.* **2012**, 124, 2281–2285; *Angew. Chem. Int. Ed.* **2012**, 51, 2238–2242; d) D. Katayev, M. Nakanishi, T. Bürgi, E. P. Kündig, *Chem. Sci.* **2012**, 3, 1422–1425; e) N. Martin, C. Pierre, M. Davi, R. Jazzar, O. Baudoin, *Chem. Eur. J.* **2012**, 18, 4480–4484; for asymmetric Pd^{II}-catalyzed enantioselective C–(sp³)–H functionalization, see: f) B.-F. Shi, N. Maugel, Y.-H. Zhang, J.-Q. Yu, *Angew. Chem.* **2008**, 120, 4960–4964; *Angew. Chem. Int. Ed.* **2008**, 47, 4882–4886; g) M. Wasa, K. M. Engle, D. W. Lin, E. J. Yoo, J.-Q. Yu, *J. Am. Chem. Soc.* **2011**, 133, 19598–19601; for a review, see: Ref. [2d].
- [7] a) M. Catellani, E. Motti, S. Ghelli, *Chem. Commun.* **2000**, 2003–2004; for cyclization leading to cyclopropanes, see: b) F. Liron, P. Knochel, *Tetrahedron Lett.* **2007**, 48, 4943–4946; c) H.-S. Kim, S. Gowrisankar, S. H. Kim, J. N. Kim, *Tetrahedron Lett.* **2008**, 49, 3858–3861; d) Q. Huang, R. C. Larock, *Tetrahedron Lett.* **2009**, 50, 7235–7238; for reactions under oxidative conditions (Pd^{II}/Pd^{IV}), see: e) T. Wu, X. Mu, G. Liu, *Angew. Chem.* **2011**, 123, 12786–12789; *Angew. Chem. Int. Ed.* **2011**, 50, 12578–12581; f) X. Mu, T. Wu, H.-Y. Wang, Y.-L. Guo, G. Liu, *J. Am. Chem. Soc.* **2012**, 134, 878–881; for Ir-catalyzed C(sp³)–H silylation, see: g) E. M. Simmons, J. F. Hartwig, *Nature* **2012**, 483, 70–73; for Rh-catalyzed C(sp³)–C(sp³) bond formation triggered by a metallacycle intermediate, see: h) Y. Oonishi, Y. Kitano, Y. Sato, *Angew. Chem.* **2012**, 124, 7417–7420; *Angew. Chem. Int. Ed.* **2012**, 51, 7305–7308; for the cross-coupling of C(sp³)–H with organometallic reagents, see: i) D.-H. Wang, M. Wasa, R. Giri, J.-Q. Yu, *J. Am. Chem. Soc.* **2008**, 130, 7190–7191; j) D. Shabashov, O. Daugulis, *J. Am. Chem. Soc.* **2010**, 132, 3965–3972.
- [8] a) G. Cuny, M. Bois-Choussy, J. Zhu, *Angew. Chem.* **2003**, 115, 4922–4925; *Angew. Chem. Int. Ed.* **2003**, 42, 4774–4777; b) A. Pinto, L. Neuville, J. Zhu, *Angew. Chem.* **2007**, 119, 3355–3359; *Angew. Chem. Int. Ed.* **2007**, 46, 3291–3295; c) A. Salcedo, L. Neuville, C. Rondot, P. Retailleau, J. Zhu, *Org. Lett.* **2008**, 10, 857–860; d) T. Gerfaud, L. Neuville, J. Zhu, *Angew. Chem.* **2009**, 121, 580–585; *Angew. Chem. Int. Ed.* **2009**, 48, 572–577; e) A. Pinto, L. Neuville, J. Zhu, *Tetrahedron Lett.* **2009**, 50, 3602–3605; f) S. Jaegli, W. Erb, P. Retailleau, J.-P. Vors, L. Neuville, J. Zhu, *Chem. Eur. J.* **2010**, 16, 5863–5867; g) S. Jaegli, J. Dufour, H.-L. Wei, T. Piou, X.-H. Duan, J.-P. Vors, L. Neuville, J. Zhu, *Org. Lett.* **2010**, 12, 4498–4501; h) T. Piou, L. Neuville, J. Zhu, *Org. Lett.* **2012**, 14, 3760–3763.

- [9] a) C. Marti, E. M. Carreira, *Eur. J. Org. Chem.* **2003**, 2209–2219; b) C. V. Galliford, K. A. Scheidt, *Angew. Chem.* **2007**, *119*, 8902–8912; *Angew. Chem. Int. Ed.* **2007**, *46*, 8748–8758; c) B. M. Trost, M. K. Brennan, *Synthesis* **2009**, 3003–3025; d) A. Millemaggi, R. J. K. Taylor, *Eur. J. Org. Chem.* **2010**, 4527–4547; e) F. Zhou, Y.-L. Liu, J. Zhou, *Adv. Synth. Catal.* **2010**, *352*, 1381–1407; f) R. Rios, *Chem. Soc. Rev.* **2012**, *41*, 1060–1074.
- [10] For Pd-catalyzed domino Heck/C(sp²)-H functionalization, see: a) D. Brown, R. Grigg, V. Sridharan, V. Tambyrajah, *Tetrahedron Lett.* **1995**, *36*, 8137–8140; b) R. T. Ruck, M. A. Huffman, M. M. Kim, M. Shevlin, W. V. Kandur, I. W. Davies, *Angew. Chem.* **2008**, *120*, 4789–4792; *Angew. Chem. Int. Ed.* **2008**, *47*, 4711–4714; c) Y. Hu, F. Song, F. Wu, Cheng, S. Wang, *Chem. Eur. J.* **2008**, *14*, 3110–3117; d) Y. Hu, C. Yu, D. Ren, Q. Hu, L. Zhang, D. Cheng, *Angew. Chem.* **2009**, *121*, 5556–5559; *Angew. Chem. Int. Ed.* **2009**, *48*, 5448–5451; e) Y. Hu, Y. Ouyang, Y. Qu, Q. Hu, H. Yao, *Chem. Commun.* **2009**, 4575–4577; f) O. René, D. Lapointe, K. Fagnou, *Org. Lett.* **2009**, *11*, 4560–4563; g) G. Satyanarayana, C. Maichle-Mössmer, M. E. Maier, *Chem. Commun.* **2009**, 1571–1573; h) T. T. Jayanth, L. Zhang, T. S. Johnson, H. C. Malinakova, *Org. Lett.* **2009**, *11*, 815–818; i) K. H. Kim, H. S. Lee, S. H. Kim, S. H. Kim, J. N. Kim, *Chem. Eur. J.* **2010**, *16*, 2375–2380.
- [11] For details, see the Supporting Information. The 2-(2,6-dimethylphenyl)acrylate and other 2-arylacrylates used in this work were prepared from pyruvate and aryl halides; see: J. Barluenga, M. Tomás-Gamasa, F. Aznar, C. Valdés, *Chem. Eur. J.* **2010**, *16*, 12801–12803.
- [12] I. Gorokhovik, L. Neuville, J. Zhu, *Org. Lett.* **2011**, *13*, 5536–5539.
- [13] *The Mizoroki-Heck Reaction* (Ed.: M. Oestreich), Wiley, Chichester, **2009**.
- [14] For a detailed survey of reaction conditions, see the Supporting Information.
- [15] L.-C. Campeau, M. Parisien, A. Jean, K. Fagnou, *J. Am. Chem. Soc.* **2006**, *128*, 581–590.
- [16] S. Pascual, P. de Mendoza, A. A. C. Braga, F. Maseras, A. M. Echavarren, *Tetrahedron* **2008**, *64*, 6021–6029.
- [17] J. Mo, J. Ruan, L. Xu, Z. Hyder, O. Saidi, S. Liu, W. Pei, J. Xiao, *J. Mol. Catal. A* **2007**, *261*, 267–275.
- [18] Inversely, “ligandless” conditions are indeed known to favor the 6-*endo* mode of cyclization, presumably by accelerating the β -hydride elimination process; see: a) J. H. Rigby, R. C. Hughes, M. J. Heeg, *J. Am. Chem. Soc.* **1995**, *117*, 7834–7835; b) J. W. Dankwardt, L. A. Flippin, *J. Org. Chem.* **1995**, *60*, 2312–2313; c) A. Bombrun, O. Sageot, *Tetrahedron Lett.* **1997**, *38*, 1057–1060; for an earlier example of formal 6-*endo* cyclization by rearrangement of a 5-*exo* carbopalladation product, see: d) M. O. Terpkko, R. F. Heck, *J. Am. Chem. Soc.* **1979**, *101*, 5281–5286.
- [19] J. E. Marcone, K. G. Moloy, *J. Am. Chem. Soc.* **1998**, *120*, 8527–8528.
- [20] For a recent review on the bite angle effects of diphosphines in a metal-catalyzed process, see: M.-N. Birkholz née Gensow, Z. Freixa, P. W. N. M. van Leeuwen, *Chem. Soc. Rev.* **2009**, *38*, 1099–1118.
- [21] a) M. Gómez-Gallego, M. A. Sierra, *Chem. Rev.* **2011**, *111*, 4857–4963; b) E. M. Simmons, J. F. Hartwig, *Angew. Chem.* **2012**, *124*, 3120–3126; *Angew. Chem. Int. Ed.* **2012**, *51*, 3066–3072.
- [22] For an inverse isotope effect on the reductive elimination of an alkane from a metal hydride complex, see: W. D. Jones, *Acc. Chem. Res.* **2003**, *36*, 140–146.
- [23] J. I. Seeman, *Chem. Rev.* **1983**, *83*, 83–134.