

Enantioselective Construction of α -Quaternary Cyclobutanones by Catalytic Asymmetric Allylic Alkylation**

Corey M. Reeves, Christian Eidamshaus, Jimin Kim, and Brian M. Stoltz*

Dedicated to Dr. Michael Mara

The asymmetric alkylation of enolates to generate α -quaternary carbonyl compounds has become a mainstay transformation in organic synthesis.^[1] In this domain, cyclobutanones have received far less attention relative to their five-, six- and seven-membered congeners, despite the fact that these compounds and their derivatives are prevalent in important biologically active natural products (Figure 1 A).^[2] Additionally, cyclobutanes have been shown to serve as highly valuable synthetic intermediates for a variety of transformations.^[3] The dearth of reports describing the asymmetric alkylation of cyclobutanones may be attributed to the fact that these compounds possess an estimated 26–28.6 kcal mol⁻¹ of ringstrain^[4] and, in turn, exhibit enhanced carbonyl electrophilicity.^[5] The propensity of cyclobutanones to alleviate this strain through electrophilic ring opening is often a limiting challenge during their manipulation. Moreover, the energetic requirements for enolization of cyclobutanones are compounded by a concomitant increase in ringstrain to 31–34 kcal mol⁻¹ (calculated for cyclobutene, Figure 1 B)^[3] as well as enforced deviation from the more favorable puckered conformation (Figure 1 C).^[6] In the case of α -substituted cyclobutanones, enolization is further impeded by the development of torsional strain between the putative enolate substituents (Figure 1 C).^[7]

Given these data, it is not surprising that previous methods for the preparation of enantioenriched cyclobutanes have relied primarily on either [2+2] cycloaddition reactions^[8] or ring expansion from various cyclopropane derivatives.^[9] Recent reports from Baudoin on annulating C–H activation^[10] as well as disclosures from Toste^[11] and Echavarren^[12] that employ gold(I) catalysis to affect cyclopropa-




Figure 1. A) Representative bioactive natural products possessing cyclobutanoid core structures. B) Increased ring strain generated upon enolization. C) Increased conformational and torsional strain upon unsaturation of four-membered rings.

noid rearrangements have emerged as significant new methods for the construction of cyclobutanes. Despite these advances, transformations that produce chiral cyclobutanones remain limited in scope and very few methods exist for the catalytic construction of chiral cyclobutanones from achiral starting materials.^[9b,13] In order to address these limitations and to further develop the nucleophilic chemistry of these unusually reactive compounds we report herein the first direct transition metal-catalyzed asymmetric α -alkylation of cyclobutanones to form all-carbon quaternary centers.


A longstanding interest of our research group is the asymmetric transition metal-catalyzed α -functionalization of carbonyl compounds to form all-carbon quaternary centers.^[14] In the course of our studies, we have developed a series of phosphinoxazoline (PHOX) ligands with varied steric and electronic properties that exhibit a range of reactivities and selectivities. We have found that the use of electron-deficient ligands (e.g. **L2** and **L3**, Table 1) often results in superior asymmetric induction in instances where electron-rich or electron-neutral ligands perform poorly.^[15] Examination of ligand electronic effects would, therefore, help to inform our development of a method for the asymmetric allylic alkylation of cyclobutanones.

[*] C. M. Reeves, C. Eidamshaus, J. Kim, Prof. B. M. Stoltz
The Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology
1200 E. California Blvd, MC 101-20, Pasadena, CA 91125 (USA)
E-mail: stoltz@caltech.edu

[**] The authors wish to thank NIH-NIGMS (R01GM080269-01), Deutsche Forschungsgemeinschaft (DFG postdoctoral fellowship to C.E.), Amgen, Abbott, Boehringer Ingelheim, and Caltech for financial support. Materia, Inc. is gratefully acknowledged for the donation of catalysts. Dr. Mona Shahgholi and Naseem Torian are acknowledged for high-resolution mass spectrometry assistance. Jonny Gordon is acknowledged for insightful discussions. Robert Craig is gratefully acknowledged for providing compound **L3** and for insightful discussion.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201301815>.

We first established a simple and efficient reaction sequence to access allyl 1-alkyl-2-oxocyclobutanecarboxylate substrates (**4**) (Scheme 1). Diazotization of commercially available 1,3-cyclopentane dione (**1**) with *para*-acetamidobenzenesulfonyl azide (*p*-ABSA)^[16] delivered the corresponding diazodiketone (**2**) in consistently good yields. Microwave-promoted Wolff rearrangement of diketone **2** in



Scheme 1. Construction of allyl 1-benzyl-2-oxocyclobutane-carboxylate (**4l**).


the presence of an allylic alcohol^[17] (e.g., allyl alcohol (**3**)), followed by alkylation with an alkyl halide (e.g., benzyl bromide) furnished the allyl 1-alkyl-2-oxocyclobutanecarboxylates in good yields over two steps. With a quick and efficient method to access the desired substrates in hand, we next examined reaction parameters to identify optimal conditions for reactivity and enantioselectivity.

Our initial experiments revealed that treatment of allyl 1-benzyl-2-oxocyclobutanecarboxylate (**4l**) with catalytic [Pd₂(pmdba)₃] in the presence of (*S*)-*t*BuPHOX (**L1**) in THF delivered the desired α -quaternary (*S*)-2-allyl-2-benzylcyclobutanone (**5l**)^[18] in 90% yield, albeit in moderate enantioselectivity (Table 1, entry 1). The use of electron-deficient ligands resulted in considerably improved enantioinduction (Table 1, entries 2 and 5). Although the reaction proceeds well in a number of solvents, toluene was identified as optimal for inducing asymmetry. This solvent effect is likely due to an

Table 1: Reaction optimization studies.^[a]

Entry	Ligand	Solvent	T [°C]	ee [%]
1	L1	THF	25	58
2	L2	THF	25	75
3	L2	<i>p</i> -dioxane	25	84
4	L2	benzene	25	84
5	L3	toluene	25	84
6	L2	toluene	25	85
7	L2	toluene	20	86

[a] Conditions: cyclobutanone **4l** (1.0 equiv), [Pd₂(pmdba)₃] (5 mol %), Ligand (**L1**–**L3**) (12.5 mol %) in stated solvent (0.033 M) at indicated temperature for 12–48 h. ee determined by chiral supercritical fluid chromatography (SFC). pmdba = 4,4'-dimethoxydibenzylidene acetone.



enhanced binding between the enolate and the electrophilic α -allyl-Pd^{II} center in the catalytic cycle, which may reinforce a tight ion pair and lead to an inner-sphere mechanism.^[19] Finally, at temperatures just below ambient the reaction was found to proceed at a reasonable rate and with high enantioselectivity.

With these optimized conditions identified, we next explored the influence of different α -substituents (R¹, Table 2A) on the efficacy of the allylic alkylation process. To aid in the isolation of these highly volatile compounds, we chose β -ketoesters of higher molecular weights, bearing substitution at both the α -position and allyl fragment (i.e. 2-phenylallyl 1-benzyl-2-oxocyclobutanecarboxylate) as base substrates. We were pleased to find that α -alkyl substituents were well tolerated with enantiomeric excess up to 99% (Table 2A; **5a**, **5b**). α -Benzyl substituents were found to give the respective α -quaternary cyclobutanones with uniformly excellent enantioselectivity regardless of the electronic nature of the benzyl moiety (**5c**, **5e**). In addition to alkyl and benzyl

Table 2: Catalytic enantioselective cyclobutanone alkylation: A) scope of α -keto substituent tolerance; B) functional diversity incorporated at the 2-allyl position.^[a]

A)

5a , 62% yield 99% ee	5b , 71% yield 90% ee	5c , 89% yield 95% ee
5d , 71% yield 93% ee	5e , 83% yield 97% ee	5f , 86% yield 92% ee
5g , 90% yield 93% ee	5h , 82% yield 92% ee	

B)

5i , 92% yield 91% ee	5j , 82% yield 90% ee	5k , 67% yield 93% ee	5l , 72% yield 86% ee
5m , 95% yield 89% ee	5n , 91% yield 92% ee	5o , 95% yield 94% ee	


[a] Conditions: cyclobutanone **4** (1.0 equiv), [Pd₂(pmdba)₃] (5 mol %), (*S*)-**L2** (12.5 mol %) in toluene (0.033 M) at 20 °C for 12–48 h. All reported yields are for isolated products.

substituents, allyl-, TMS-protected propargyl- and heteroaryl-substituted 2-carboxyallyl cyclobutanones proved to be eligible substrates in the asymmetric allylic alkylation reaction providing cyclobutanones **5f–5h** in high yields and enantiomeric excess.

Having surveyed the scope of the process with respect to various substituents at the quaternary center, we were poised to investigate the influence of different allyl substitution on the process (R^2 , Table 2B). In accord with previous studies on the palladium-catalyzed asymmetric allylic alkylation, the catalytic system was found to be relatively inactive when terminally substituted or cyclic allyl fragments were employed.^[14a,b] As such, we limited our survey to carboxyallyl fragments bearing substituents at the 2-allyl position. Gratifyingly, diverse substituents were found to be well tolerated (Table 2B). All α -quaternary cyclobutanones were obtained in moderate to high yield and with outstanding enantiopurity. Particularly interesting are compounds **5i**, **5k**, and **5m** featuring a butadiene, a vinyl chloro, and a benzyl ether moiety, respectively. Each of these diverse functional groups may potentially serve as handles for various derivatization reactions (e.g., cycloaddition, annulation or transition metal-catalyzed cross-coupling).

Myriad studies have shown cyclobutanoids to be highly valuable synthetic intermediates, allowing access to enantioenriched oxazepines,^[20] piperidines,^[21] tetrahydropyrans,^[22] α - and β -quaternary cyclopentanones,^[10] benzannulated polycycles,^[23] as well as β -quaternary linear ketones.^[2] Cyclobutanones may participate directly in a variety of robust classical transformations, such as Baeyer–Villiger oxidation and Beckmann rearrangement,^[2] as well as transition metal-catalyzed ring expansion,^[24] ring contraction^[25] and ring-opening processes.^[26] To demonstrate the utility of our asymmetric synthesis of cyclobutanones within this domain, we carried out a number of transformations on the chiral cyclobutanones generated in this study. Ring expansion by Baeyer–Villiger oxidation, treatment with trimethylsilyldiazomethane and Beckmann rearrangement all proceeded smoothly to deliver dialkyl γ -lactone **6**, α -quaternary cyclopentanone **7**, and dialkyl γ -lactam **8**, respectively. Additionally, ring closing metathesis of diallyl-substituted cyclobutanone **5f** cleanly furnished quaternary [4.5]-spirocycle **9** (Scheme 2).

In summary, we have developed the first transition metal-catalyzed enantioselective α -alkylation of cyclobutanones. This method employs palladium catalysis and an electron-deficient PHOX-type ligand to afford α -quaternary cyclobutanones in good to excellent yields and enantioselectivities. A wide variety of substituents are tolerated at both the α -keto and 2-allyl positions. The mild nature of our method is reflected in its compatibility with otherwise highly electrophilic cyclobutanones. We have further demonstrated the utility of chiral cyclobutanones as synthetic building blocks to access a variety of enantioenriched derivative compounds including dialkyl γ -lactams, dialkyl γ -lactones, α -quaternary cyclopentanones, and quaternary [4.5]-spirocycles. We believe that this novel synthetic method will enable the expeditious synthesis of complex bioactive natural products and pharmaceutical components by providing unique access to previously



Scheme 2. Derivatization of enantioenriched α -quaternary cyclobutanones. Conditions: a) **5c**, H_2O_2 (55 wt% in H_2O), 1 M NaOH, MeOH, 23 °C, 80% yield. b) **5g**, $TMSCHN_2$, $BF_3 \cdot Et_2O$, Et_2O , HCl (aq.), DCM, 69% yield, two steps. c) **5o**, $HONH_2 \cdot HCl$, pyridine, EtOH; p -TsCl, Et_3N , DMAP, DCM, 22% yield, two steps. d) **5f**, Grubbs–Hoveyda G2, PhH, 97% yield.

unknown and inaccessible enantioenriched α -quaternary cyclobutanones. Efforts toward this end are currently underway in our laboratory.

Received: March 4, 2013

Revised: April 17, 2013

Published online: May 17, 2013

Keywords: allylic alkylation · asymmetric catalysis · cyclobutanone · palladium · quaternary center

- [1] For general reviews see: a) D. Cain in *Carbon-Carbon Bond Formation*, Vol. 5 (Ed.: R. L. Augustine), Marcel Dekker, New York, **1979**, pp. 85–250; b) D. Seebach, *Angew. Chem.* **1988**, *100*, 1685–1715; *Angew. Chem. Int. Ed. Engl.* **1988**, *27*, 1624–1654; c) D. L. Hughes in *Comprehensive Asymmetric Catalysis*, Vol. 2 (Eds.: E. N. Jacobsen, A. Pfaltz, H. Yamamoto), Springer, Berlin, **1999**, pp. 1273–1294.
- [2] a) D. Belluš, B. Ernst, *Angew. Chem.* **1988**, *100*, 820–850; *Angew. Chem. Int. Ed. Engl.* **1988**, *27*, 797–827; b) A. Sergeiko, V. V. Poroikov, L. O. Hanus, V. M. Dembitsky, *Open Med. Chem. J.* **2008**, *2*, 26–37; c) V. M. Dembitsky, *J. Nat. Med.* **2008**, *62*, 1–33; d) K. Yoshikawa, A. Kaneko, Y. Matsumoto, H. Hama, S. Arihara, *J. Nat. Prod.* **2006**, *69*, 1267–1270; e) J. Marrero, A. D. Rodríguez, P. Baran, R. G. Raptis, J. A. Sánchez, E. Ortega-Barria, T. L. Capson, *Org. Lett.* **2004**, *6*, 1661–1664; f) Y.-S. Li, K. Matsunaga, M. Ishibashi, Y. Ohizumi, *J. Org. Chem.* **2001**, *66*, 2165–2167; g) N. Tanaka, M. Okasaka, Y. Ishimaru, Y. Takaishi, M. Sato, M. Okamoto, T. Oshikawa, S. U. Ahmed, L. M. Consentino, K.-H. Lee, *Org. Lett.* **2005**, *7*, 2997–2999; h) B. Jiang, J. Qiao, Y. Yang, Y. Lu, *J. Cell. Biochem.* **2012**, *113*, 2560–2566.
- [3] a) T. Seiser, T. Saget, D. N. Tran, N. Cramer, *Angew. Chem.* **2011**, *123*, 7884–7896; *Angew. Chem. Int. Ed.* **2011**, *50*, 7740–7752; b) N.-Y. Fu, S.-H. Chan, H. N. C. Wong in *Patai Series: the Chemistry of Functional Groups* (Eds.: Z. Rappoport, J. F. Liebman, S. Patai), Wiley, Chichester, **2005**, pp. 357–440.
- [4] A. Bauzá, D. Quiñero, P. M. Deyà, A. Frontera, *Chem. Phys. Lett.* **2012**, *536*, 165–169.
- [5] K. B. Wiberg in *Patai Series: the Chemistry of Functional Groups* (Eds.: Z. Rappoport, J. F. Liebman, S. Patai), Wiley, Chichester, **2005**, pp. 1–15.
- [6] N. L. Bauld, J. Cessac, R. L. Holloway, *J. Am. Chem. Soc.* **1977**, *99*, 8140–8144.

- [7] U. Berg in *Patai Series: the Chemistry of Functional Groups* (Eds.: Z. Rappoport, J. F. Liebman, S. Patai), Wiley, Chichester, **2005**, pp. 83–131.
- [8] a) J.-P. Pete, *Advances in Photochemistry*, Wiley, Hoboken, **1996**; b) J. P. Hehn, C. Mueller, T. Bach in *Handbook of Synthetic Photochemistry* (Eds.: A. Albini, M. Fagnoni), Wiley-VCH, Weinheim, **2010**, pp. 171–215; c) N. Hoffmann in *Handbook of Synthetic Photochemistry* (Eds.: A. Albini, M. Fagnoni), Wiley-VCH, Weinheim, **2010**, pp. 137–169; d) E. Lee-Ruff in *Methods of Organic Chemistry, Vol. E17e* (Ed.: A. De Meijere), Thieme, Stuttgart, **1997**, pp. 179–187.
- [9] a) L. Fitjer in *Methods of Organic Chemistry Vol. E17e* (Ed.: A. De Meijere), Thieme, Stuttgart, **1997**, pp. 251–316; b) E. Lee-Ruff in *Patai Series: the Chemistry of Functional Groups* (Eds.: Z. Rappoport, J. F. Liebman, S. Patai), Wiley, Chichester, **2005**, pp. 281–355.
- [10] M. Chaumontet, R. Piccardi, N. Audic, J. Hitce, J.-L. Peglion, E. Clot, O. Baudoin, *J. Am. Chem. Soc.* **2008**, *130*, 15157–15166.
- [11] a) J. P. Markham, S. T. Staben, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 9708–9709; b) F. Kleinbeck, F. D. Toste, *J. Am. Chem. Soc.* **2009**, *131*, 9178–9179.
- [12] V. López-Carrillo, A. M. Echavarren, *J. Am. Chem. Soc.* **2010**, *132*, 9292–9294.
- [13] a) E. Lee-Ruff, G. Mladenova, *Chem. Rev.* **2003**, *103*, 1449–1484; b) B. M. Trost, T. Yasukata, *J. Am. Chem. Soc.* **2001**, *123*, 7162–7165; c) H. Nemoto, H. Ishibashi, M. Nagamochi, K. Fukumoto, *J. Org. Chem.* **1992**, *57*, 1707–1712.
- [14] a) Cursory studies on substrates bearing terminally substituted and cyclic allyl fragments were conducted. Subjecting compounds **10** and **11** to the reaction conditions described in Scheme 2 resulted in minimal conversion to the desired products (< 5% and < 1%, respectively, as determined by ¹H NMR of the crude reaction mixtures) and these substrate classes were not pursued further.
- [15] N. T. McDougal, S. C. Virgil, B. M. Stoltz, *Synlett* **2010**, 1712–1716.
- [16] J. S. Baum, D. A. Shook, H. M. L. Davies, H. D. Smith, *Synth. Commun.* **1987**, *14*, 1709–1716.
- [17] M. Presset, Y. Coquerel, J. Rodriguez, *J. Org. Chem.* **2009**, *74*, 415–418.
- [18] Derivatization of the allylic alkylation product 2-allyl-2-methylcyclobutanone (**13**) to known compound (5*S*)-5-allyl-5-methyl-dihydrofuran-2(3*H*)-one (**14**) allowed for comparison of optical rotation data and confirmation of the (5*S*) absolute configuration (see Supporting Information). The absolute stereoconfiguration of all other products generated in this study were assigned by analogy. These data are consistent with the absolute configurations observed in previous studies on 5-, 6- and 7-membered rings (see Ref. [14]).

