


Angewandte Corrigendum

In Scheme 1 of this communication, the structure of notoamide C (**3**) should be corrected as shown below.



The configuration at C3 of **3** was not determined by spectroscopic methods and the co-occurrence of notoamides A and B indicated the 3*R* configuration of **3** based on biogenetic considerations. Recently, the absolute configuration of notoamide C was corrected to be 3*S* based on the biochemical conversion of notoamide E into notoamide C by recombinant NotB.^[1] This correction affects the structure that appeared in Figure S3 of the Supporting Information. Recently, the relative and absolute configuration of notoamide C was independently reported by Chen et al. on the basis of a single crystal X-ray analysis of notoamide C isolated from *Aspergillus* sp. XS-20090066.^[2]

- [1] S. Li, J. M. Finefield, J. D. Sunderhaus, T. J. McAfoos, R. M. Williams, D. H. Sherman, *J. Am. Chem. Soc.* **2012**, *134*, 788–791.
- [2] M. Chen, C.-L. Shao, X.-M. Fu, R.-F. Xu, J.-J. Zheng, D.-L. Zhao, Z.-G. She, C.-Y. Wang, *J. Nat. Prod.* **2013**, *76*, 547–553.

Notoamides A–D: Prenylated Indole Alkaloids Isolated from a Marine-Derived Fungus, *Aspergillus* sp.

H. Kato, T. Yoshida, T. Tokue, Y. Nojiri, H. Hirota, T. Ohta, R. M. Williams, S. Tsukamoto* _____ **2254–2256**

Angew. Chem. Int. Ed. **2007**, *46*

DOI: 10.1002/anie.200604381