


# Arylation of Diazoesters by a Transient N–H Insertion Organocascade\*\*

Tyler J. Auvil, Sonia S. So, and Anita E. Mattson\*


Cascade catalysis is a powerful tool demonstrated both by nature and chemists alike to generate complex products through sequential catalytic processes.<sup>[1]</sup> These processes benefit from the efficiency of catalyzing multiple steps in a single flask for the rapid construction of molecular complexity without the necessity of intermediate isolation. Autotandem catalytic methods are cascade processes employed to promote two or more distinct reactions with a single catalyst.<sup>[2]</sup> Inspired by the demonstrated power of autotandem organocascade catalysis,<sup>[3]</sup> we sought to invoke a fundamentally novel sequence taking advantage of catalytic transient carbene activation. More specifically, we envisioned activating a species having carbene character by a urea-catalyzed<sup>[4]</sup> insertion reaction with an appropriate carbene activator to generate a transient intermediate which would then enter into a second distinct cycle for further functionalization (Scheme 1). Successful development of this process would



**Scheme 1.** Transient carbene activation cascade catalysis.

afford a straightforward organocatalytic route to synthetically challenging structural motifs from simple starting materials, and also expose the potential of transient carbene insertion reactions as innovative tools in organocatalytic method development. Herein, we detail our breakthrough discoveries in urea-catalyzed transient N–H insertion reactions for the controlled functionalization of nitrodiazoesters.

The geminal diaryl functionality is frequently found in bioactive target molecules, and several drugs already commercially available to treat a variety of diseases contain this structural motif.<sup>[5]</sup> Specific examples include the antidepressant Zoloft, and anticancer agents Femara and Etoposide. Aware of the therapeutic significance and synthetic challenges associated with preparing unsymmetrical geminal diaryl frameworks,<sup>[6]</sup> we focused the development of transient insertion organocascade catalysis on arylation reactions of diazo compounds. The working hypothesis was that an appropriate amine could trigger urea-catalyzed arylation reactions of diazo compounds (**1**)<sup>[7]</sup> through transient N–H insertion intermediates<sup>[8,9]</sup> (**4**; Scheme 2). To the best of our



**Scheme 2.** Double arylation of nitrodiazoesters through transient N–H insertion organocascade catalysis.

knowledge, cascade reactions proceeding through transient carbene insertion intermediates had not been reported at the onset of our investigations. The testing of the feasibility of our hypothesis began with investigating anilines as carbene activators of nitrodiazoesters.<sup>[10]</sup>

Our studies began with the preparation of **2a** by the double arylation of ethyl nitrodiazoacetate (**1a**) with 5-methoxyindole and 5-bromoindole in the presence of 20 mol % of the boronate urea **3a**<sup>[11,12]</sup> and an aniline activator (Table 1). Our first attempt afforded a disappointing 12 % of the desired adduct **2a** when aniline was employed as the carbene activator (entry 1). Further examination of the reaction led us to identify participation of the aniline activator in the diarylation reaction. Next, we turned to *p*-substituted anilines, activators less likely to be involved in undesired arylation pathways. Indeed, the strategic choice of activators proved fruitful and we were delighted to find that activation of **1a** with *p*-anisidine gave rise to 73 % of **2a** (entry 2). Additional tuning of the aniline led us to 4-fluoroaniline as an even better carbene activator (entry 3), potentially because of its better leaving ability. The necessity of an aniline activator able to undergo N–H insertion with the diazoester **1a** is clear: a 0 % yield of **2a** was observed when 2,6-dimethylaniline, an aniline too sterically encumbered to undergo N–H insertion,

[\*] T. J. Auvil, S. S. So, Prof. Dr. A. E. Mattson  
Department of Chemistry and Biochemistry  
The Ohio State University  
100 W. 18th Ave., Columbus, OH 43210 (USA)  
E-mail: mattson@chemistry.ohio-state.edu  
Homepage: <http://mattson.group.chemistry.ohio-state.edu/>

[\*\*] The Ohio State University Department of Chemistry and Biochemistry and Donors of the American Chemical Society Petroleum Research Fund are gratefully acknowledged for supporting this research. Prof. J. Fox (University of Delaware) is thanked for insightful discussions.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201304921>.

**Table 1:** Study of aniline activators in transient N–H insertion catalysis.<sup>[a]</sup>

| Entry                    | 1  | 2  | 3 <sup>[c]</sup> | 4 | 5    |
|--------------------------|----|----|------------------|---|------|
| Ar-NH <sub>2</sub>       |    |    |                  |   | none |
| Yield [%] <sup>[b]</sup> | 12 | 73 | 78               | 0 | 0    |

[a] Reactions performed at 2 M in DCE. [b] Yield of isolated product.


[c] 0% of **2a** was generated in the absence of **3a**.

was tested as the activator (entry 4). From the results of two critical control experiments, we concluded that the urea catalyst and aniline activator working in combination are uniquely able to effect the double arylation of **1a**. Specifically, the first control experiment resulted in no desired **2a** when an N–H insertion activator was excluded from the reaction system (entry 5). Similarly, in the second control, none of **2a** was observed if the boronate urea **3a** was excluded from the optimized reaction conditions (entry 3).

A plausible organocascade for the unsymmetrical diarylation of nitrodiazoesters, depicted in Scheme 3, takes advantage of the urea catalyst (**3a**) in two sequential roles: 1) nitrodiazoester activation for N–H insertion/multicomponent coupling and 2)  $\alpha$ -amino ester activation for arylation. In the first cycle, **1a** is activated by **3a** and 4-fluoroaniline, thus affording the N–H insertion adduct **B** through the hydrogen-bonded complex **A**. Upon liberation of nitrous acid, **B** is converted into the iminoglyoxylate **C**. The arylation of the activated iminoglyoxylate then generates the transient N–H insertion intermediate **4a**.<sup>[13]</sup> In Cycle II, the  $\alpha$ -amino ester is


activated through dual binding with the urea catalyst, conceivably through the complex **D**. The urea-assisted substitution of the N-bound 4-fluoroaniline with 5-bromoindole then generates the desired double arylated ester product **2a** upon aromatization, thus introducing **3a** and 4-fluoroaniline back into the catalytic cycle.<sup>[14,15]</sup>

In support of the proposed reaction pathway, the enantioenriched transient N–H insertion intermediate **4a** was isolated and subjected to the reaction conditions (Scheme 4).<sup>[13]</sup> From this experiment we found that **4a** is



**Scheme 4.** Urea-catalyzed arylation of the enantioenriched **4a**.

converted into a racemic mixture of **2a** in 91% yield under the influence of 10 mol% **3a**. Importantly, the urea is essential for this reaction: no reaction was observed if the urea catalyst was excluded. The loss of enantiomeric excess in the conversion of **4a** into **2a** suggests that an S<sub>N</sub>1-like mechanism is operating for the conversion of **D** into **E**. Additional control experiments conducted in our laboratory suggest the stereocenter in the di- $\alpha$ -arylester **2** is stable to the reaction conditions, thus further supporting an S<sub>N</sub>1-like reaction pathway as opposed to an S<sub>N</sub>2 and subsequent racemization.<sup>[16]</sup> Stoichiometric amounts of inexpensive 4-fluoroaniline were necessary to prevent the generation of undesired symmetrical  $\alpha$ -diarylated esters, which were




**Scheme 3.** Plausible reaction pathway for the unsymmetric double arylation of nitrodiazoesters through a transient N–H insertion intermediate.

formed when using catalytic quantities of the carbene activator. However, the feasibility of transient insertion cascades requiring only catalytic amounts of the carbene activator is a promising avenue for future studies not featuring double functionalization.

The autotandem catalytic double arylation of **1a** was easily extended to a remarkably wide array of substrates to give rise to the di- $\alpha$ -aryl esters **2** in excellent yields (Scheme 5). Unsymmetrical double indolylolation was efficiently achieved in up to 97 % yield when indoles were used as both the first and second nucleophiles (**2a–e**). Anilines were also easily incorporated into the diarylation reaction. Examples of indoles working in combination with anilines afforded the products **2f–j** in excellent yields of up to 98 % yield. A variety of anilines, as both the first and second nucleophiles, are excellent reactants for this coupling reaction as well. Even sterically encumbered, electron-rich anilines were well tolerated in the diarylation reaction. For instance, 3,5-dimethoxyaniline gave rise to 98 % of **2n**. Select halogenated anilines were capable of being integrated into the double arylation sequence, albeit longer reaction times were required for step 2. *o*-Iodoaniline gave rise to 56 % of **2o** after 72 hours at 60 °C. Additionally, pyrroles operated moderately well in the cascade sequence (**2q,r**).

The reaction also proved tolerant of a wide range of nitro diazoesters (**1**; Table 2). Sterically demanding nitrodi-



**Scheme 5.** Substrate scope for the double arylation of ethyl nitro diazoacetate.<sup>[a,b]</sup> [a] Reactions performed at 2 M in DCE in step 2. See the Supporting Information for details [b] Yield of isolated products. [c] 72 h reaction time in step 2. DCE = 1,2-dichloroethane.

**Table 2:** Diazoester substrate scope.<sup>[a]</sup>

| Entry            | Diazoester | <b>1</b>  | Product   | Yield [%] <sup>[b]</sup> |
|------------------|------------|-----------|-----------|--------------------------|
| 1                |            | <b>1b</b> | <b>2s</b> | 88                       |
| 2                |            | <b>1c</b> | <b>2t</b> | 75                       |
| 3                |            | <b>1d</b> | <b>2u</b> | 63                       |
| 4                |            | <b>1e</b> | <b>2v</b> | 60                       |
| 5 <sup>[c]</sup> |            | <b>1f</b> | <b>2w</b> | 68                       |

[a] Reactions performed at 2 M in DCE in step 2. [b] Yield of isolated product. [c] **2w** isolated as a 1:1 mixture of diastereomers.

azoacetates (**1b–d**) were well tolerated, thus providing the products **2s–u** in up to 88 % yield. Substituted benzyl esters were also easily incorporated into the double arylation sequence. For instance, the product **2v** was prepared in 60 % yield from the diazoester **1e** (entry 4). The (–)-menthol-derived nitro diazoester **1f** gave rise to **2w** in 68 % yield as a 1:1 mixture of diastereomers.

In summary, we have discovered that the intermolecular double arylation of nitro diazoesters is achieved through a unique urea-catalyzed transient N–H insertion organocascade. The reaction can accommodate a remarkable number of anilines and indoles, thus enabling the efficient preparation of a large family of unsymmetric  $\alpha$ -diaryl esters. The proposed reaction pathway invokes the urea catalyst in two distinct, sequential catalytic cycles: N–H insertion and  $\alpha$ -arylation. We anticipate transient insertion catalysis may be a fruitful new direction in organocatalysis. Ongoing investigations in our laboratory are focused on exploring the possibility of enantioselective transient N–H insertion catalysis.

Received: June 7, 2013

Revised: July 12, 2013

Published online: August 26, 2013

**Keywords:** carbenes · diazo compounds · insertion · organocatalysis · synthetic methods

- [1] For recent reviews on cascade catalysis, see: a) Z. Du, Z. Shao, *Chem. Soc. Rev.* **2013**, 42, 1337–1378; b) H. Pellissier, *Chem. Rev.* **2013**, 113, 442–524; c) N. Patil, V. Shinde, B. Gajula, *Org. Biomol. Chem.* **2012**, 10, 211–224; d) L. Ambrosini, T. Lambert,

- ChemCatChem* **2010**, *2*, 1373–1380; e) J. Zhou, *Chem. Asian J.* **2010**, *5*, 422–434; f) K. Nicolaou, D. Edmonds, P. Bulger, *Angew. Chem.* **2006**, *118*, 7292–7344; *Angew. Chem. Int. Ed.* **2006**, *45*, 7134–7186; g) A. Bruggink, R. Schoevarrt, T. Kieboom, *Org. Process Res. Dev.* **2003**, *7*, 622–640; For recent reviews on organocascade catalysis, see: h) R. Wende, P. Schreiner, *Green Chem.* **2012**, *14*, 1821–1849; i) S. Jones, B. Simmons, A. Mastracchio, D. MacMillan, *Nature* **2011**, *475*, 183–188; j) C. Grondal, M. Jeanty, D. Enders, *Nat. Chem.* **2010**, *2*, 167–178; k) X. Yu, W. Wang, *Org. Biomol. Chem.* **2008**, *6*, 2037–2046; For examples of organocascade catalysis, see: l) B. Horning, D. MacMillan, *J. Am. Chem. Soc.* **2013**, *135*, 6442–6445; m) X. Zeng, Q. Ni, G. Raabe, D. Enders, *Angew. Chem.* **2013**, *125*, 3050–3054; *Angew. Chem. Int. Ed.* **2013**, *52*, 2977–2980; n) X. Tan, P. Melchiorre, *Angew. Chem.* **2013**, *125*, 5468–5471; *Angew. Chem. Int. Ed.* **2013**, *52*, 5360–5363; o) B. Tan, N. Candeias, C. Barbas III, *Nat. Chem.* **2011**, *3*, 473–477; p) Z. Jia, H. Jiang, B. Gschwend, Q. Li, X. Yin, J. Grouleff, Y. Chen, K. Jørgensen, *J. Am. Chem. Soc.* **2011**, *133*, 5053–5061; q) H. Ishikawa, M. Honma, Y. Hayashi, *Angew. Chem.* **2011**, *123*, 2876–2879; *Angew. Chem. Int. Ed.* **2011**, *50*, 2824–2827; r) H. Uehara, R. Imashiro, G. Hernández-Torres, C. Barbas III, *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 20672–20677; s) X. Zhang, S. Zjang, W. Wang, *Angew. Chem.* **2010**, *122*, 1523–1526; *Angew. Chem. Int. Ed.* **2010**, *49*, 1481–1484; t) B. Simmons, A. Walji, D. MacMillan, *Angew. Chem.* **2009**, *121*, 4413–4417; *Angew. Chem. Int. Ed.* **2009**, *48*, 4349–4353; u) J. Zhou, B. List, *J. Am. Chem. Soc.* **2007**, *129*, 7498–7499; v) M. Rueping, A. Antonchick, T. Theissmann, *Angew. Chem.* **2006**, *118*, 3765–3768; *Angew. Chem. Int. Ed.* **2006**, *45*, 3683–3686; w) Y. Huang, A. Walji, C. Larsen, D. MacMillan, *J. Am. Chem. Soc.* **2005**, *127*, 15051–15053.
- [2] For recent reviews on tandem catalysis, see: a) N. Shindoh, Y. Takemoto, K. Takasu, *Chem. Eur. J.* **2009**, *15*, 12168–12179; b) D. Fogg, E. dos Santos, *Coord. Chem. Rev.* **2004**, *248*, 2365–2379; c) J. Wasilke, S. Obrey, R. Baker, G. Bazan, *Chem. Rev.* **2005**, *105*, 1001–1020.
- [3] For recent examples of autotandem catalysis, see: a) Q. Nguyen, T. Nguyen, T. Driver, *J. Am. Chem. Soc.* **2013**, *135*, 620–623; b) N. Kanbayashi, K. Takenaka, T. Okamura, K. Onitsuka, *Angew. Chem. Int. Ed.* **2013**, *52*, 4897–4901; c) Y. Yang, W. Shu, S. Yu, F. Ni, M. Gao, A. Wu, *Chem. Commun.* **2013**, *49*, 1729–1731; d) A. Yadav, S. Verbeek, S. Hostyn, P. Franck, S. Sergeyev, B. Maes, *Org. Lett.* **2013**, *15*, 1060–1063; e) J. Clark, J. Griffiths, S. Diver, *J. Am. Chem. Soc.* **2013**, *135*, 3327–3330; f) K. Takasu, T. Tanaka, T. Azuma, Y. Takemoto, *Chem. Commun.* **2010**, *46*, 8246–8248.
- [4] For information on urea and thiourea catalysis, see: a) Y. Takemoto, *Chem. Pharm. Bull.* **2010**, *58*, 593–601; b) Z. Zhang, P. Schreiner, *Chem. Soc. Rev.* **2009**, *38*, 1187–1198; c) S. Connon, S. Kavanagh, A. Piccinini, *Org. Biomol. Chem.* **2008**, *6*, 1339–1343; d) A. Berkessel, H. Groger in *Asymmetric Organocatalysis: From Biomimetic Concepts to Applications in Asymmetric Synthesis*, Wiley-VCH, Weinheim, **2005**; e) Y. Take-moto, *Org. Biomol. Chem.* **2005**, *3*, 4299–4306.
- [5] For examples of the geminal diaryl moiety in bioactive molecules, see: a) S. Yokoshima, T. Ueda, S. Kobayashi, A. Sato, T. Kuboyama, H. Tokuyama, T. Fukuyama, *J. Am. Chem. Soc.* **2002**, *124*, 2137–2139; b) L. Nilvebrant, K. Andersson, P. Gillberg, M. Stahl, B. Sparf, *Eur. J. Pharmacol.* **1997**, *327*, 195–207; c) B. Hughes, D. Howat, H. Lisle, M. Holbrook, T. James, N. Gozzard, K. Bleas, P. Hughes, R. Kingbay, G. Warrellow, R. Alexander, J. Head, E. Boyd, M. Eaton, M. Perry, M. Wales, B. Smith, R. Owens, C. Catterall, S. Lumb, A. Russell, R. Allen, M. Merriman, D. Bloxham, D. Bloxham, G. Higgs, *Br. J. Pharmacol.* **1996**, *118*, 1183–1191; d) J. Lowe III, D. Hageman, S. Drozda, S. McLean, D. Bryce, R. Crawford, S. Zorn, J. Morrone, J. Bordner, *J. Med. Chem.* **1994**, *37*, 3789–3811; e) K. Bogeso, A. Christensen, J. Hyttel, T. Liljefors, *J. Med. Chem.* **1985**, *28*, 1817–1828.
- [6] For recent reviews on transition metal catalyzed  $\alpha$ -arylation reactions to synthesize the geminal diaryl functionality, see: a) O. Baudoin, *Chem. Soc. Rev.* **2011**, *40*, 4902–4911; b) C. Johansson, T. Colacot, *Angew. Chem.* **2010**, *122*, 686–718; *Angew. Chem. Int. Ed.* **2010**, *49*, 676–707; c) F. Bellina, R. Rossi, *Chem. Rev.* **2010**, *110*, 1082–1146; d) D. Culkin, J. Hartwig, *Acc. Chem. Res.* **2003**, *36*, 234–245; For examples of syntheses of geminal diaryl compounds, see: e) R. Martín, S. Buchwald, *Angew. Chem.* **2007**, *119*, 7374–7377; *Angew. Chem. Int. Ed.* **2007**, *46*, 7236–7239; f) N. Beare, J. Hartwig, *J. Org. Chem.* **2002**, *67*, 541–555; g) O. Gaertzen, S. Buchwald, *J. Org. Chem.* **2002**, *67*, 465–475; h) W. Moradi, S. Buchwald, *J. Am. Chem. Soc.* **2001**, *123*, 7996–8002; i) S. Stauffer, N. Beare, J. Stambuli, J. Hartwig, *J. Am. Chem. Soc.* **2001**, *123*, 4641–4642; j) H. Davies, D. Stafford, T. Hansen, *Org. Lett.* **1999**, *1*, 233–236; k) D. Old, J. Wolfe, S. Buchwald, *J. Am. Chem. Soc.* **1998**, *120*, 9722–9723.
- [7] For recent reviews on diazo compound insertion chemistry, see: a) D. Morton, H. Davies, *Chem. Soc. Rev.* **2011**, *40*, 1857–1869; b) C. Slattery, A. Ford, A. Maguire, *Tetrahedron* **2010**, *66*, 6681–6705; c) C. Moody, *Angew. Chem.* **2007**, *119*, 9308–9310; *Angew. Chem. Int. Ed.* **2007**, *46*, 9148–9150; d) H. Davies, R. J. Beckwith, *Chem. Rev.* **2003**, *103*, 2861–2903.
- [8] For a recent example of urea-catalyzed N–H insertion reactions of nitrodiazoesters, see: S. So, A. Mattson, *J. Am. Chem. Soc.* **2012**, *134*, 8798–8801.
- [9] For recent examples of diazo compounds undergoing N–H insertion reactions, see: a) S. Hansen, J. Spangler, J. Hansen, H. Davies, *Org. Lett.* **2012**, *14*, 4626–4629; b) S. Zhu, B. Xu, G. Wang, Q. Zhou, *J. Am. Chem. Soc.* **2012**, *134*, 436–442; c) B. Xu, S. Zhu, X. Xie, J. Shen, Q. Zhou, *Angew. Chem.* **2011**, *123*, 11685–11688; *Angew. Chem. Int. Ed.* **2011**, *50*, 11483–11486; d) E. Lee, G. Fu, *J. Am. Chem. Soc.* **2007**, *129*, 12066–12067; e) B. Lui, S. Zhu, W. Zhang, C. Chen, Q. Zhou, *J. Am. Chem. Soc.* **2007**, *129*, 5834–5835.
- [10] For examples of transition metal catalyzed approaches toward controlling  $\alpha$ -nitro- $\alpha$ -diazocarbonyl reactivity, see: a) S. Vanier, G. Larouche, R. Wurz, A. Charette, *Org. Lett.* **2010**, *12*, 672–675; b) V. Lindsay, W. Lin, A. Charette, *J. Am. Chem. Soc.* **2009**, *131*, 16383–16385; c) S. Zhu, J. Perman, X. Zhang, *Angew. Chem.* **2008**, *120*, 8588–8591; *Angew. Chem. Int. Ed.* **2008**, *47*, 8460–8463; d) R. Wurz, A. Charette, *J. Org. Chem.* **2004**, *69*, 1262–1269; e) P. O'Bannon, W. Dailey, *J. Org. Chem.* **1989**, *54*, 3096–3101; f) A. Charette, R. Wurz, T. Ollevier, *Helv. Chim. Acta* **2002**, *85*, 4468–4484.
- [11] For boronate ureas as enhanced hydrogen-bond-donor catalysts, see: a) D. Nickerson, V. Angeles, T. Auvil, S. So, A. Mattson, *Chem. Commun.* **2013**, *49*, 4289–4291; b) S. So, T. Auvil, V. Garza, A. Mattson, *Org. Lett.* **2012**, *14*, 444–447; c) S. So, J. Burkett, A. Mattson, *Org. Lett.* **2011**, *13*, 716–719.
- [12] For pioneering work on boronate ureas in anion recognition, see: a) M. Hughes, B. Smith, *J. Org. Chem.* **1997**, *62*, 4492–4499; b) M. Hughes, M. Shang, B. Smith, *J. Org. Chem.* **1996**, *61*, 4510–4511.
- [13] The N–H insertion component of the proposed organocascade has previously been demonstrated to be catalyzed by **3a**. See Ref. [8].
- [14] For examples of Lewis acid catalyzed substitutions of anilines, see: a) G. Desimoni, G. Faita, M. Mariella, M. Toscanini, M. Biocchi, *Eur. J. Org. Chem.* **2008**, 6232–6238; b) S. Shirakawa, S. Kobayashi, *Org. Lett.* **2006**, *8*, 4939–4942.
- [15] For an example of a thiourea catalyst stabilizing alkoxides through proposed partial protonation by means of double hydrogen bonding, see: M. Kotke, P. R. Schreiner, *Tetrahedron* **2006**, *62*, 434–439.
- [16] See the Supporting Information for details.