

Rhodium-Catalyzed Asymmetric Synthesis of Silicon-Stereogenic Dibenzosiloles by Enantioselective [2+2+2] Cycloaddition**

Ryo Shintani,* Chihiro Takagi, Tomoaki Ito, Masanobu Naito, and Kyoko Nozaki*


Abstract: A rhodium-catalyzed asymmetric synthesis of silicon-stereogenic dibenzosiloles has been developed through a [2+2+2] cycloaddition of silicon-containing prochiral triynes with internal alkynes. High yields and enantioselectivities have been achieved by employing an axially chiral monophosphine ligand, and the present catalysis is also applicable to the asymmetric synthesis of a germanium-stereogenic dibenzogermole. Preliminary studies on the optical properties of these compounds are also described.

Dibenzosiloles are widely found as a structural motif in various useful materials,^[1] such as light-emitting diodes,^[2] thin-film transistors,^[3] solar cells,^[4] and detectors for explosives,^[5] because of their optoelectronic properties derived from the π -conjugated system. Several efficient synthetic methods for dibenzosiloles using transition-metal catalysts have therefore been actively developed in recent years.^[6] In contrast, the preparation of enantioenriched chiral dibenzosiloles has been scarcely investigated^[7] despite their potential future applications, for example as materials for circularly polarized luminescence (CPL).^[8] This lack of available methods is presumably due to the planar nature of these compounds except at the silicon atom. However, the introduction of chirality at the silicon center, particularly in a catalytic asymmetric manner, is not a trivial task compared with the construction of carbon stereocenters.^[9–11] In fact, only two intramolecular approaches have been reported for the

transition-metal-catalyzed enantioselective synthesis of dibenzosiloles.^[7a,b] In this context, herein we describe the first intermolecular asymmetric synthesis of silicon-stereogenic dibenzosiloles through a rhodium-catalyzed [2+2+2] cycloaddition of silicon-containing prochiral triynes with internal alkynes.^[12,13]


Initially, we employed prochiral triyne **1a**, which has two identical alkynyl groups on the silicon atom, as a model substrate for the rhodium-catalyzed [2+2+2] cycloaddition with 1,4-dimethoxy-2-butyne (**2a**) in the presence of (*R*)-Binap^[14] as the ligand (Table 1, entry 1). The reaction

Table 1: The effect of the ligand in the rhodium-catalyzed asymmetric [2+2+2] cycloaddition of **1a** with **2a**.



Entry	Ligand	Yield [%] ^[a]	ee ^[b]
1	(<i>R</i>)-Binap	74	18
2	(<i>R</i>)-H ₈ -Binap	83	9
3	(<i>R</i>)-Segphos	95	73
4	(<i>R</i>)-MeO-Mop	63	80
5	(<i>R</i>)-L	98	88

[a] Yield of isolated product. [b] Determined by chiral HPLC on a Chiralpak IF-3 column with hexane/2-propanol = 500:1 v/v.



proceeded at 25°C to give the desired silicon-stereogenic dibenzosilole **3aa** in 74% yield, but the enantiomeric excess was only 18%.^[15] The change of ligand to (*R*)-H₈-Binap^[16] resulted in even lower enantioselectivity (9% ee; entry 2), whereas the use of (*R*)-Segphos^[17] improved the enantioselectivity to 73% ee (Table 1, entry 3). Compared to these axially chiral bisphosphine ligands that are typically used for rhodium-catalyzed asymmetric [2+2+2] cycloaddition reactions,^[12] (*R*)-MeO-mop,^[18] an axially chiral monophosphine

[*] Dr. R. Shintani, C. Takagi, Prof. Dr. K. Nozaki
Department of Chemistry and Biotechnology
Graduate School of Engineering, The University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656 (Japan)
E-mail: shintani@chembio.t.u-tokyo.ac.jp
nozaki@chembio.t.u-tokyo.ac.jp

T. Ito
Department of Chemistry, Graduate School of Science
Kyoto University, Sakyo, Kyoto 606-8502 (Japan)

Dr. M. Naito
National Institute of Materials Science
1-1 Namiki, Tsukuba, Ibaraki 305-0044 (Japan)












[**] Support has been provided in part by a Grant-in-Aid for Young Scientists (B), the Ministry of Education, Culture, Sports, Science, and Technology (Japan) and in part by the Asahi Glass Foundation. We thank Prof. Takuzo Aida at the University of Tokyo for the measurement of fluorescence and CD spectra. We thank Prof. Yoshiaki Nishibayashi and Dr. Kazunari Nakajima at the University of Tokyo for the measurement of optical rotations. We thank Masahiro Miyamoto at Daicel Corporation for the help of the chiral HPLC analysis for compound **3ac**. We thank Dr. Shingo Ito and Dr. Shuhei Kusumoto for X-ray crystallographic analysis.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201409733>.

ligand, gave **3aa** with higher enantioselectivity (80% *ee*; entry 4). By introducing a methyl group at the 3'-position of (*R*)-MeO-mop ((*R*)-**L**),^[19] the enantioselectivity of **3aa** was further improved to 88% *ee* in 98% yield (entry 5).^[20]

Under the conditions described in Table 1 entry 5, various silicon-stereogenic dibenzosiloles can be synthesized with high yields and enantioselectivities as summarized in Table 2.


Table 2: Reaction scope of the rhodium-catalyzed asymmetric synthesis of dibenzosiloles **3**.

			
Entry	Product	Yield [%] ^[a]	<i>ee</i> ^[b]
1		98	88
2		96	91
3		99	86
4		91	89
5 ^[c]		72	85
6		95	85
7		91	59
8		80	87
9 ^[c]		92	88
10 ^[d]		90	88
11 ^[e]		91 ^[f]	88
12		60	96
		26	77

[a] Yield of isolated product. [b] Determined by chiral HPLC. [c] The reaction was conducted at 30 °C. [d] The regioselectivity was 99:1. [e] The regioselectivity was 93:7. [f] Yield of the isolated major regioisomer.



Thus, the 1-pentynyl group on **1a** (entry 1) can be replaced by other alkynyl groups, such as 1-propynyl (**1b**) and 4-methyl-1-pentynyl (**1c**) groups to give **3ba** and **3ca** in high yields with 86–91% *ee* (entries 2 and 3). Triynes having two identically substituted arylethynyl groups on the silicon atom (**1d** and **1e**) are also suitable substrates for the reaction with alkyne **2a** to give **3da** and **3ea** with similar enantiomeric excesses (85–89% *ee*; entries 4 and 5). Replacement of the *tert*-butyl group

on the silicon atom of **1a** by the less bulky cyclohexyl group (**1f**) provides the corresponding dibenzosilole **3fa** with a relatively high *ee* value of 85% (entry 6), but the use of the phenyl-substituted variant (**1g**) results in the formation of dibenzosilole **3ga** with significantly lower enantioselectivity (59% *ee*; entry 7). With regard to the internal alkyne, symmetrical alkynes, such as **2a**, 1,4-diacetoxy-2-butyne (**2b**), and bis(4-methoxyphenyl)acetylene (**2c**), as well as unsymmetrical alkynes, such as dimethyl 2-(2-butynyl)-2-methylmalonate (**2d**), methyl tetrolate (**2e**), and 1-phenyl-1-propyne (**2f**), can be employed for the present catalysis to give **3aa–3ae** and **3df/3d'** in high yields (80–92% yield) with up to 96% *ee* (entries 8–12). It is worth mentioning that the reactions of **1a** with unsymmetrical **2d** and **2e** proceed with high regioselectivity to give **3ad** (99:1; entry 10) and **3ae** (93:7; entry 11) respectively,^[21] although the reaction of **1d** with **2f** gives a 70:30 mixture of regioisomers (entry 12).^[22] More functionalized dibenzosiloles **3hg/3hg'** bearing polymerizable sites through the (sila-)Sonogashira coupling can also be prepared enantioselectively (96% *ee* and 70% *ee*, respectively) as shown in Eq. (1) with moderate regioselectivity (76:24).^[21,23] In addition to triynes **1**, thiophene-tethered triyne **4** can also react with alkyne **2a**, leading to the formation of benzosilolothiophene **5** in 72% yield with 92% *ee* [Eq. (2)].



We have also found that the present asymmetric catalysis can be applied to the preparation of enantioenriched germanium-stereogenic dibenzogermoles by employing germanium-containing prochiral triynes as substrates. For example, the reaction of triyne **6** with alkyne **2a** under our standard conditions gives dibenzogermole **7** in 73% yield with 86% *ee* [Eq. (3)]. To our knowledge, this result demonstrates the first example of the enantioselective construction of a germanium stereogenic center under transition-metal catalysis.^[24] Furthermore, the reaction of prochiral hexayne

8 gives doubly cyclized product **9** in an 81 % combined yield of chiral and *meso* isomers (ratio = 83:17) with 94 % *ee* for the chiral isomer through the twofold enantioselective [2+2+2] cycloaddition with alkyne **2a** [Eq. (4)].



Having established the catalytic asymmetric synthesis of various dibenzosiloles and related compounds, we examined the optical properties of these compounds, considering the lack of literature precedent for the study of the optical properties of silicon-stereogenic dibenzosiloles. In our preliminary investigation, we focused on comparing the properties of **3aa** and chiral **9**. As summarized in Table 3, the UV/Vis

Table 3: Optical properties of compounds **3aa** and **9** in CH_2Cl_2 at 25 °C.

Compound	UV/Vis absorption $\lambda_{\text{max}}/\text{nm}$ ($\epsilon/10^4 \text{ M}^{-1} \text{ cm}^{-1}$)	Emission $\lambda_{\text{max}}/\text{nm}^{[a]}$	$\Phi_{\text{F}}^{[a]}$
(<i>R</i>)- 3aa (88 % <i>ee</i>)	245 (10.1), 330 (0.7) ^[b]	358 ^[c]	0.19 ^[c]
(<i>R,R</i>)- 9 (94 % <i>ee</i>)	256 (9.2), 331 (3.6), 368 (1.9) ^[d]	404 ^[e]	0.54 ^[e]

[a] Excited at $\lambda = 250 \text{ nm}$. [b] At $1.1 \times 10^{-5} \text{ M}$. [c] At $2.3 \times 10^{-6} \text{ M}$. [d] At $5.7 \times 10^{-6} \text{ M}$. [e] At $2.2 \times 10^{-6} \text{ M}$.

absorption and emission band maxima are red-shifted for doubly cyclized (*R,R*)-**9** compared to dibenzosilole (*R*)-**3aa**. Additionally, the absolute quantum yield of (*R,R*)-**9** was significantly higher than that of (*R*)-**3aa** (0.54 vs. 0.19).^[25] We also examined the CD spectra of these compounds and found that (*R*)-**3aa** showed a relatively weak negative Cotton effect at $\lambda = 259 \text{ nm}$ ($\Delta\epsilon = -3.0 \text{ M}^{-1} \text{ cm}^{-1}$), whereas (*R,R*)-**9** showed stronger positive and negative Cotton effects at $\lambda = 282 \text{ nm}$ ($\Delta\epsilon = +7.3 \text{ M}^{-1} \text{ cm}^{-1}$) and $\lambda = 256 \text{ nm}$ ($\Delta\epsilon = -10.8 \text{ M}^{-1} \text{ cm}^{-1}$), respectively (Figure 1). Based on these data, we attempted to measure the circularly polarized luminescence (CPL) of (*R,R*)-**9**, but, unfortunately, the CPL activity was only close to the measurement limit ($g_{\text{lum}} \sim 10^{-4}$) for this particular compound.^[26]

In summary, we have developed a rhodium-catalyzed asymmetric synthesis of silicon-stereogenic dibenzosiloles through a [2+2+2] cycloaddition of silicon-containing prochiral triynes with internal alkynes. High yields and enantio-




Figure 1. CD spectra of (*R*)-**3aa** (gray line; $1.1 \times 10^{-5} \text{ M}$) and (*R,R*)-**9** (black line; $5.7 \times 10^{-6} \text{ M}$) in CH_2Cl_2 at 25 °C.

selectivities have been achieved by employing axially chiral monophosphine (*R*)-**L** as the ligand, and this process could also be applied to the asymmetric synthesis of a germanium-stereogenic dibenzogermole. We have also conducted preliminary investigations on the optical properties of these compounds. Future studies will be directed toward further expansion of the scope of this catalysis including the synthesis of more CPL-active compounds.

Experimental Section

General procedure for Table 2: A solution of $[(\text{RhCl}(\text{C}_2\text{H}_4)_2)_2]$ (1.5 mg, 7.7 μmol Rh) and (*R*)-**L** (3.6 mg, 7.5 μmol) in CH_2Cl_2 (0.50 mL) were stirred for 20 min at 25 °C, and a mixture of triyne **1** (0.150 mmol), alkyne **2** (0.150 mmol), and NaBARF_4 (13.3 mg, 15.0 μmol) in CH_2Cl_2 (0.50 mL) was added to it with additional CH_2Cl_2 (0.50 mL). The reaction mixture was stirred for 13–23 h at 25 °C, and this was directly passed through a pad of silica gel with EtOAc. After removal of the solvent under vacuum, the residue was purified by silica gel preparative TLC to afford compound **3**.

Received: October 3, 2014

Revised: November 14, 2014

Published online: December 9, 2014

Keywords: asymmetric catalysis · cycloaddition · dibenzosilole · rhodium · silicon

- [1] For reviews, see: a) J. Chen, Y. Cao, *Macromol. Rapid Commun.* **2007**, 28, 1714; b) W. W. H. Wong, J. F. Hooper, A. B. Holmes, *Aust. J. Chem.* **2009**, 62, 393; c) H. Fu, Y. Cheng, *Curr. Org. Chem.* **2012**, 16, 1423.
- [2] For recent examples, see: a) K. L. Chan, M. J. McKiernan, C. R. Towns, A. B. Holmes, *J. Am. Chem. Soc.* **2005**, 127, 7662; b) S. H. Lee, B.-B. Jang, Z. H. Kafafi, *J. Am. Chem. Soc.* **2005**, 127, 9071; c) E. Wang, C. Li, W. Zhuang, J. Peng, Y. Cao, *J. Mater. Chem.* **2008**, 18, 797; d) M. Shimizu, H. Tatsumi, K. Mochida, K. Oda, T. Hiyama, *Chem. Asian J.* **2008**, 3, 1238; e) Y.-Q. Mo, X.-Y. Deng, X. Jiang, Q.-H. Cui, *J. Polym. Sci. Part A* **2009**, 47, 3286; f) L. Li, C. Xu, S. Li, *Tetrahedron Lett.* **2010**, 51, 622; g) C. W. Keyworth, K. L. Chan, J. G. Labram, T. D. Anthopoulos, S. E. Watkins, M. MaKiernan, A. J. P. White, A. B. Holmes, C. K. Williams, *J. Mater. Chem.* **2011**, 21, 11800.
- [3] a) H. Usta, G. Lu, A. Facchetti, T. J. Marks, *J. Am. Chem. Soc.* **2006**, 128, 9034; b) G. Lu, H. Usta, C. Risko, L. Wang, A. Facchetti, M. A. Ratner, T. J. Marks, *J. Am. Chem. Soc.* **2008**, 130, 7670.

- [4] For recent examples, see: a) J. Hou, H.-Y. Chen, S. Zhang, G. Li, Y. Yang, *J. Am. Chem. Soc.* **2008**, *130*, 16144; b) L. Huo, H.-Y. Chen, J. Hou, T. L. Chen, Y. Yang, *Chem. Commun.* **2009**, 5570; c) C. Duan, W. Cai, F. Huang, J. Zhang, M. Wang, T. Yang, C. Zhong, X. Gong, Y. Cao, *Macromolecules* **2010**, *43*, 5262; d) M. Akhtaruzzaman, Y. Seya, N. Asao, A. Islam, E. Kwon, A. El-Shafei, L. Han, Y. Yamamoto, *J. Mater. Chem.* **2012**, *22*, 10771; e) C. Bathula, C. E. Song, W.-H. Lee, J. Lee, S. Badgujar, R. Koti, I.-N. Kang, W. S. Shin, T. Ahn, J.-C. Lee, S.-J. Moon, S. K. Lee, *Thin Solid Films* **2013**, *537*, 231; f) Y. Lin, Y. Li, X. Zhan, *Adv. Energy Mater.* **2013**, *3*, 724; g) H. J. Song, J. Y. Lee, E. J. Lee, D. K. Moon, *Eur. Polym. J.* **2013**, *49*, 3261.
- [5] a) J. C. Sanchez, A. G. DiPasquale, A. L. Rheingold, W. C. Trogler, *Chem. Mater.* **2007**, *19*, 6459; b) J. C. Sanchez, S. A. Urbas, S. J. Toal, A. G. DiPasquale, A. L. Rheingold, W. C. Trogler, *Macromolecules* **2008**, *41*, 1237; c) J. Yang, S. Aschmeyer, H. P. Martinez, W. C. Trogler, *Chem. Commun.* **2010**, 6804.
- [6] For reviews, see: a) J. Y. Corey, *Adv. Organomet. Chem.* **2011**, *59*, 181; b) T. Jin, J. Zhao, N. Asao, Y. Yamamoto, *Chem. Eur. J.* **2014**, *20*, 3554. For leading references, see: c) T. Matsuda, S. Kadowaki, T. Goya, M. Murakami, *Org. Lett.* **2007**, *9*, 133; d) M. Shimizu, K. Mochida, T. Hiyama, *Angew. Chem. Int. Ed.* **2008**, *47*, 9760; *Angew. Chem.* **2008**, *120*, 9906; e) K. Mochida, M. Shimizu, T. Hiyama, *J. Am. Chem. Soc.* **2009**, *131*, 8350; f) M. Tobisu, M. Onoe, Y. Kita, N. Chatani, *J. Am. Chem. Soc.* **2009**, *131*, 7506; g) Y. Yabusaki, N. Ohshima, H. Kondo, T. Kusamoto, Y. Yamanoi, H. Nishihara, *Chem. Eur. J.* **2010**, *16*, 5581; h) T. Ureshino, T. Yoshida, Y. Kuninobu, K. Takai, *J. Am. Chem. Soc.* **2010**, *132*, 14324; i) Y. Liang, S. Zhang, Z. Xi, *J. Am. Chem. Soc.* **2011**, *133*, 9204.
- [7] a) R. Shintani, H. Otomo, K. Ota, T. Hayashi, *J. Am. Chem. Soc.* **2012**, *134*, 7305; b) Y. Kuninobu, K. Yamauchi, N. Tamura, T. Seiki, K. Takai, *Angew. Chem. Int. Ed.* **2013**, *52*, 1520; *Angew. Chem.* **2013**, *125*, 1560. See also: c) S. Yasuike, T. Iida, S. Okajima, K. Yamaguchi, H. Seki, J. Kurita, *Tetrahedron* **2001**, *57*, 10047; d) T. Shibata, T. Uchiyama, Y. Yoshinami, S. Takayasu, K. Tsuchikama, K. Endo, *Chem. Commun.* **2012**, *48*, 1311; e) H. Oyama, K. Nakano, T. Harada, R. Kuroda, M. Naito, K. Nobusawa, K. Nozaki, *Org. Lett.* **2013**, *15*, 2104.
- [8] For reviews, see: a) F. S. Richardson, *Chem. Rev.* **1977**, *77*, 773; b) J. P. Riehl, F. S. Richardson, *Chem. Rev.* **1986**, *86*, 1.
- [9] For reviews on asymmetric synthesis of silicon-stereogenic organosilanes, see: a) M. Oestreich, *Synlett* **2007**, 1629; b) L.-W. Xu, L. Li, G.-Q. Lai, J.-X. Jiang, *Chem. Soc. Rev.* **2011**, *40*, 1777; See also: c) L.-W. Xu, *Angew. Chem. Int. Ed.* **2012**, *51*, 12932; *Angew. Chem.* **2012**, *124*, 13106.
- [10] For examples of catalytic asymmetric preparation of silicon-stereogenic organosilanes, see: a) R. J. P. Corriu, J. J. E. Moreau, *Tetrahedron Lett.* **1973**, *14*, 4469; b) T. Hayashi, K. Yamamoto, M. Kumada, *Tetrahedron Lett.* **1974**, *15*, 331; c) R. J. P. Corriu, J. J. E. Moreau, *J. Organomet. Chem.* **1975**, *85*, 19; d) R. J. P. Corriu, J. J. E. Moreau, *J. Organomet. Chem.* **1976**, *120*, 337; e) T. Ohta, M. Ito, A. Tsuneto, H. Takaya, *J. Chem. Soc. Chem. Commun.* **1994**, 2525; f) Y. Yasutomi, H. Suematsu, T. Katsuki, *J. Am. Chem. Soc.* **2010**, *132*, 4510; g) R. Shintani, K. Moriya, T. Hayashi, *J. Am. Chem. Soc.* **2011**, *133*, 16440; h) R. Shintani, K. Moriya, T. Hayashi, *Org. Lett.* **2012**, *14*, 2902; i) R. Shintani, E. E. Maciver, F. Tamakuni, T. Hayashi, *J. Am. Chem. Soc.* **2012**, *134*, 16955; j) Y. Kurihara, M. Nishikawa, Y. Yamanoi, H. Nishihara, *Chem. Commun.* **2012**, *48*, 11564; k) K. Igawa, D. Yoshihiro, N. Ichikawa, N. Kokan, K. Tomooka, *Angew. Chem. Int. Ed.* **2012**, *51*, 12745; *Angew. Chem.* **2012**, *124*, 12917. See also: l) K. Tamao, K. Nakamura, H. Ishii, S. Yamaguchi, M. Shiro, *J. Am. Chem. Soc.* **1996**, *118*, 12469; m) D. R. Schmidt, S. J. O'Malley, J. L. Leighton, *J. Am. Chem. Soc.* **2003**, *125*, 1190; n) K. Igawa, J. Takada, T. Shimono, K. Tomooka, *J. Am. Chem. Soc.* **2008**, *130*, 16132; o) M. Onoe, K. Baba, Y. Kim, Y. Kita, M. Tobisu, N. Chatani, *J. Am. Chem. Soc.* **2012**, *134*, 19477; p) X. Lu, L. Li, W. Yang, K. Jiang, K.-F. Yang, Z.-J. Zheng, L.-W. Xu, *Eur. J. Org. Chem.* **2013**, 5814.
- [11] Stoichiometric construction of chiral silicon stereocenters has been somewhat more investigated. For pioneering work, see: a) L. H. Sommer, *Stereochemistry, Mechanism and Silicon; an Introduction to the Dynamic Stereochemistry and Reaction Mechanisms of Silicon Centers*, McGraw-Hill, New York, **1965**. For selected recent examples, see: b) C. Strohmann, J. Hörnig, D. Auer, *Chem. Commun.* **2002**, 766; c) M. Trzoss, J. Shao, S. Bienz, *Tetrahedron: Asymmetry* **2004**, *15*, 1501; d) S. Rendler, G. Auer, M. Oestreich, *Angew. Chem. Int. Ed.* **2005**, *44*, 7620; *Angew. Chem.* **2005**, *117*, 7793; e) S. Rendler, G. Auer, M. Keller, M. Oestreich, *Adv. Synth. Catal.* **2006**, *348*, 1171; f) S. Rendler, M. Oestreich, C. P. Butts, G. C. Lloyd-Jones, *J. Am. Chem. Soc.* **2007**, *129*, 502; g) K. Igawa, N. Kokan, K. Tomooka, *Angew. Chem. Int. Ed.* **2010**, *49*, 728; *Angew. Chem.* **2010**, *122*, 740.
- [12] For recent reviews on the rhodium-catalyzed [2+2+2] cycloaddition reactions, see: a) K. Tanaka, *Chem. Asian J.* **2009**, *4*, 508; b) N. Weding, M. Hapke, *Chem. Soc. Rev.* **2011**, *40*, 4525; c) Y. Shibata, K. Tanaka, *Synthesis* **2012**, *44*, 323; d) D. L. J. Broere, E. Ruijter, *Synthesis* **2012**, *44*, 2639. For selected recent enantioselective examples, see: e) T. Shibata, T. Uchiyama, K. Endo, *Org. Lett.* **2009**, *11*, 3906; f) F. Mori, N. Fukawa, K. Noguchi, K. Tanaka, *Org. Lett.* **2011**, *13*, 362; g) N. Sakiyama, D. Hojo, K. Noguchi, K. Tanaka, *Chem. Eur. J.* **2011**, *17*, 1428; h) S. Ogaki, Y. Shibata, K. Noguchi, K. Tanaka, *J. Org. Chem.* **2011**, *76*, 1926; i) L. Garcia, A. Roglans, R. Laurent, J.-P. Majoral, A. Pla-Quintana, A.-M. Caminade, *Chem. Commun.* **2012**, *48*, 9248; j) Y. Kimura, N. Fukawa, Y. Miyauchi, K. Noguchi, K. Tanaka, *Angew. Chem. Int. Ed.* **2014**, *53*, 8480; *Angew. Chem.* **2014**, *126*, 8620.
- [13] For a pioneering work on the metal-catalyzed enantioselective desymmetrization through [2+2+2] cycloaddition reactions, see: Y. Sato, T. Nishimata, M. Mori, *J. Org. Chem.* **1994**, *59*, 6133.
- [14] H. Takaya, K. Mashima, K. Koyano, M. Yagi, H. Kumobayashi, T. Taketomi, S. Akutagawa, R. Noyori, *J. Org. Chem.* **1986**, *51*, 629.
- [15] A nonasymmetric variant was first reported by Murakami and co-workers using iridium catalysis at elevated temperatures. See Ref. [6c]. The use of $[\text{IrCl}(\text{C}_2\text{H}_4)_2]_2$ in place of $[\text{RhCl}(\text{C}_2\text{H}_4)_2]_2$ for the present catalysis resulted in no reaction.
- [16] X. Zhang, K. Mashima, K. Koyano, N. Sayo, H. Kumobayashi, S. Akutagawa, H. Takaya, *Tetrahedron Lett.* **1991**, *32*, 7283.
- [17] T. Saito, T. Yokozawa, T. Ishizaki, T. Moroi, N. Sayo, T. Miura, H. Kumobayashi, *Adv. Synth. Catal.* **2001**, *343*, 264.
- [18] a) T. Hayashi, *Acc. Chem. Res.* **2000**, *33*, 354; b) Y. Uozumi, A. Tanahashi, S.-Y. Lee, T. Hayashi, *J. Org. Chem.* **1993**, *58*, 1945.
- [19] B. Saha, T. V. RajanBabu, *J. Org. Chem.* **2007**, *72*, 2357.
- [20] The reaction using 5 mol % of NaBARF_4 instead of 10 mol % gave **3aa** in a lower yield of 38 % (47 % conversion of **1a**) with the same *ee* value of 88 %.
- [21] The structures of **3ad**, **3ae**, and **3hg/3hg'** were determined by NOE NMR spectroscopy measurements. See the Supporting Information for details.
- [22] The structure of **3df** including the absolute configuration was determined by X-ray crystallographic analysis. CCDC-1027469 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [23] We are now actively investigating the synthesis and properties of the macromolecules based on this and related strategies. The results will be compiled and reported in due course.

- [24] An enzymatic approach for the construction of germanium seterogenic centers has been reported, see: a) R. Tacke, U. Kosub, S. A. Wagner, R. Bertermann, S. Schwarz, S. Merget, K. Günter, *Organometallics* **1998**, *17*, 1687. For examples of the optical resolution of racemic compounds, see: b) R. Tacke, D. Reichel, P. G. Jones, X. Hou, M. Waelbroeck, J. Gross, E. Mutschler, G. Lambrecht, *J. Organomet. Chem.* **1996**, *521*, 305; c) R. Tacke, T. Kornek, T. Heinrich, C. Burschka, M. Penka, M. Pülm, C. Keim, E. Mutschler, G. Lambrecht, *J. Organomet. Chem.* **2001**, *640*, 140.
- [25] For a similar observation, see Ref. [6c].
- [26] For recent examples of CPL-active small organic molecules, see: a) J. E. Field, G. Muller, J. P. Riehl, D. Venkataraman, *J. Am. Chem. Soc.* **2003**, *125*, 11808; b) R. A. van Delden, N. P. M. Huck, J. J. Piet, J. M. Warman, S. C. J. Meskers, H. P. J. M. Dekkers, B. L. Feringa, *J. Am. Chem. Soc.* **2003**, *125*, 15659; c) T. Kawai, K. Kawamura, H. Tsumatori, M. Ishikawa, M. Naito, M. Fujiki, T. Nakashima, *ChemPhysChem* **2007**, *8*, 1465; d) F. C. Spano, S. C. J. Meskers, E. Hennebicq, D. Beljonne, *J. Am. Chem. Soc.* **2007**, *129*, 7044; e) H. Tsumatori, T. Nakashima, T. Kawai, *Org. Lett.* **2010**, *12*, 2362; f) T. Kaseyama, S. Furumi, X. Zhang, K. Tanaka, M. Takeuchi, *Angew. Chem. Int. Ed.* **2011**, *50*, 3684; *Angew. Chem.* **2011**, *123*, 3768; g) H. Maeda, Y. Bando, K. Shimomura, I. Yamada, M. Naito, K. Nobusawa, H. Tsumatori, T. Kawai, *J. Am. Chem. Soc.* **2011**, *133*, 9266; h) Y. Sawada, S. Furumi, A. Takai, M. Takeuchi, K. Noguchi, K. Tanaka, *J. Am. Chem. Soc.* **2012**, *134*, 4080; i) Y. Haketa, Y. Bando, K. Takaishi, M. Uchiyama, A. Muranaka, M. Naito, H. Shibaguchi, T. Kawai, H. Maeda, *Angew. Chem. Int. Ed.* **2012**, *51*, 7967; *Angew. Chem.* **2012**, *124*, 8091; j) K. Goto, R. Yamaguchi, S. Hiroto, H. Ueno, T. Kawai, H. Shinokubo, *Angew. Chem. Int. Ed.* **2012**, *51*, 10333; *Angew. Chem.* **2012**, *124*, 10479; k) See Ref. [7e]; l) J. Kumar, T. Nakashima, H. Tsumatori, M. Mori, M. Naito, T. Kawai, *Chem. Eur. J.* **2013**, *19*, 14090; m) H. Maeda, W. Hane, Y. Bando, Y. Terashima, Y. Haketa, H. Shibaguchi, T. Kawai, M. Naito, K. Takaishi, M. Uchiyama, A. Muranaka, *Chem. Eur. J.* **2013**, *19*, 16263; n) Y. Morisaki, M. Gon, T. Sasamori, N. Tokitoh, Y. Chujo, *J. Am. Chem. Soc.* **2014**, *136*, 3350; o) K. Nakamura, S. Furumi, M. Takeuchi, T. Shibuya, K. Tanaka, *J. Am. Chem. Soc.* **2014**, *136*, 5555.