

Nitrosopurines En Route to Potently Cytotoxic Asmarines**

Kanny K. Wan, Kotaro Iwasaki, Jeffrey C. Umotoy, Dennis W. Wolan,* and Ryan A. Shenvi*

Abstract: A nitrosopurine ene reaction easily assembles the asmarine pharmacophore and transmits remote stereochemistry to the diazepine-purine heterocycle. This reaction generates potent cytotoxins which exceed the potency of asmarine A ($1.2 \mu\text{M}$ IC_{50}) and supersede the metabolites as useful leads for biological discovery.

Asmarines A and B (**1** and **2**; Figure 1) were identified in 1998 by Kashman and co-workers as the bioactive constituents of a Red Sea sponge (*Raspailia* sp.) extract, and exhibited cytotoxicity against several cancer cell lines with a minimum EC_{50} value of $1.2 \mu\text{M}$ and 120 nM , respectively.^[1] The asmarines are unique among alkaloids by virtue of the embedded *N*-hydroxypurine diazepine (primary pharmacophore)^[2] connected by an ethyl bridge to a clerodane decalin (putative secondary pharmacophore). Biosynthetically, the asmarines derive from agelasines (e.g. **3**), which are thought to exert cytotoxicity by Na^+/K^+ ATPase inhibition^[3] or by membrane disruption.^[4] However, the uncharged purines **1** and **2** are likely to exert cytotoxicity through different mechanisms than

3. Thus, the driving force for investigation of the asmarines is to determine their mechanism of action and chemical reactivity within the cell. However, no material remains from the original isolation work^[5] and the exact identity of the sponge is unknown.^[1b] Chemical synthesis is therefore the most feasible way to access material for biological study.

Procurement of these cytotoxic molecules represents a challenge for synthesis despite efforts by Kashman, Schauss, Tashiro, and Gundersen.^[6] Their difficulty arises partly from the remoteness of the stereogenic *tert*-alkyl *N*-hydroxyamine moiety (Figure 1), which frustrates stereocontrol relative to the clerodane subunit. Furthermore, synthesis of the *tert*-alkyl *N*-hydroxy-diazepinepurine pharmacophore is a challenge in itself^[6a,b,d] and only one route exists. This successful strategy by Kashman^[2] utilizes a carbocationic mode of ring closure (30% HBr/AcOH at 100°C), which limits functional-group compatibility and stereocontrol, and delivers analogues (e.g. **4–7**) which, while bioactive, reach a maximum potency of $4 \mu\text{M}$ ^[7] (GI_{50} , HT-29; Figure 2a). We aimed to secure efficient access to the asmarines to understand their structure–activity relationship (SAR) and to provide simplified, high potency analogues for further biological study.

A biomimetic strategy to close the seven-membered ring from an agelasine-type intermediate **8** (Figure 2b) appeared to offer high efficiency. However, control of the diazepine stereochemistry during C–N bond formation presented





Figure 1. Asmarines may be derived from agelasines. [a] Against HT-29 cells. [b] Against MCF-7 cells.^[3b]

a) Kashman analogues



b) This work




Figure 2. a) Prior analogue work. b) Stereochemical relay and high potency analogues through nitrosopurine ene reaction. [a] Cytotoxicity against HT-29 cells (EC_{50} more appropriate).

[*] K. K. Wan, Dr. K. Iwasaki, Prof. Dr. R. A. Shenvi
Department of Chemistry, The Scripps Research Institute
10550 N. Torrey Pines Rd., La Jolla, CA 92037 (USA)
E-mail: rshenvi@scripps.edu

J. C. Umotoy, Prof. Dr. D. W. Wolan
Department of Molecular and Experimental Medicine and
Chemical Physiology, The Scripps Research Institute
10550 N. Torrey Pines Rd., La Jolla, CA 92037 (USA)
E-mail: wolan@scripps.edu

[**] This work was supported by the NIH (GM104180 and the NSF GRFP (K.K.W.; DGE-1346837). We thank Dr. C. E. Moore and Prof. A. L. Rheingold (UCSD) for X-ray crystallographic analysis. We thank TSRI, Eli Lilly, Boehringer Ingelheim, Amgen, the Baxter Foundation, and the Sloan Foundation for additional financial support.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201411493>.

a serious obstacle to this strategy. We wondered if a nitroso ene reaction^[8] of **8** might relay stereochemistry^[9] of the decalin core to the *tert*-alkylamine stereocenter in **9**, and thus achieve the necessary Markovnikov selectivity.^[10] Herein we report 1) the successful implementation of this strategy to relay clerodane stereochemistry to the remote stereocenter, 2) the ability of this strategy to probe the role of stereochemistry in SARs, and 3) the use of this nitrosopurine ene reaction to synthesize simplified high potency asmarine analogues which exhibit cytotoxicity at nanomolar levels against multiple cancer cell lines.

A short route to the targeted **8** began with 6-chloro-4,5-pyrimidinediamine (**11**) and 4-iodo-1-butyne (**12**; Scheme 1). The diamine **11** was monoformylated to amide **13**, and **12** was subjected to the Wipf-modified^[11] Negishi carboalumination,^[12] with subsequent alkylation of the intermediate organoaluminum with gaseous formaldehyde^[13] to yield the iodoalcohol **14**. The formamide **13** was alkylated with **14** and heating effected ring closure to the chloropurine **15**.^[14] This alcohol was converted into the allylbromide **16** (see the Supporting Information), which was then trapped with the enolate generated from dissolving metal reduction of the (methyl)-Wieland–Miescher ketone^[15] ketal **17** (99% *ee*) in the presence of bis(2-methoxyethyl)amine^[16] to yield **18** as a single stereoisomer. Selective displacement of the arylchloride of **18** with hydroxylamine proved challenging since condensation with the decalin ketone to form an oxime might be cleaved later, and thus decided to push the reaction towards the bis(hydroxyamin)ated product **19**, an unforeseen but providential choice (see Table 1).

Having obtained the agelasine scaffold **19**, oxidation of the 6-*N*-hydroxyaminopurine (HAP) moiety to a nitrosopur-

ine was explored in a model system (Figure 3). The parent compound 6-nitrosopurine **20**^[17] proved unstable and in our hands could not be isolated. However, we found that treatment of the 6-*N*-hydroxyaminopurine **21** with MnO₂ in DMSO and in the presence of excess tetramethylethylenegave the expected *tert*-alkyl methallyl hydroxylamine **22**. A variety of oxidants [I₂, Mn(OAc)₃, PhI(OAc)₂] gave similar results and on large scale PhI(OAc)₂ proved more amenable to purification. When these oxidation conditions were applied to the *N*-hydroxyaminopurine **23** in a mixture of methanol and dichloromethane, the targeted seven-membered diazepine purine **24** was obtained as the sole product with no trace of the eight-membered diazocine. Notably, this reaction could be performed on gram-scale, thus demonstrating that the





Figure 3. Proof-of-principle for nitrosopurine ene reactions. DMSO = dimethylsulfoxide.



Scheme 1. Union of clerodane and purine cores, and a nitrosopurine ene reaction with remote stereocontrol. Cp = cyclopentadiene, Ms = methanesulfonyl, THF = tetrahydrofuran.

chemistry of nitrosopurines (e.g. **25**) is a practical means of procuring material.

When applied to the agelasine-type compound **19**, we also obtained the diazepine **26** exclusively, but initial experiments generated this as a 1:1 mixture of diastereomers (entry 1, Table 1). Fortunately, we discovered two influences on diastereoselectivity in the ene reaction. There was a marked solvent effect associated with diastereoselectivity, whereby more polar protic solvents increased selectivity, and water proved to be the most selective (ethanol was added for solubility). This effect was not due to protonation or hydrogen bonding, since neither acetic acid had a pronounced effect (entry 6), nor

did the strongly hydrogen-bond-donating solvent trifluoroethanol^[18] (entry 7) when compared to ethanol (entry 3). There was also a stark and serendipitous substituent effect: although the incidental oxime caused high stereoselectivity, the ketone **27** and exocyclic methylene **28** showed almost no selectivity in formation of **29** and **30**, respectively (entries 10 and 11). Application of the polarized diradical (PD) model for the nitroso ene reaction^[19] of **19** suggests that developing positive charge at the carbon atom in the transition state might be stabilized by the proximal oxime (Figure 4). The conformer **32** should be lower in energy than **33**, which suffers a steric clash between the methyl and the aromatic ring. These polarized transition states would be more populated in solvents of high polarity (e.g., water), as predicted by Leach and Houk.^[19]

This strategy allowed us to directly probe the contribution of diazepine stereochemistry to cytotoxicity. The major diastereomer **26** (Figure 5) exhibited higher potency against




Figure 4. Water might increase the polar character in a nitroso ene reaction and the transition state leading to it. Conformers of different energy are stabilized by the proximal oxime. X = (-OC₂H₄O-).

Table 1: Solvent and substituent control stereoselection.

19:	X = (-OC ₂ H ₄ O-), Y = NOH	26:	X = (-OC ₂ H ₄ O-), Y = NOH
27:	X = O, Y = O	29:	X = O, Y = O
28:	X = (-OC ₂ H ₄ O-), Y = CH ₂	30:	X = (-OC ₂ H ₄ O-), Y = CH ₂

Entry	Y	Solvent	T [°C]	Additive	(S)-13/(R)-13 ^[a]
1	NOH	DMF	22	–	1.5:1.0
2	NOH	CH ₂ Cl ₂	22	–	1.5:1.0
3	NOH	EtOH	22	–	3.8:1.0
4	NOH	EtOH	–20	–	3.0:1.0
5	NOH	MeOH/CH ₂ Cl ₂ (1:9)	22	NH ₄ OH ^[b]	2.1:1.0
6	NOH	EtOH	22	AcOH ^[c]	2.7:1.0
7	NOH	TFE	22	–	2.3:1.0
8	NOH	EtOH/H ₂ O (1:1)	22	–	4.7:1.0
9	NOH	EtOH/H ₂ O (1:3)	22	–	7.9:1.0
10	O	EtOH/H ₂ O (1:4)	22	–	1.2:1.0
11	CH ₂	EtOH/H ₂ O (1:4)	22	–	1.1:1.0

[a] Determined by NMR spectroscopy and LCMS. [b] 1% NH₄OH.

[c] 1 equiv AcOH. DMF = N,N-dimethylformamide.

HT-29 cells (EC₅₀ = 8 μM) than the minor isomer **34** (17 μM), even though **34** corresponds to the stereochemistry of the natural asmarines (determined by X-ray analysis;^[20] see the Supporting Information). Since the difference in potency is not profound, the stereochemistry in the diazepine seems to




Figure 5. Activity (EC₅₀, 48 h) against HT-29 cells. Although there is clear SAR, stereochemistry does not play a major role. X = (-OC₂H₄O-).

play only a minor role in the mechanism of action. Synthesis also allowed us to probe the role of absolute stereochemistry. We were surprised to find that *ent*-**26** and *ent*-**34** possessed similar biological activity as their enantiomers ($EC_{50} = 10 \mu\text{M}$ for both). Similarly, the diketones **35** and *ent*-**35** possess activity which differs only twofold. Whereas this similarity might suggest a nonspecific mechanism of action, it seems more likely that the differences in potencies reflect cell permeability, thus affecting accessibility to the target. As such, replacement of the oxime with a more lipophilic methylene delivers compounds with greater potency (**36** and **37**). In this case, the ‘natural’ diazepine **37** is fivefold more potent than the ‘unnatural’ diastereomer **36**, and in fact matches the potency against HT-29 cells reported for asmarine A ($EC_{50} = 1 \mu\text{M}$). As control compounds, we also tested the N-H purine **38** and the uncyclized *N*-hydroxyaminopurine **19**. Both compounds were unable to effectively induce cell death after 72 hours. The inactivity of **19** and **37** suggests a different mechanism than that of the inhibition of ATPases ascribed to the agelasines and their simplified analogues,^[21] although further study will be necessary to rule out this target.

Since the ultimate goal of our work was procurement of material for biological study and it was clear that stereo-

chemistry had little influence on the activity of the asmarines, we adapted the route shown in Scheme 1 to the late-stage, divergent appendage of unnatural hydrophobic cores. As shown in Figure 6, a short convergent sequence from **13** and **39** was devised to arrive at the iodide **40** on gram scale. From **39**, a variety of substituents were added by a very effective Negishi coupling,^[22] a noteworthy step given the acidic proton on the hydroxyamino group and our observation that the *N*-hydroxyamino purine will chelate metals. Each adduct (**41–48**) was then cyclized by the nitrosopurine ene reaction to its *N*-hydroxydiazepine purine (**50–57**) to generate a small library of cytotoxic compounds.

Each molecule in Figure 6 was screened for cytotoxicity against HT-29, Jurkat, and HeLa cells. While compounds with truncated side chains (**24**, **49–53**) show cytotoxicity, the potency is weak, with or without an unsaturated linker (**24** versus **49**). Small rings like a methylenecyclobutane can be generated in the ene reaction, thus highlighting the mildness of the reaction conditions, which are in stark contrast to the previously reported high temperature/strong acid method of ring closure. Therefore, esters (**52**) and amino-acid (**53**) motifs can be incorporated; the β,γ -unsaturated amino ester carbamate does not migrate into conjugation under the reaction conditions. This latter example shows some small stereose-




Figure 6. A late-stage divergent route to unnatural asmarines and their activities (EC_{50} , 48 h) against HT-29 and HeLa cells. Potencies are on a logarithmic scale.

lectivity (2:1) associated with the presence of the carbamate (compare to methyl; **52**), thus supporting the idea that a Lewis-basic group is necessary for stereoinduction. However, none of these short, polar side chains show significant potency. In contrast, large hydrophobic groups similar to the clerodane cores of asmarines A and B induce high potency. For example, compounds with 1- and 2-naphthyl substituents (**54** and **55**) possess single-digit micromolar activity; **54** is equipotent to asmarine A (1.1 μM against HT-29). 1-Adamantyl (**56**) and 4-biphenyl (**57**) asmarines are more potent with sub-micromolar activity against all three cell lines (471–714 nM; **56** also inhibits HL60 cells at 199 nM, see Figure 8), thus approaching the 120 nM IC_{50} value reported for asmarine B (**2**) against HT29 cells. In fact, this study represents a rare example of the deprioritization of isolated metabolites in light of near-equipotent but simpler analogues. Readily accessible asmarines **56** and **57** supersede the scarce metabolites **1** and **2** as useful tools for biochemical inquiry.

Whereas there is some latitude in the choice of the hydrophobic lobe, the *N*-hydroxy diazepine purine is more conservative. The acyclic *tert*-alkyl *N*-hydroxyaminopurine **19** was completely inactive (Figure 7), thus indicating that a ring




Figure 7. Structural specificity for activity.

is necessary. However, the ring-expanded *N*-hydroxy diazocine purine **51** was similarly inactive. This eight-membered ring was the unexpected anti-Markovnikov product of nitropurine ene reaction of cyclopropane **42**, which reacts with “twix” selectivity and avoids the alternative, highly strained methylenecyclopropane (Figure 7).

The potency of **56** and **57** is general, thus showing sub-micromolar cytotoxicity against HL60 (leukemia), HEK 293, MCF7 (breast cancer), and MDA-MB 231 (breast cancer) cell lines, in addition to the HT29, Jurkat, and HeLa cell lines (Figure 8). The activity of **56** is surprising, since the saturated analogue was reported by Kashman to possess very weak activity against two cancer cell lines (NSCL A549 and PANC1, $\text{EC}_{50} > 27 \mu\text{M}$). The activity of **57** is noteworthy since installation of functional groups on the aromatic rings should allow a variety of pull-down experiments.

This work builds a platform for the discovery of the mechanism of action of the asmarines. We are entertaining three hypotheses: noncovalent, covalent, and radical. The latter two possibilities are supported by some experimental data. Kashman and co-workers^[1b] and Ohba and Tashiro^[6b] have reported the instability of the *N*-hydroxypurine upon acylation, wherein methanol adds to the purine ring and cleaves the N–O bond (**2**→**58**; Figure 9). This mode of reactivity may also be effected in vivo by acetylation or




Figure 8. 1-Adamantyl-asmarine (**56**) and 4-biphenyl-asmarine (**57**) possess nanomolar activity against seven cell lines.




Figure 9. Reactivity of potential relevance to bioactivity.

phosphorylation. Alternatively, we observed that the *tert*-alkyl *N*-hydroxypurine is readily oxidized with mild reagents. The acyclic purine **22** generates the stable nitroxide radical **59**, a vibrant red-orange crystalline solid whose structure was confirmed by X-ray analysis.^[23] Interestingly, the nitroxides of the diazepines (e.g. **24**) are very unstable, cannot be isolated, and instead appear to disproportionate to the corresponding N–H diazepines, thus suggesting that, if generated, they may react with a target in vivo.

To summarize, we have established a chemical platform for the study of asmarine cytotoxins enabled by an unusual but highly practical nitrosopurine ene reaction. This reaction exhibits both high regioselectivity for the Markovnikov product and exhibits high chemoselectivity. Identification of a Lewis base (oxime) as a stereoelectronic control element in the nitroso ene reaction may be generally useful for linear stereocontrol. Use of the nitrosopurine ene reaction as a simplifying element in the synthesis of cytotoxic asmarine analogues enables very short syntheses with few redox manipulations and no protecting groups. Notably, this route is diversifiable at a late stage, and has generated potent analogues with cytotoxicity in the nanomolar range. We have also found that 1) increased potency may result from cell

permeability, but the *N*-hydroxypurine diazepine is required; 2) the similar potency of **54** and **55** suggests the hydrophobic moiety does not play a major role in target binding; 3) stereochemistry at the diazepine plays a minor, but not insignificant role in potency; 4) the N–O bond is required for activity, since an N–H analogue is inactive; and 5) the seven-membered ring is required for activity, since acyclic variants and a diazocine analogue show no cytotoxicity. These findings, coupled with the development of a short, scalable, and divergent route to asmarine analogues, and especially the identification of potent the biphenyl asmarine **57** lay the groundwork for identification of the mechanism of action of these enigmatic metabolites.

Received: November 27, 2014

Published online: January 7, 2015

Keywords: alkaloids · ene reaction · natural products · structure–activity relationships · terpenoids

- [1] a) T. Yosief, A. Rudi, Z. Stein, I. Goldberg, G. M. D. Gravalos, M. Schleyer, Y. Kashman, *Tetrahedron Lett.* **1998**, 39, 3323–3326; b) T. Yosief, A. Rudi, Y. Kashman, *J. Nat. Prod.* **2000**, 63, 299–304.
- [2] D. Pappo, S. Shimony, Y. Kashman, *J. Org. Chem.* **2005**, 70, 199–206.
- [3] a) H. Nakamura, H. Wu, Y. Ohizumi, Y. Hirata, *Tetrahedron Lett.* **1984**, 25, 2989; b) A. A. Pimentel, P. Felibertt, F. Sojo, L. Colman, A. Mayora, M. L. Silva, H. Rojas, R. Dipolo, A. I. Suarez, R. S. Compagnone, F. Arvelo, I. Galindo-Castro, J. B. De Sanctis, P. Chirino, G. Benaim, *Cancer Chemother. Pharmacol.* **2012**, 69, 71; c) R. Fathi-Afshar, T. M. Allen, C. A. Krueger, D. A. Cook, A. S. Clanachan, R. Vriend, H. P. Baer, C. E. Cass, *Can. J. Physiol. Pharmacol.* **1989**, 67, 276.
- [4] A. Vik, E. Hedner, C. Charnock, L. W. Tangen, Ø Samuelsen, R. Larsson, L. Bohlin, L.-L. Gundersen, *Bioorg. Med. Chem.* **2007**, 15, 4016; Samuelsen, R. Larsson, L. Bohlin, L.-L. Gundersen, *Bioorg. Med. Chem.* **2007**, 15, 4016.
- [5] Y. Kashman, personal communication.
- [6] a) D. Pappo, A. Rudi, Y. Kashman, *Tetrahedron Lett.* **2001**, 42, 5941; b) M. Ohba, T. Tashiro, *Heterocycles* **2002**, 57, 1235; c) S. A. Rodgen, S. E. Schaus, *Angew. Chem. Int. Ed.* **2006**, 45, 4929; *Angew. Chem.* **2006**, 118, 5051; d) A. Vik, L.-L. Gundersen, *Tetrahedron Lett.* **2007**, 48, 1931.
- [7] Although labeled as cytotoxicity in Ref. [2], the use of GI₅₀ indicates cytostatic effect. Based on the activity of our analogues, we assume the authors observed cytotoxicity and meant to use EC₅₀.
- [8] W. Adam, O. Krebs, *Chem. Rev.* **2003**, 103, 4131.
- [9] a) G. E. Keck, R. R. Webb, *J. Org. Chem.* **1982**, 47, 1302; b) W. Adam, H.-G. Degen, O. Krebs, C. R. Saha-Möller, *J. Am. Chem. Soc.* **2002**, 124, 12938; c) W. Oppolzer, G. Poli, C. Starkemann, G. Bernardinelli, *Tetrahedron Lett.* **1988**, 29, 3559; d) G. KreBe, A. Vasella, H. Felber, A. Ritter, B. Ascherl, *Recl. Trav. Chim. Pays-Bas* **1986**, 105, 295; e) W. Adam, N. Bottke, *J. Am. Chem. Soc.* **2000**, 122, 9846; f) W. Adam, N. Bottke, O. Krebs, I. Lykakis, M. Orfanopoulos, M. Stratakis, *J. Am. Chem. Soc.* **2002**, 124, 14403.
- [10] G. E. Keck, R. R. Webb, J. B. Yates, *Tetrahedron Lett.* **1981**, 37, 4007.
- [11] P. Wipf, S. Lim, *Angew. Chem. Int. Ed. Engl.* **1993**, 32, 1068; *Angew. Chem.* **1993**, 105, 1095.
- [12] J. Chen, S. Ma, *J. Org. Chem.* **2009**, 74, 5595.
- [13] N. Okukado, E. Negishi, *Tetrahedron Lett.* **1978**, 19, 2357.
- [14] J. A. Montgomery, K. Hewson, *J. Org. Chem.* **1961**, 26, 4469.
- [15] T. Tokoroyama, *Synthesis* **2000**, 611.
- [16] T. J. Donohoe, P. M. Guyo, M. Helliwell, *Tetrahedron Lett.* **1999**, 40, 435.
- [17] R. O. Duthaler, J. D. Roberts, *J. Am. Chem. Soc.* **1978**, 100, 4969.
- [18] A. Giner-Sorolla, *J. Heterocycl. Chem.* **1970**, 7, 75.
- [19] A. G. Leach, K. N. Houk, *J. Am. Chem. Soc.* **2002**, 124, 14820.
- [20] CCDC 1036462 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [21] L.-L. Gundersen, *Phytochem. Rev.* **2013**, 12, 467.
- [22] E.-I. Negishi, Q. Hu, Z. Huang, M. Qian, G. Wang, *Aldrichimica Acta* **2005**, 38, 2005.
- [23] CCDC 1036478 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.